

Ethyl 2-(3,4-dimethoxybenzyl)-1-phenyl-sulfonyl-1*H*-indole-3-carboxylate

B. Gunasekaran,^a Radhakrishnan Sureshbabu,^b
A. K. Mohanakrishnan,^b G. Chakkaravarthi^c and
V. Manivannan^{d*}

^aDepartment of Physics, AMET University, Kanathur, Chennai 603 112, India,
^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai
600 025, India, ^cDepartment of Physics, CPCL Polytechnic College, Chennai 600
068, India, and ^dDepartment of Research and Development, PRIST University,
Vallam, Thanjavur 613 403, Tamil Nadu, India
Correspondence e-mail: manivan_1999@yahoo.com


Received 7 July 2009; accepted 7 July 2009

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å;
 R factor = 0.043; wR factor = 0.124; data-to-parameter ratio = 16.3.

In the title compound, $\text{C}_{26}\text{H}_{25}\text{NO}_6\text{S}$, the phenyl ring forms a dihedral angle of $82.5(1)^\circ$ with the indole ring system. The molecular structure is stabilized by weak intramolecular $\text{C}-\text{H}\cdots\text{O}$ interactions and the crystal structure is stabilized by weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ interactions.

Related literature

For the biological activity of indoles see: Macor *et al.* (1992); Williams *et al.* (1993); For related structures, see: Chakkaravarthi *et al.* (2007, 2008). For graph set notation see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{26}\text{H}_{25}\text{NO}_6\text{S}$
 $M_r = 479.53$
Triclinic, $P\bar{1}$
 $a = 9.2914(3)$ Å

$b = 9.3008(3)$ Å
 $c = 14.1561(5)$ Å
 $\alpha = 87.367(2)^\circ$
 $\beta = 76.158(2)^\circ$

$\gamma = 87.877(2)^\circ$
 $V = 1186.13(7)$ Å³
 $Z = 2$
Mo $K\alpha$ radiation

$\mu = 0.18$ mm⁻¹
 $T = 295$ K
 $0.24 \times 0.20 \times 0.16$ mm

Data collection

Bruker Kappa APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.958$, $T_{\max} = 0.972$

26965 measured reflections
5046 independent reflections
3632 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.124$
 $S = 1.03$
5046 reflections

310 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.18$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.29$ e Å⁻³

Table 1
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{O1}$	0.93	2.56	2.911 (3)	103
$\text{C8}-\text{H8}\cdots\text{O2}$	0.93	2.31	2.894 (3)	121
$\text{C11}-\text{H11}\cdots\text{O4}$	0.93	2.36	2.885 (2)	115
$\text{C18}-\text{H18A}\cdots\text{O1}$	0.97	2.23	2.855 (3)	122
$\text{C18}-\text{H18B}\cdots\text{O3}$	0.97	2.33	2.930 (3)	119
$\text{C25}-\text{H25B}\cdots\text{O1}^{\text{i}}$	0.96	2.38	3.231 (3)	147
$\text{C9}-\text{H9}\cdots\text{O2}^{\text{ii}}$	0.93	2.58	3.503 (3)	174

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, -y, -z$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009); software used to prepare material for publication: SHELXL97.

Author thanks AMET University management, India, for their kind support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT2993).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Chakkaravarthi, G., Dhayalan, V., Mohanakrishnan, A. K. & Manivannan, V. (2007). *Acta Cryst.* **E63**, o3698.
Chakkaravarthi, G., Sureshbabu, R., Mohanakrishnan, A. K. & Manivannan, V. (2008). *Acta Cryst.* **E64**, o732.
Macor, J. E., Fox, C. B., Johnson, C., Koe, B. K., Lebel, L. A. & Zorn, S. H. (1992). *J. Med. Chem.* **35**, 3625–3632.
Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Williams, T. M., Ciccarone, T. M., MacTough, S. C., Rooney, C. S., Balani, S. K., Condra, J. H., Emini, E. A., Goldman, M. E., Greenlee, W. J. & Kauffman, L. R. (1993). *J. Med. Chem.* **36**, 1291–1294.

supplementary materials

Acta Cryst. (2009). E65, o1856 [doi:10.1107/S1600536809026506]

Ethyl 2-(3,4-dimethoxybenzyl)-1-phenylsulfonyl-1*H*-indole-3-carboxylate

B. Gunasekaran, R. Sureshbabu, A. K. Mohanakrishnan, G. Chakkaravarthi and V. Manivannan

Comment

The chemistry of indole has been of increasing interest, since several compounds of this type possess diverse biological activities (Macor *et al.*, 1992). In addition, phenylsulfonyl indole compounds inhibit the HIV-1 RT enzyme *in vitro* and HTLVIIIb viral spread in MT-4 human T-lymphoid cells (Williams *et al.*, 1993).

The geometric parameters of the title compound, (I), (Fig. 1) agree well with the reported similar structures (Chakkaravarthi *et al.*, 2007; Chakkaravarthi *et al.*, 2008). The phenyl ring makes a dihedral angle of 82.5 (1)° with the indole ring system. The two aromatic rings C1—C6 and C19—C24 are inclined at an angle of 44.2 (1)° with respect to each other. The sum of the bond angles around N1 [358.8 (5)°] indicate the sp^2 hybridized state. The torsion angles O1—S1—N1—C14 and O2—S1—N1—C7 [27.8 (2)° and -37.1 (2)°, respectively] indicate the *syn* conformation of the sulfonyl moiety.

A distorted tetrahedral geometry [O1—S1—O2 = 120.4 (1)° and O1—S1—N1 = 106.9 (1)°] around S1 is observed. The widening of the angles may be due to repulsive interactions between the two short S=O bonds.

The molecular structure is stabilized by weak intramolecular C—H...O interactions and the crystal packing is stabilized by weak intermolecular C—H...O interactions. The C6—H6...O1 interaction generate an S(5) graph set motif and C8—H8...O2 and C11—H11...O4 interactions generate S(6) graph set motifs (Bernstein *et al.*, 1995).

Experimental

Ethyl 2-(acetoxymethyl)-1-(phenylsulfonyl)-1*H*-indole-3-carboxylate (0.39 g, 0.97 mmol) was dissolved in dry 1,2-dichloroethane (15 ml). To this, anhydrous Ferric chloride (0.02 g, 0.09 mmol) and 1,2-dimethoxy benzene (0.15 ml, 1.16 mmol) were added under nitrogen atmosphere. It was refluxed for 5 hr and cooled to room temperature. Ferric chloride was carefully filtered off and the filtrate was poured to water (50 ml) and extracted with chloroform (30 ml). The organic layer was separated and dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure to give the product. It was recrystallized from methanol. Yield: 0.28 g (61%), M.Pt: 134–136°C.

Refinement

H atoms were positioned geometrically and refined using riding model with C—H = 0.93Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic C—H, C—H = 0.97Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for CH₂, C—H = 0.96Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for CH₃.

Figures




Fig. 1. The molecular structure of the title compound, with atom labels and 30% probability displacement ellipsoids for non-H atoms.




Fig. 2. The packing of the title compound, viewed down the *a* axis. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted.

Ethyl 2-(3,4-dimethoxybenzyl)-1-phenylsulfonyl-1*H*-indole-3-carboxylate

Crystal data

$C_{26}H_{25}NO_6S$

$M_r = 479.53$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 9.2914(3) \text{ \AA}$

$b = 9.3008(3) \text{ \AA}$

$c = 14.1561(5) \text{ \AA}$

$\alpha = 87.367(2)^\circ$

$\beta = 76.158(2)^\circ$

$\gamma = 87.877(2)^\circ$

$V = 1186.13(7) \text{ \AA}^3$

$Z = 2$

$F_{000} = 504$

$D_x = 1.343 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5120 reflections

$\theta = 2.4\text{--}25.1^\circ$

$\mu = 0.18 \text{ mm}^{-1}$

$T = 295 \text{ K}$

Block, colourless

$0.24 \times 0.20 \times 0.16 \text{ mm}$

Data collection

Bruker Kappa APEX2
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295 \text{ K}$

ω and ϕ scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.958$, $T_{\max} = 0.972$

5046 independent reflections

3632 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\text{max}} = 26.8^\circ$

$\theta_{\text{min}} = 1.5^\circ$

$h = -11 \rightarrow 11$

$k = -11 \rightarrow 11$

26965 measured reflections

$l = -17 \rightarrow 17$

Refinement

Refinement on F^2

Secondary atom site location: difference Fourier map

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.043$

H-atom parameters constrained

$wR(F^2) = 0.124$

$$w = 1/[\sigma^2(F_o^2) + (0.0563P)^2 + 0.284P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$S = 1.03$

$(\Delta/\sigma)_{\max} < 0.001$

5046 reflections

$\Delta\rho_{\max} = 0.18 \text{ e } \text{\AA}^{-3}$

310 parameters

$\Delta\rho_{\min} = -0.29 \text{ e } \text{\AA}^{-3}$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.13534 (6)	-0.05077 (5)	0.22554 (4)	0.06267 (17)
O4	-0.19660 (14)	0.34677 (15)	-0.01641 (10)	0.0687 (4)
N1	0.06324 (16)	0.04353 (16)	0.14286 (11)	0.0546 (4)
O2	0.26358 (17)	-0.12263 (15)	0.17097 (12)	0.0813 (4)
O1	0.01860 (18)	-0.13039 (16)	0.28540 (12)	0.0863 (5)
C12	0.05829 (19)	0.2019 (2)	0.01735 (12)	0.0534 (4)
O5	-0.2124 (3)	0.48992 (18)	0.40802 (13)	0.1186 (7)
O3	-0.34719 (17)	0.2500 (2)	0.11472 (13)	0.1079 (6)
C24	-0.3063 (2)	0.0625 (2)	0.41242 (14)	0.0635 (5)
H24	-0.3268	-0.0350	0.4158	0.076*
C19	-0.24165 (19)	0.12692 (19)	0.32427 (13)	0.0548 (4)
C13	-0.08945 (18)	0.1877 (2)	0.07827 (13)	0.0540 (4)
C1	0.19069 (19)	0.07726 (18)	0.29540 (13)	0.0532 (4)
C14	-0.08487 (18)	0.09319 (19)	0.15310 (13)	0.0536 (4)
C2	0.3292 (2)	0.1354 (2)	0.26491 (14)	0.0611 (5)
H2	0.3928	0.1083	0.2068	0.073*
C20	-0.2098 (2)	0.2712 (2)	0.32182 (13)	0.0633 (5)
H20	-0.1652	0.3168	0.2628	0.076*
C18	-0.2119 (2)	0.0430 (2)	0.23233 (14)	0.0632 (5)
H18A	-0.1932	-0.0572	0.2489	0.076*
H18B	-0.3006	0.0477	0.2075	0.076*
C7	0.15157 (19)	0.11180 (19)	0.05862 (13)	0.0530 (4)
O6	-0.3423 (2)	0.3687 (2)	0.57147 (12)	0.1124 (7)
C8	0.3023 (2)	0.0979 (2)	0.01668 (15)	0.0659 (5)
H8	0.3635	0.0364	0.0443	0.079*
C15	-0.2254 (2)	0.2614 (2)	0.06311 (15)	0.0637 (5)
C10	0.2676 (2)	0.2679 (3)	-0.10809 (15)	0.0771 (6)
H10	0.3086	0.3210	-0.1648	0.092*
C23	-0.3416 (2)	0.1395 (3)	0.49619 (15)	0.0721 (6)

supplementary materials

H23	-0.3862	0.0937	0.5552	0.087*
C22	-0.3116 (2)	0.2824 (2)	0.49306 (14)	0.0734 (6)
C3	0.3721 (2)	0.2339 (2)	0.32159 (16)	0.0724 (6)
H3	0.4653	0.2739	0.3019	0.087*
C11	0.1178 (2)	0.2816 (2)	-0.06740 (14)	0.0677 (5)
H11	0.0577	0.3429	-0.0961	0.081*
C6	0.0959 (2)	0.1151 (3)	0.38187 (15)	0.0730 (6)
H6	0.0031	0.0744	0.4024	0.088*
C21	-0.2426 (3)	0.3481 (2)	0.40464 (15)	0.0719 (6)
C9	0.3577 (2)	0.1780 (3)	-0.06701 (16)	0.0741 (6)
H9	0.4584	0.1711	-0.0965	0.089*
C16	-0.3218 (3)	0.4260 (2)	-0.03882 (19)	0.0812 (6)
H16A	-0.3649	0.4908	0.0128	0.097*
H16B	-0.3971	0.3602	-0.0453	0.097*
C5	0.1408 (3)	0.2136 (3)	0.43708 (17)	0.0882 (7)
H5	0.0777	0.2405	0.4955	0.106*
C4	0.2778 (3)	0.2730 (3)	0.40702 (17)	0.0809 (6)
H4	0.3068	0.3404	0.4449	0.097*
C17	-0.2667 (3)	0.5085 (3)	-0.1315 (2)	0.1058 (9)
H17A	-0.1923	0.5732	-0.1241	0.159*
H17B	-0.3475	0.5627	-0.1484	0.159*
H17C	-0.2247	0.4433	-0.1820	0.159*
C25	-0.1090 (5)	0.5491 (3)	0.3324 (2)	0.170 (2)
H25A	-0.0150	0.5004	0.3284	0.255*
H25B	-0.1003	0.6493	0.3430	0.255*
H25C	-0.1390	0.5392	0.2728	0.255*
C26	-0.4361 (4)	0.3220 (4)	0.65636 (19)	0.1280 (12)
H26A	-0.5290	0.2974	0.6437	0.192*
H26B	-0.4524	0.3971	0.7024	0.192*
H26C	-0.3927	0.2388	0.6826	0.192*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0670 (3)	0.0450 (3)	0.0787 (3)	-0.0027 (2)	-0.0224 (2)	-0.0013 (2)
O4	0.0577 (8)	0.0723 (9)	0.0770 (9)	0.0095 (6)	-0.0196 (7)	-0.0045 (7)
N1	0.0505 (8)	0.0554 (8)	0.0594 (9)	-0.0013 (6)	-0.0142 (7)	-0.0106 (7)
O2	0.0858 (10)	0.0559 (8)	0.1053 (12)	0.0212 (7)	-0.0288 (9)	-0.0201 (8)
O1	0.0913 (11)	0.0628 (9)	0.1081 (12)	-0.0266 (8)	-0.0310 (9)	0.0210 (8)
C12	0.0492 (9)	0.0633 (11)	0.0486 (10)	0.0017 (8)	-0.0110 (7)	-0.0180 (8)
O5	0.183 (2)	0.0696 (10)	0.0809 (11)	-0.0411 (12)	0.0196 (12)	-0.0157 (9)
O3	0.0499 (8)	0.1632 (18)	0.0973 (12)	0.0196 (10)	0.0004 (8)	0.0213 (12)
C24	0.0580 (11)	0.0606 (11)	0.0689 (13)	-0.0168 (9)	-0.0097 (9)	0.0126 (10)
C19	0.0467 (9)	0.0563 (10)	0.0592 (11)	-0.0086 (8)	-0.0079 (8)	0.0014 (8)
C13	0.0466 (9)	0.0643 (11)	0.0515 (10)	0.0006 (8)	-0.0106 (7)	-0.0143 (9)
C1	0.0537 (10)	0.0507 (10)	0.0561 (10)	0.0013 (8)	-0.0166 (8)	0.0045 (8)
C14	0.0486 (9)	0.0563 (10)	0.0577 (10)	-0.0051 (8)	-0.0130 (8)	-0.0162 (8)
C2	0.0586 (11)	0.0645 (11)	0.0599 (11)	-0.0039 (9)	-0.0135 (9)	-0.0014 (9)

C20	0.0734 (12)	0.0599 (11)	0.0489 (10)	-0.0141 (9)	0.0015 (9)	0.0043 (8)
C18	0.0546 (10)	0.0623 (11)	0.0725 (12)	-0.0128 (9)	-0.0123 (9)	-0.0068 (9)
C7	0.0496 (9)	0.0565 (10)	0.0540 (10)	-0.0004 (8)	-0.0112 (8)	-0.0185 (8)
O6	0.1424 (17)	0.1171 (14)	0.0607 (10)	-0.0368 (12)	0.0179 (10)	-0.0204 (9)
C8	0.0511 (10)	0.0770 (13)	0.0694 (13)	0.0075 (9)	-0.0123 (9)	-0.0190 (10)
C15	0.0533 (11)	0.0792 (13)	0.0597 (12)	0.0057 (9)	-0.0148 (9)	-0.0144 (10)
C10	0.0677 (13)	0.1025 (17)	0.0533 (11)	-0.0056 (12)	0.0022 (10)	-0.0074 (11)
C23	0.0682 (12)	0.0859 (15)	0.0555 (12)	-0.0210 (11)	-0.0023 (9)	0.0174 (11)
C22	0.0777 (14)	0.0842 (15)	0.0511 (11)	-0.0149 (11)	0.0008 (10)	-0.0037 (10)
C3	0.0688 (13)	0.0759 (13)	0.0768 (14)	-0.0156 (10)	-0.0245 (11)	0.0005 (11)
C11	0.0611 (11)	0.0893 (15)	0.0502 (11)	0.0036 (10)	-0.0086 (9)	-0.0067 (10)
C6	0.0588 (11)	0.0993 (16)	0.0610 (12)	-0.0080 (11)	-0.0135 (9)	-0.0048 (11)
C21	0.0886 (15)	0.0598 (12)	0.0582 (12)	-0.0186 (10)	0.0031 (10)	-0.0015 (9)
C9	0.0508 (10)	0.0985 (16)	0.0684 (13)	0.0021 (11)	-0.0018 (10)	-0.0234 (12)
C16	0.0718 (13)	0.0725 (14)	0.1077 (18)	0.0176 (11)	-0.0386 (13)	-0.0152 (13)
C5	0.0776 (15)	0.124 (2)	0.0647 (13)	0.0054 (14)	-0.0168 (11)	-0.0265 (13)
C4	0.0899 (16)	0.0852 (15)	0.0768 (15)	-0.0024 (13)	-0.0356 (13)	-0.0175 (12)
C17	0.122 (2)	0.0814 (17)	0.124 (2)	0.0123 (16)	-0.0544 (19)	0.0124 (16)
C25	0.293 (5)	0.085 (2)	0.100 (2)	-0.092 (3)	0.029 (3)	0.0008 (17)
C26	0.153 (3)	0.151 (3)	0.0598 (15)	-0.021 (2)	0.0180 (17)	-0.0110 (16)

Geometric parameters (Å, °)

S1—O1	1.4160 (15)	O6—C26	1.368 (3)
S1—O2	1.4182 (15)	O6—C22	1.368 (3)
S1—N1	1.6809 (16)	C8—C9	1.371 (3)
S1—C1	1.7485 (18)	C8—H8	0.9300
O4—C15	1.326 (2)	C10—C9	1.369 (3)
O4—C16	1.445 (2)	C10—C11	1.377 (3)
N1—C14	1.411 (2)	C10—H10	0.9300
N1—C7	1.414 (2)	C23—C22	1.365 (3)
C12—C11	1.387 (3)	C23—H23	0.9300
C12—C7	1.392 (3)	C22—C21	1.388 (3)
C12—C13	1.443 (2)	C3—C4	1.369 (3)
O5—C21	1.363 (3)	C3—H3	0.9300
O5—C25	1.365 (3)	C11—H11	0.9300
O3—C15	1.196 (2)	C6—C5	1.369 (3)
C24—C19	1.370 (2)	C6—H6	0.9300
C24—C23	1.379 (3)	C9—H9	0.9300
C24—H24	0.9300	C16—C17	1.479 (4)
C19—C20	1.382 (2)	C16—H16A	0.9700
C19—C18	1.511 (3)	C16—H16B	0.9700
C13—C14	1.353 (3)	C5—C4	1.370 (3)
C13—C15	1.471 (3)	C5—H5	0.9300
C1—C6	1.378 (3)	C4—H4	0.9300
C1—C2	1.378 (3)	C17—H17A	0.9600
C14—C18	1.492 (3)	C17—H17B	0.9600
C2—C3	1.375 (3)	C17—H17C	0.9600
C2—H2	0.9300	C25—H25A	0.9600

supplementary materials

C20—C21	1.367 (3)	C25—H25B	0.9600
C20—H20	0.9300	C25—H25C	0.9600
C18—H18A	0.9700	C26—H26A	0.9600
C18—H18B	0.9700	C26—H26B	0.9600
C7—C8	1.388 (2)	C26—H26C	0.9600
O1—S1—O2	120.38 (10)	C11—C10—H10	119.2
O1—S1—N1	106.84 (9)	C22—C23—C24	120.41 (18)
O2—S1—N1	105.33 (9)	C22—C23—H23	119.8
O1—S1—C1	108.93 (9)	C24—C23—H23	119.8
O2—S1—C1	108.65 (9)	C23—C22—O6	125.20 (19)
N1—S1—C1	105.74 (8)	C23—C22—C21	118.95 (19)
C15—O4—C16	116.22 (16)	O6—C22—C21	115.85 (19)
C14—N1—C7	108.32 (15)	C4—C3—C2	120.0 (2)
C14—N1—S1	127.51 (13)	C4—C3—H3	120.0
C7—N1—S1	122.95 (12)	C2—C3—H3	120.0
C11—C12—C7	119.17 (17)	C10—C11—C12	118.6 (2)
C11—C12—C13	133.86 (18)	C10—C11—H11	120.7
C7—C12—C13	106.97 (16)	C12—C11—H11	120.7
C21—O5—C25	117.9 (2)	C5—C6—C1	118.8 (2)
C19—C24—C23	121.22 (18)	C5—C6—H6	120.6
C19—C24—H24	119.4	C1—C6—H6	120.6
C23—C24—H24	119.4	O5—C21—C20	124.45 (18)
C24—C19—C20	118.07 (17)	O5—C21—C22	115.43 (18)
C24—C19—C18	120.49 (17)	C20—C21—C22	120.12 (19)
C20—C19—C18	121.41 (16)	C10—C9—C8	121.29 (19)
C14—C13—C12	109.01 (16)	C10—C9—H9	119.4
C14—C13—C15	124.31 (16)	C8—C9—H9	119.4
C12—C13—C15	126.68 (17)	O4—C16—C17	107.4 (2)
C6—C1—C2	121.12 (18)	O4—C16—H16A	110.2
C6—C1—S1	119.04 (15)	C17—C16—H16A	110.2
C2—C1—S1	119.81 (14)	O4—C16—H16B	110.2
C13—C14—N1	108.19 (15)	C17—C16—H16B	110.2
C13—C14—C18	127.54 (17)	H16A—C16—H16B	108.5
N1—C14—C18	124.23 (17)	C6—C5—C4	120.5 (2)
C3—C2—C1	119.09 (19)	C6—C5—H5	119.7
C3—C2—H2	120.5	C4—C5—H5	119.7
C1—C2—H2	120.5	C3—C4—C5	120.4 (2)
C21—C20—C19	121.20 (17)	C3—C4—H4	119.8
C21—C20—H20	119.4	C5—C4—H4	119.8
C19—C20—H20	119.4	C16—C17—H17A	109.5
C14—C18—C19	115.27 (15)	C16—C17—H17B	109.5
C14—C18—H18A	108.5	H17A—C17—H17B	109.5
C19—C18—H18A	108.5	C16—C17—H17C	109.5
C14—C18—H18B	108.5	H17A—C17—H17C	109.5
C19—C18—H18B	108.5	H17B—C17—H17C	109.5
H18A—C18—H18B	107.5	O5—C25—H25A	109.5
C8—C7—C12	121.86 (18)	O5—C25—H25B	109.5
C8—C7—N1	130.62 (18)	H25A—C25—H25B	109.5
C12—C7—N1	107.51 (15)	O5—C25—H25C	109.5

C26—O6—C22	119.5 (2)	H25A—C25—H25C	109.5
C9—C8—C7	117.6 (2)	H25B—C25—H25C	109.5
C9—C8—H8	121.2	O6—C26—H26A	109.5
C7—C8—H8	121.2	O6—C26—H26B	109.5
O3—C15—O4	122.87 (19)	H26A—C26—H26B	109.5
O3—C15—C13	126.1 (2)	O6—C26—H26C	109.5
O4—C15—C13	111.03 (16)	H26A—C26—H26C	109.5
C9—C10—C11	121.5 (2)	H26B—C26—H26C	109.5
C9—C10—H10	119.2		
O1—S1—N1—C14	27.85 (17)	C14—N1—C7—C8	-178.86 (17)
O2—S1—N1—C14	156.95 (14)	S1—N1—C7—C8	12.9 (3)
C1—S1—N1—C14	-88.10 (15)	C14—N1—C7—C12	-0.29 (18)
O1—S1—N1—C7	-166.29 (13)	S1—N1—C7—C12	-168.52 (11)
O2—S1—N1—C7	-37.18 (15)	C12—C7—C8—C9	0.8 (3)
C1—S1—N1—C7	77.77 (14)	N1—C7—C8—C9	179.23 (17)
C23—C24—C19—C20	1.3 (3)	C16—O4—C15—O3	-0.6 (3)
C23—C24—C19—C18	-176.42 (18)	C16—O4—C15—C13	-179.67 (16)
C11—C12—C13—C14	179.75 (19)	C14—C13—C15—O3	1.4 (3)
C7—C12—C13—C14	0.14 (19)	C12—C13—C15—O3	-179.1 (2)
C11—C12—C13—C15	0.2 (3)	C14—C13—C15—O4	-179.51 (16)
C7—C12—C13—C15	-179.39 (16)	C12—C13—C15—O4	-0.1 (3)
O1—S1—C1—C6	-19.05 (18)	C19—C24—C23—C22	-0.5 (3)
O2—S1—C1—C6	-151.89 (16)	C24—C23—C22—O6	179.8 (2)
N1—S1—C1—C6	95.47 (16)	C24—C23—C22—C21	-1.2 (3)
O1—S1—C1—C2	159.04 (15)	C26—O6—C22—C23	-14.4 (4)
O2—S1—C1—C2	26.20 (17)	C26—O6—C22—C21	166.6 (3)
N1—S1—C1—C2	-86.45 (16)	C1—C2—C3—C4	0.0 (3)
C12—C13—C14—N1	-0.32 (19)	C9—C10—C11—C12	0.2 (3)
C15—C13—C14—N1	179.22 (16)	C7—C12—C11—C10	0.3 (3)
C12—C13—C14—C18	-178.34 (16)	C13—C12—C11—C10	-179.24 (19)
C15—C13—C14—C18	1.2 (3)	C2—C1—C6—C5	0.9 (3)
C7—N1—C14—C13	0.38 (18)	S1—C1—C6—C5	178.94 (18)
S1—N1—C14—C13	167.92 (12)	C25—O5—C21—C20	-18.9 (5)
C7—N1—C14—C18	178.48 (15)	C25—O5—C21—C22	161.3 (3)
S1—N1—C14—C18	-14.0 (2)	C19—C20—C21—O5	179.0 (2)
C6—C1—C2—C3	-0.7 (3)	C19—C20—C21—C22	-1.2 (3)
S1—C1—C2—C3	-178.79 (15)	C23—C22—C21—O5	-178.2 (2)
C24—C19—C20—C21	-0.5 (3)	O6—C22—C21—O5	0.9 (3)
C18—C19—C20—C21	177.3 (2)	C23—C22—C21—C20	2.0 (4)
C13—C14—C18—C19	-93.4 (2)	O6—C22—C21—C20	-178.9 (2)
N1—C14—C18—C19	88.8 (2)	C11—C10—C9—C8	-0.2 (3)
C24—C19—C18—C14	-152.33 (18)	C7—C8—C9—C10	-0.3 (3)
C20—C19—C18—C14	30.0 (3)	C15—O4—C16—C17	-177.62 (19)
C11—C12—C7—C8	-0.9 (3)	C1—C6—C5—C4	-0.3 (4)
C13—C12—C7—C8	178.82 (16)	C2—C3—C4—C5	0.6 (4)
C11—C12—C7—N1	-179.58 (15)	C6—C5—C4—C3	-0.4 (4)
C13—C12—C7—N1	0.10 (18)		

supplementary materials

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C6—H6 \cdots O1	0.93	2.56	2.911 (3)	103
C8—H8 \cdots O2	0.93	2.31	2.894 (3)	121
C11—H11 \cdots O4	0.93	2.36	2.885 (2)	115
C18—H18A \cdots O1	0.97	2.23	2.855 (3)	122
C18—H18B \cdots O3	0.97	2.33	2.930 (3)	119
C25—H25B \cdots O1 ⁱ	0.96	2.38	3.231 (3)	147
C9—H9 \cdots O2 ⁱⁱ	0.93	2.58	3.503 (3)	174

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y, -z$.

Fig. 1




Fig. 2

