

2-(1*H*-Benzotriazol-1-yl)-1-(3-methylbenzoyl)ethyl benzoate

Wu-Lan Zeng,^a Lian-Cai Du,^b Lei Zhang^b and Fang-Fang Jian^{b*}

^aMicroScale Science Institute, Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and ^bMicroscale Science Institute, Department of Biological Engineering, Weifang University, Weifang 261061, People's Republic of China

Correspondence e-mail: wulanpeng@163.com


Received 7 June 2009; accepted 9 July 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.044; wR factor = 0.110; data-to-parameter ratio = 12.6.

In the title molecule, $\text{C}_{23}\text{H}_{19}\text{N}_3\text{O}_3$, the dihedral angles between the mean plane of the benzotriazole ring system and the benzene and phenyl rings are $9.67(9)$ and $86.08(10)^\circ$, respectively. The dihedral angle between the benzene and phenyl rings is $85.89(11)^\circ$. In the crystal structure, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link molecules into chains along [010].

Related literature

For the pharmacological activities of 1H -benzotriazoles and their derivatives, see: Chen & Wu (2005). For standard bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

$\text{C}_{23}\text{H}_{19}\text{N}_3\text{O}_3$
 $M_r = 385.41$
 Monoclinic, $P2_1/c$
 $a = 10.1095(5)\text{ \AA}$
 $b = 9.3849(4)\text{ \AA}$
 $c = 20.7091(10)\text{ \AA}$
 $\beta = 99.061(4)^\circ$

$V = 1940.29(16)\text{ \AA}^3$
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.30 \times 0.10 \times 0.10\text{ mm}$

Data collection

Siemens SMART CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.974$, $T_{\max} = 0.991$

16102 measured reflections
 3301 independent reflections
 2071 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.110$
 $S = 1.00$
 3301 reflections

262 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.19\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.22\text{ e \AA}^{-3}$

Table 1
 Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$C7-\text{H7B}\cdots\text{O2}^{\text{i}}$	0.97	2.42	3.062 (2)	123
$C11-\text{H11A}\cdots\text{O1}^{\text{ii}}$	0.93	2.60	3.375 (2)	141

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2841).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Chen, Z.-Y. & Wu, M.-J. (2005). *Org. Lett.* **7**, 475–477.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o1882 [doi:10.1107/S1600536809026853]

2-(1*H*-Benzotriazol-1-yl)-1-(3-methylbenzoyl)ethyl benzoate

W.-L. Zeng, L.-C. Du, L. Zhang and F.-F. Jian

Comment

1*H*-Benzotriazoles and their derivatives are an important class of compounds because they exhibit a broad spectrum of pharmacological activities such as antifungal, antitumor and antineoplastic activities (Chen & Wu., 2005). Herein, we present the crystal structure of the title compound (I). In (I) (Fig. 1) all bond lengths (Allen *et al.*, 1987) and angles within normal ranges. The benzotriazole ring system is essentially planar. The dihedral angles between the mean plane of the benzotriazole ring system and rings C10—C15 and C18—C23 are 9.67 (9) and 86.08 (10) $^{\circ}$, respectively. The dihedral angle between rings C10—C15 and C19—C23 is 85.89 (11) $^{\circ}$. In the crystal structure weak intermolecular C—H \cdots O hydrogen bonds link molecules into chains along [010] (see Fig. 2).

Experimental

Bromine (3.2 g, 0.02 mol) was added dropwise to a solution of 3-(1*H*-benzo[*d*][1,2,3]triazol-1-yl)-1-*m*-tolylpropan-1-one (5.30 g, 0.02 mol) and sodium acetate (1.6 g, 0.02 mol) in acetic acid (50 ml). The reaction proceeded for 8 h. Water (50 ml) and chloroform (20 ml) were then added. The organic layer was washed successively with saturated sodium bicarbonate solution and brine, dried over anhydrous magnesium sulfate and the chloroform solution filtered. It was cooled with ice-water, and then an acetone solution (10 ml) of benzoic acid (2.24 g, 0.02 mol) and triethylamine (2.8 ml) was added. The mixture was stirred with ice-water for about 6 h. The solution was then filtered and concentrated. Single crystals were obtained by slow evaporation of an acetone-ethylacetate (1:1 *v/v*) solution of (I) at room temperature over a period of one week.

Refinement

All H atoms were located in difference Fourier maps and were subsequently constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.98 Å, and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $1.5 U_{\text{eq}}(\text{methyl C})$ H atoms.

Figures




Fig. 1. The molecular structure of (I), drawn with 30% probability ellipsoids.

supplementary materials




Fig. 2. Part of the crystal structure with hydrogen bonds shown as dashed lines.

2-(1*H*-Benzotriazol-1-yl)-1-(3-methylbenzoyl)ethyl benzoate

Crystal data

C ₂₃ H ₁₉ N ₃ O ₃	$F_{000} = 808$
$M_r = 385.41$	$D_x = 1.319 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3301 reflections
$a = 10.1095 (5) \text{ \AA}$	$\theta = 2.0\text{--}25.0^\circ$
$b = 9.3849 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 20.7091 (10) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 99.061 (4)^\circ$	Block, colorless
$V = 1940.29 (16) \text{ \AA}^3$	$0.30 \times 0.10 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Siemens SMART CCD diffractometer	3301 independent reflections
Radiation source: fine-focus sealed tube	2071 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.051$
$T = 298 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -11 \rightarrow 11$
$T_{\text{min}} = 0.974$, $T_{\text{max}} = 0.991$	$k = -11 \rightarrow 11$
16102 measured reflections	$l = -23 \rightarrow 24$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0483P)^2 + 0.2024P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
3301 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
262 parameters	$\Delta\rho_{\text{max}} = 0.19 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct Extinction correction: none
methods

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.09785 (14)	0.45987 (15)	0.21346 (7)	0.0543 (4)
O2	0.17138 (13)	0.14124 (15)	0.23494 (7)	0.0508 (4)
O3	0.18369 (12)	0.29378 (14)	0.31866 (6)	0.0444 (4)
C1	0.0844 (2)	0.2945 (3)	0.51412 (11)	0.0562 (6)
C2	0.1674 (3)	0.2855 (3)	0.57479 (12)	0.0757 (8)
H2B	0.1432	0.2312	0.6086	0.091*
C3	0.2831 (3)	0.3582 (4)	0.58230 (14)	0.0886 (10)
H3A	0.3400	0.3535	0.6222	0.106*
C4	0.3204 (3)	0.4402 (3)	0.53236 (16)	0.0884 (9)
H4A	0.4022	0.4876	0.5397	0.106*
C5	0.2409 (3)	0.4538 (3)	0.47244 (13)	0.0677 (7)
H5A	0.2654	0.5103	0.4394	0.081*
C6	0.1216 (2)	0.3781 (2)	0.46443 (10)	0.0464 (5)
C7	-0.0004 (2)	0.4194 (2)	0.34864 (9)	0.0518 (6)
H7A	0.0510	0.5066	0.3485	0.062*
H7B	-0.0943	0.4431	0.3358	0.062*
C8	0.04164 (19)	0.3168 (2)	0.29917 (9)	0.0431 (5)
H8A	-0.0068	0.2265	0.3000	0.052*
C9	0.0140 (2)	0.3818 (2)	0.23043 (9)	0.0402 (5)
C10	-0.11527 (19)	0.3540 (2)	0.18760 (9)	0.0391 (5)
C11	-0.2093 (2)	0.2566 (2)	0.20324 (11)	0.0521 (6)
H11A	-0.1922	0.2043	0.2418	0.062*
C12	-0.3285 (2)	0.2383 (3)	0.16091 (12)	0.0634 (7)
H12A	-0.3912	0.1729	0.1711	0.076*
C13	-0.3551 (2)	0.3158 (2)	0.10391 (11)	0.0586 (6)
H13A	-0.4359	0.3025	0.0762	0.070*
C14	-0.2632 (2)	0.4134 (2)	0.08716 (10)	0.0479 (5)
C15	-0.1438 (2)	0.4296 (2)	0.12935 (10)	0.0442 (5)
H15A	-0.0804	0.4932	0.1184	0.053*
C16	-0.2919 (3)	0.4998 (3)	0.02543 (11)	0.0743 (8)
H16A	-0.3786	0.4747	0.0022	0.112*

supplementary materials

H16B	-0.2250	0.4804	-0.0015	0.112*
H16C	-0.2905	0.5993	0.0362	0.112*
C17	0.2376 (2)	0.1978 (2)	0.28153 (10)	0.0395 (5)
C18	0.38089 (19)	0.1709 (2)	0.30448 (9)	0.0412 (5)
C19	0.4567 (2)	0.2509 (2)	0.35264 (11)	0.0599 (7)
H19A	0.4178	0.3263	0.3720	0.072*
C20	0.5906 (2)	0.2187 (3)	0.37213 (12)	0.0713 (8)
H20A	0.6414	0.2733	0.4044	0.086*
C21	0.6490 (2)	0.1080 (3)	0.34462 (12)	0.0637 (7)
H21A	0.7390	0.0871	0.3581	0.076*
C22	0.5745 (2)	0.0282 (3)	0.29719 (12)	0.0604 (6)
H22A	0.6140	-0.0479	0.2787	0.072*
C23	0.4409 (2)	0.0590 (2)	0.27624 (11)	0.0514 (6)
H23A	0.3915	0.0049	0.2433	0.062*
N1	-0.0359 (2)	0.2318 (2)	0.49227 (10)	0.0793 (7)
N2	-0.07420 (19)	0.2723 (2)	0.43222 (10)	0.0730 (6)
N3	0.01911 (17)	0.36109 (19)	0.41419 (8)	0.0487 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0460 (9)	0.0623 (10)	0.0531 (10)	-0.0116 (8)	0.0031 (7)	0.0094 (8)
O2	0.0483 (9)	0.0567 (9)	0.0447 (9)	-0.0043 (7)	-0.0009 (7)	-0.0106 (8)
O3	0.0398 (8)	0.0517 (9)	0.0394 (8)	0.0079 (7)	-0.0005 (6)	-0.0068 (7)
C1	0.0569 (15)	0.0742 (17)	0.0380 (14)	0.0125 (13)	0.0090 (12)	0.0009 (12)
C2	0.085 (2)	0.102 (2)	0.0395 (15)	0.0252 (18)	0.0090 (14)	0.0056 (14)
C3	0.078 (2)	0.136 (3)	0.0470 (18)	0.021 (2)	-0.0036 (16)	-0.0203 (18)
C4	0.0663 (18)	0.119 (3)	0.076 (2)	-0.0130 (17)	-0.0001 (17)	-0.040 (2)
C5	0.0687 (17)	0.0741 (17)	0.0603 (17)	-0.0128 (14)	0.0101 (14)	-0.0133 (14)
C6	0.0483 (13)	0.0536 (14)	0.0371 (13)	0.0106 (11)	0.0059 (11)	-0.0030 (11)
C7	0.0539 (14)	0.0633 (15)	0.0375 (13)	0.0198 (11)	0.0047 (11)	0.0069 (11)
C8	0.0360 (12)	0.0521 (13)	0.0397 (13)	0.0054 (10)	0.0008 (9)	0.0040 (10)
C9	0.0402 (12)	0.0422 (12)	0.0382 (12)	0.0050 (10)	0.0055 (10)	-0.0008 (10)
C10	0.0384 (12)	0.0415 (12)	0.0371 (12)	0.0015 (9)	0.0046 (9)	-0.0018 (10)
C11	0.0444 (13)	0.0583 (14)	0.0517 (14)	-0.0003 (11)	0.0021 (11)	0.0108 (11)
C12	0.0461 (14)	0.0701 (17)	0.0704 (18)	-0.0134 (12)	-0.0017 (12)	0.0155 (14)
C13	0.0472 (14)	0.0683 (16)	0.0547 (15)	-0.0041 (12)	-0.0090 (11)	-0.0019 (13)
C14	0.0500 (14)	0.0533 (14)	0.0389 (13)	0.0071 (11)	0.0021 (11)	-0.0007 (10)
C15	0.0403 (12)	0.0511 (14)	0.0405 (13)	-0.0014 (10)	0.0039 (10)	0.0014 (10)
C16	0.0744 (18)	0.092 (2)	0.0504 (15)	0.0029 (14)	-0.0092 (13)	0.0170 (14)
C17	0.0450 (13)	0.0398 (12)	0.0332 (12)	-0.0006 (10)	0.0045 (10)	0.0024 (10)
C18	0.0416 (12)	0.0432 (12)	0.0391 (12)	0.0001 (10)	0.0072 (10)	0.0032 (10)
C19	0.0476 (14)	0.0673 (16)	0.0611 (16)	0.0089 (12)	-0.0033 (12)	-0.0190 (13)
C20	0.0485 (15)	0.0864 (19)	0.0734 (18)	0.0059 (14)	-0.0077 (13)	-0.0231 (15)
C21	0.0421 (14)	0.0818 (18)	0.0669 (17)	0.0119 (13)	0.0079 (13)	0.0018 (15)
C22	0.0508 (15)	0.0639 (16)	0.0683 (17)	0.0132 (12)	0.0151 (13)	-0.0050 (13)
C23	0.0490 (14)	0.0518 (14)	0.0538 (14)	0.0006 (11)	0.0094 (11)	-0.0057 (11)
N1	0.0728 (15)	0.1137 (19)	0.0527 (14)	-0.0127 (14)	0.0142 (12)	0.0184 (13)

N2	0.0516 (12)	0.1129 (18)	0.0558 (14)	-0.0102 (12)	0.0118 (10)	0.0085 (13)
N3	0.0431 (11)	0.0660 (13)	0.0367 (11)	0.0073 (9)	0.0057 (9)	0.0051 (9)

Geometric parameters (\AA , $^{\circ}$)

O1—C9	1.213 (2)	C11—H11A	0.9300
O2—C17	1.207 (2)	C12—C13	1.377 (3)
O3—C17	1.353 (2)	C12—H12A	0.9300
O3—C8	1.445 (2)	C13—C14	1.387 (3)
C1—N1	1.363 (3)	C13—H13A	0.9300
C1—C6	1.392 (3)	C14—C15	1.382 (3)
C1—C2	1.399 (3)	C14—C16	1.503 (3)
C2—C3	1.342 (4)	C15—H15A	0.9300
C2—H2B	0.9300	C16—H16A	0.9600
C3—C4	1.388 (4)	C16—H16B	0.9600
C3—H3A	0.9300	C16—H16C	0.9600
C4—C5	1.374 (4)	C17—C18	1.474 (3)
C4—H4A	0.9300	C18—C19	1.380 (3)
C5—C6	1.387 (3)	C18—C23	1.387 (3)
C5—H5A	0.9300	C19—C20	1.384 (3)
C6—N3	1.357 (2)	C19—H19A	0.9300
C7—N3	1.448 (2)	C20—C21	1.363 (3)
C7—C8	1.515 (3)	C20—H20A	0.9300
C7—H7A	0.9700	C21—C22	1.364 (3)
C7—H7B	0.9700	C21—H21A	0.9300
C8—C9	1.534 (3)	C22—C23	1.382 (3)
C8—H8A	0.9800	C22—H22A	0.9300
C9—C10	1.482 (3)	C23—H23A	0.9300
C10—C15	1.390 (3)	N1—N2	1.300 (3)
C10—C11	1.393 (3)	N2—N3	1.355 (2)
C11—C12	1.385 (3)		
?...?	?		
C17—O3—C8	114.33 (15)	C11—C12—H12A	119.7
N1—C1—C6	109.2 (2)	C12—C13—C14	121.0 (2)
N1—C1—C2	130.5 (2)	C12—C13—H13A	119.5
C6—C1—C2	120.3 (2)	C14—C13—H13A	119.5
C3—C2—C1	117.5 (3)	C15—C14—C13	117.8 (2)
C3—C2—H2B	121.3	C15—C14—C16	120.7 (2)
C1—C2—H2B	121.3	C13—C14—C16	121.4 (2)
C2—C3—C4	122.0 (3)	C14—C15—C10	122.27 (19)
C2—C3—H3A	119.0	C14—C15—H15A	118.9
C4—C3—H3A	119.0	C10—C15—H15A	118.9
C5—C4—C3	122.4 (3)	C14—C16—H16A	109.5
C5—C4—H4A	118.8	C14—C16—H16B	109.5
C3—C4—H4A	118.8	H16A—C16—H16B	109.5
C4—C5—C6	115.8 (2)	C14—C16—H16C	109.5
C4—C5—H5A	122.1	H16A—C16—H16C	109.5
C6—C5—H5A	122.1	H16B—C16—H16C	109.5
N3—C6—C5	134.1 (2)	O2—C17—O3	121.65 (18)

supplementary materials


N3—C6—C1	103.80 (19)	O2—C17—C18	125.15 (19)
C5—C6—C1	122.1 (2)	O3—C17—C18	113.20 (17)
N3—C7—C8	112.51 (16)	C19—C18—C23	119.12 (19)
N3—C7—H7A	109.1	C19—C18—C17	123.06 (19)
C8—C7—H7A	109.1	C23—C18—C17	117.82 (18)
N3—C7—H7B	109.1	C18—C19—C20	119.9 (2)
C8—C7—H7B	109.1	C18—C19—H19A	120.1
H7A—C7—H7B	107.8	C20—C19—H19A	120.1
O3—C8—C7	106.18 (16)	C21—C20—C19	120.8 (2)
O3—C8—C9	110.28 (15)	C21—C20—H20A	119.6
C7—C8—C9	110.21 (16)	C19—C20—H20A	119.6
O3—C8—H8A	110.0	C22—C21—C20	119.5 (2)
C7—C8—H8A	110.0	C22—C21—H21A	120.2
C9—C8—H8A	110.0	C20—C21—H21A	120.2
O1—C9—C10	121.57 (18)	C21—C22—C23	120.9 (2)
O1—C9—C8	118.38 (18)	C21—C22—H22A	119.6
C10—C9—C8	119.99 (18)	C23—C22—H22A	119.6
C15—C10—C11	118.72 (19)	C22—C23—C18	119.8 (2)
C15—C10—C9	118.17 (18)	C22—C23—H23A	120.1
C11—C10—C9	123.11 (18)	C18—C23—H23A	120.1
C12—C11—C10	119.5 (2)	N2—N1—C1	107.9 (2)
C12—C11—H11A	120.3	N1—N2—N3	109.15 (19)
C10—C11—H11A	120.3	C6—N3—N2	109.98 (17)
C13—C12—C11	120.7 (2)	C6—N3—C7	130.49 (19)
C13—C12—H12A	119.7	N2—N3—C7	119.52 (18)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C7—H7B \cdots O2 ⁱ	0.97	2.42	3.062 (2)	123
C11—H11A \cdots O1 ⁱⁱ	0.93	2.60	3.375 (2)	141

Symmetry codes: (i) $-x, y+1/2, -z+1/2$; (ii) $-x, y-1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

