

4-[(*1E,3E,5E*)-6-(4-Pyridyl)hexa-1,3,5-trienyl]pyridine

Mamoun M. Bader

Department of Chemistry, Pennsylvania State University at Hazleton, 76 University Drive, Hazleton, PA 18202, USA
Correspondence e-mail: mmb11@psu.edu


Received 19 July 2009; accepted 22 July 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.032; wR factor = 0.089; data-to-parameter ratio = 10.3.

The two independent molecules in the asymmetric unit of the title compound, $C_{16}H_{14}N_2$, are planar [dihedral angle between the terminal pyridine rings = $1.76(2)^\circ$] and each display an all-*trans* configuration of $\text{C}=\text{C}$ double bonds. One of the two molecules lies about a center of inversion. The dihedral angle between the two pyridine rings in the molecule lying on a general position is $1.65(2)^\circ$.


Related literature

For acceptor-terminated polyenes, see: Gao *et al.* (2004). For the synthesis, see: Woitellier *et al.* (1989). For a related structure, see: Pham (2009).



Experimental

Crystal data



Monoclinic, $P2_1/n$
 $a = 5.837(1)\text{ \AA}$
 $b = 17.171(4)\text{ \AA}$
 $c = 19.227(4)\text{ \AA}$
 $\beta = 97.685(4)^\circ$
 $V = 1909.8(7)\text{ \AA}^3$

$Z = 6$
Mo $K\alpha$ radiation
 $\mu = 0.07\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.44 \times 0.24 \times 0.22\text{ mm}$

Data collection

Bruker SMART Platform CCD diffractometer
Absorption correction: none
18771 measured reflections

3366 independent reflections
2460 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.089$
 $S = 1.01$
3366 reflections

328 parameters
All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.09\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.15\text{ e \AA}^{-3}$

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SIR2002 (Burla *et al.*, 2003); program(s) used to refine structure: SHEXL97 (Sheldrick, 2008); molecular graphics: SHEXTL (Sheldrick, 2008); software used to prepare material for publication: SHEXL97.

This work was supported in part by Research Development grants from the Pennsylvania State University and partially by the MRSEC Program of the National Science Foundation under award No. DMR-0819885. The author also acknowledges William W. Brennessel, Victor G. Young Jr and the X-ray Crystallographic Laboratory at the University of Minnesota.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2615).

References

- Bruker (2001). SAINT and SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
Gao, X., Friscic, T. & MacGillivray, L. R. (2004). *Angew. Chem. Int. Ed.* **43**, 232–236.
Pham, P.-T. T. (2009). *Acta Cryst. E65*, o1806.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Woitellier, S., Launay, J. P. & Spangler, C. W. (1989). *Inorg. Chem.* **28**, 758–762.

supplementary materials

Acta Cryst. (2009). E65, o2006 [doi:10.1107/S1600536809029092]

4-[(1E,3E,5E)-6-(4-Pyridyl)hexa-1,3,5-trienyl]pyridine

M. M. Bader

Comment

Pyridyl-terminated polyenes have been investigated for studying electron reactions with applications in biology, inorganic reaction mechanisms and molecular electronics. They have also been used in the synthesis of coordination polymers as well as template for solid state reactions.

For related systems of acceptor terminated polyenes, see: Gao *et al.* (2004); Pham (2009). For literature related to the synthesis, see: Woitellier *et al.* (1989).

Experimental

Synthesis was carried out following literature procedures (Woitellier *et al.*, 1989) as follows: a solution of potassium *tert*-butoxide (1.2.1 g in 200 ml of glyme) was added dropwise to a solution of tetraethyl ((*E*)-2-butene-1,4-diyl)diphosphonate (16.4 g, 0.05 mol) and pyridine-4-carboxaldehyde (10.70 g, 0.10 mol) in 100 ml of glyme at room temperature. After the addition of the base was complete, the resulting mixture was stirred at room temperature for 16 h. Cold water (*ca* 300–400 ml) was added and the product isolated by vacuum filtration (3.5 g, 30% crude yield) and recrystallized from acetone (2.3 g, 19%): mp 196–197. Crystals were grown from DMF.

Refinement

All hydrogen atoms were found from the difference map and refined with individual isotropic displacement parameters.

Figures



Fig. 1. Ellipsoid plot



Fig. 2. 4-((1E,3E,5E)-6-(pyridin-4-yl)hexa-1,3,5-trienyl)pyridine.




Fig. 3. Crystal packing viewed along the *a* axis.

supplementary materials

4-[(1*E*,3*E*,5*E*)-6-(4-Pyridyl)hexa-1,3,5-trienyl]pyridine

Crystal data

C ₁₆ H ₁₄ N ₂	$F_{000} = 744$
$M_r = 234.29$	$D_x = 1.222 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 867 reflections
$a = 5.837 (1) \text{ \AA}$	$\theta = 2.5\text{--}27.5^\circ$
$b = 17.171 (4) \text{ \AA}$	$\mu = 0.07 \text{ mm}^{-1}$
$c = 19.227 (4) \text{ \AA}$	$T = 173 \text{ K}$
$\beta = 97.685 (4)^\circ$	Block, yellow
$V = 1909.8 (7) \text{ \AA}^3$	$0.44 \times 0.24 \times 0.22 \text{ mm}$
$Z = 6$	

Data collection

Bruker SMART Platform CCD diffractometer	2460 reflections with $I > 2\sigma(I)$
Radiation source: normal-focus sealed tube	$R_{\text{int}} = 0.028$
Monochromator: graphite	$\theta_{\max} = 25.0^\circ$
$T = 173 \text{ K}$	$\theta_{\min} = 1.6^\circ$
ω scans	$h = -6 \rightarrow 6$
Absorption correction: None	$k = -20 \rightarrow 20$
18771 measured reflections	$l = -22 \rightarrow 22$
3366 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.032$	All H-atom parameters refined
$wR(F^2) = 0.089$	$w = 1/[\sigma^2(F_o^2) + (0.048P)^2 + 0.1535P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.01$	$(\Delta/\sigma)_{\max} = 0.001$
3366 reflections	$\Delta\rho_{\max} = 0.09 \text{ e \AA}^{-3}$
328 parameters	$\Delta\rho_{\min} = -0.15 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.5720 (2)	0.58755 (7)	-0.07734 (6)	0.0569 (3)
N2	-0.45576 (19)	1.10512 (6)	0.36976 (5)	0.0481 (3)
N3	-0.00550 (19)	1.24674 (6)	0.23495 (6)	0.0497 (3)
C1	0.6991 (3)	0.65228 (9)	-0.06772 (8)	0.0543 (4)
H1	0.838 (3)	0.6547 (8)	-0.0931 (7)	0.062 (4)*
C2	0.6494 (2)	0.71361 (8)	-0.02588 (7)	0.0462 (3)
H2	0.751 (2)	0.7581 (8)	-0.0194 (7)	0.052 (4)*
C3	0.4553 (2)	0.71081 (7)	0.00886 (6)	0.0388 (3)
C4	0.3222 (2)	0.64283 (7)	-0.00011 (7)	0.0431 (3)
H4	0.185 (2)	0.6363 (7)	0.0227 (6)	0.044 (3)*
C5	0.3870 (3)	0.58440 (8)	-0.04271 (7)	0.0511 (4)
H5	0.293 (2)	0.5370 (8)	-0.0495 (6)	0.056 (4)*
C6	-0.5830 (2)	1.04076 (8)	0.35549 (7)	0.0479 (3)
H6	-0.722 (3)	1.0359 (8)	0.3800 (7)	0.060 (4)*
C7	-0.5305 (2)	0.98292 (8)	0.31014 (7)	0.0445 (3)
H7	-0.628 (2)	0.9374 (8)	0.3018 (6)	0.049 (4)*
C8	-0.3348 (2)	0.98987 (7)	0.27589 (6)	0.0385 (3)
C9	-0.2039 (2)	1.05746 (7)	0.29001 (6)	0.0411 (3)
H9	-0.064 (2)	1.0657 (7)	0.2688 (6)	0.041 (3)*
C10	-0.2692 (2)	1.11169 (8)	0.33646 (7)	0.0449 (3)
H10	-0.176 (2)	1.1575 (7)	0.3472 (6)	0.044 (3)*
C11	-0.1535 (2)	1.23391 (8)	0.17687 (7)	0.0487 (3)
H11	-0.298 (2)	1.2632 (7)	0.1723 (7)	0.055 (4)*
C12	-0.1104 (2)	1.18348 (8)	0.12416 (7)	0.0453 (3)
H12	-0.223 (2)	1.1766 (8)	0.0831 (7)	0.057 (4)*
C13	0.0985 (2)	1.14333 (7)	0.12916 (6)	0.0384 (3)
C14	0.2518 (2)	1.15541 (7)	0.19037 (7)	0.0415 (3)
H14	0.401 (2)	1.1294 (7)	0.1986 (6)	0.049 (4)*
C15	0.1931 (2)	1.20645 (8)	0.24030 (7)	0.0473 (3)
H15	0.300 (2)	1.2158 (8)	0.2839 (7)	0.059 (4)*
C16	0.3959 (2)	0.77741 (7)	0.05039 (6)	0.0414 (3)
H16	0.504 (2)	0.8205 (7)	0.0528 (6)	0.040 (3)*
C17	0.2082 (2)	0.78441 (7)	0.08307 (6)	0.0404 (3)
H17	0.100 (2)	0.7412 (7)	0.0822 (6)	0.043 (3)*
C18	0.1516 (2)	0.85257 (7)	0.12091 (6)	0.0432 (3)
H18	0.260 (2)	0.8965 (7)	0.1235 (6)	0.049 (4)*
C19	-0.0351 (2)	0.85901 (7)	0.15427 (7)	0.0431 (3)
H19	-0.140 (2)	0.8147 (7)	0.1538 (6)	0.047 (4)*

supplementary materials

C20	-0.0919 (2)	0.92526 (7)	0.19440 (6)	0.0414 (3)
H20	0.012 (2)	0.9683 (8)	0.1966 (6)	0.044 (3)*
C21	-0.2736 (2)	0.92738 (7)	0.23017 (6)	0.0420 (3)
H21	-0.376 (2)	0.8825 (8)	0.2281 (6)	0.049 (4)*
C22	0.1515 (2)	1.09529 (7)	0.07030 (7)	0.0420 (3)
H22	0.030 (2)	1.0935 (7)	0.0295 (7)	0.050 (4)*
C23	0.3498 (2)	1.05781 (7)	0.06546 (7)	0.0417 (3)
H23	0.473 (2)	1.0581 (7)	0.1052 (6)	0.044 (3)*
C24	0.3997 (2)	1.01778 (7)	0.00320 (7)	0.0439 (3)
H24	0.275 (2)	1.0192 (7)	-0.0377 (7)	0.052 (4)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0517 (7)	0.0549 (8)	0.0644 (8)	0.0045 (6)	0.0091 (6)	-0.0115 (6)
N2	0.0471 (7)	0.0512 (7)	0.0468 (6)	0.0063 (5)	0.0095 (5)	0.0009 (5)
N3	0.0557 (7)	0.0448 (6)	0.0486 (7)	0.0072 (5)	0.0073 (6)	-0.0015 (5)
C1	0.0454 (9)	0.0607 (9)	0.0581 (9)	0.0038 (7)	0.0118 (7)	-0.0031 (7)
C2	0.0441 (8)	0.0453 (8)	0.0494 (8)	-0.0044 (6)	0.0065 (6)	0.0033 (6)
C3	0.0428 (7)	0.0382 (7)	0.0340 (6)	0.0018 (5)	0.0002 (5)	0.0060 (5)
C4	0.0428 (8)	0.0420 (7)	0.0441 (7)	-0.0017 (6)	0.0049 (6)	0.0013 (6)
C5	0.0499 (9)	0.0437 (8)	0.0587 (9)	-0.0027 (6)	0.0034 (7)	-0.0047 (7)
C6	0.0413 (8)	0.0560 (9)	0.0477 (8)	0.0050 (6)	0.0107 (6)	0.0079 (7)
C7	0.0410 (8)	0.0444 (8)	0.0479 (8)	-0.0026 (6)	0.0052 (6)	0.0071 (6)
C8	0.0396 (7)	0.0377 (7)	0.0371 (7)	0.0021 (5)	0.0012 (5)	0.0065 (5)
C9	0.0384 (7)	0.0425 (7)	0.0431 (7)	-0.0015 (6)	0.0076 (6)	0.0014 (6)
C10	0.0451 (8)	0.0422 (8)	0.0473 (8)	-0.0008 (6)	0.0060 (6)	-0.0024 (6)
C11	0.0467 (8)	0.0480 (8)	0.0518 (8)	0.0099 (6)	0.0080 (7)	0.0049 (6)
C12	0.0437 (8)	0.0490 (8)	0.0417 (8)	0.0016 (6)	-0.0003 (6)	0.0050 (6)
C13	0.0437 (7)	0.0339 (6)	0.0377 (7)	-0.0030 (5)	0.0060 (6)	0.0042 (5)
C14	0.0411 (8)	0.0376 (7)	0.0450 (7)	0.0022 (6)	0.0027 (6)	-0.0003 (6)
C15	0.0531 (9)	0.0423 (7)	0.0445 (8)	0.0033 (6)	-0.0009 (6)	-0.0040 (6)
C16	0.0497 (8)	0.0359 (7)	0.0379 (7)	-0.0048 (6)	0.0029 (6)	0.0035 (5)
C17	0.0504 (8)	0.0361 (7)	0.0340 (7)	-0.0020 (6)	0.0027 (6)	0.0031 (5)
C18	0.0553 (9)	0.0364 (7)	0.0370 (7)	-0.0019 (6)	0.0026 (6)	0.0023 (6)
C19	0.0517 (8)	0.0369 (7)	0.0394 (7)	-0.0006 (6)	0.0011 (6)	0.0013 (6)
C20	0.0476 (8)	0.0364 (7)	0.0391 (7)	-0.0025 (6)	0.0022 (6)	0.0018 (5)
C21	0.0463 (8)	0.0356 (7)	0.0431 (7)	-0.0027 (6)	0.0022 (6)	0.0030 (6)
C22	0.0480 (8)	0.0395 (7)	0.0377 (7)	-0.0033 (6)	0.0028 (6)	0.0017 (5)
C23	0.0503 (8)	0.0351 (7)	0.0394 (7)	-0.0046 (6)	0.0049 (6)	0.0004 (5)
C24	0.0541 (8)	0.0363 (7)	0.0415 (7)	-0.0046 (6)	0.0073 (7)	0.0006 (6)

Geometric parameters (\AA , $^\circ$)

N1—C1	1.3355 (18)	C11—H11	0.975 (14)
N1—C5	1.3436 (18)	C12—C13	1.3931 (18)
N2—C6	1.3394 (17)	C12—H12	0.964 (14)
N2—C10	1.3396 (16)	C13—C14	1.3952 (17)
N3—C11	1.3354 (17)	C13—C22	1.4663 (17)


N3—C15	1.3422 (17)	C14—C15	1.3765 (18)
C1—C2	1.380 (2)	C14—H14	0.971 (13)
C1—H1	0.999 (15)	C15—H15	0.989 (13)
C2—C3	1.3910 (18)	C16—C17	1.3394 (18)
C2—H2	0.964 (13)	C16—H16	0.969 (12)
C3—C4	1.3999 (18)	C17—C18	1.4395 (18)
C3—C16	1.4627 (18)	C17—H17	0.973 (13)
C4—C5	1.3796 (19)	C18—C19	1.3408 (19)
C4—H4	0.970 (13)	C18—H18	0.982 (13)
C5—H5	0.980 (14)	C19—C20	1.4382 (18)
C6—C7	1.3830 (19)	C19—H19	0.976 (13)
C6—H6	0.997 (14)	C20—C21	1.3390 (18)
C7—C8	1.3974 (18)	C20—H20	0.954 (13)
C7—H7	0.969 (13)	C21—H21	0.973 (13)
C8—C9	1.3961 (17)	C22—C23	1.3380 (18)
C8—C21	1.4616 (17)	C22—H22	0.985 (13)
C9—C10	1.3788 (18)	C23—C24	1.4436 (18)
C9—H9	0.969 (12)	C23—H23	0.978 (12)
C10—H10	0.964 (13)	C24—C24 ⁱ	1.341 (3)
C11—C12	1.3815 (18)	C24—H24	0.997 (13)
C1—N1—C5	115.69 (12)	C13—C12—H12	119.3 (8)
C6—N2—C10	115.81 (12)	C12—C13—C14	116.18 (12)
C11—N3—C15	115.75 (12)	C12—C13—C22	119.88 (11)
N1—C1—C2	124.04 (14)	C14—C13—C22	123.85 (12)
N1—C1—H1	116.0 (8)	C15—C14—C13	119.46 (12)
C2—C1—H1	120.0 (8)	C15—C14—H14	118.7 (7)
C1—C2—C3	120.14 (13)	C13—C14—H14	121.9 (7)
C1—C2—H2	120.4 (8)	N3—C15—C14	124.58 (13)
C3—C2—H2	119.5 (8)	N3—C15—H15	115.0 (8)
C2—C3—C4	116.35 (12)	C14—C15—H15	120.4 (8)
C2—C3—C16	120.18 (12)	C17—C16—C3	126.49 (12)
C4—C3—C16	123.45 (12)	C17—C16—H16	118.4 (7)
C5—C4—C3	119.21 (13)	C3—C16—H16	115.1 (7)
C5—C4—H4	119.6 (7)	C16—C17—C18	124.57 (13)
C3—C4—H4	121.2 (7)	C16—C17—H17	119.5 (7)
N1—C5—C4	124.55 (13)	C18—C17—H17	115.9 (7)
N1—C5—H5	116.1 (8)	C19—C18—C17	124.39 (13)
C4—C5—H5	119.3 (8)	C19—C18—H18	118.2 (7)
N2—C6—C7	123.92 (13)	C17—C18—H18	117.4 (7)
N2—C6—H6	116.0 (8)	C18—C19—C20	125.63 (13)
C7—C6—H6	120.1 (8)	C18—C19—H19	118.8 (7)
C6—C7—C8	119.99 (13)	C20—C19—H19	115.5 (8)
C6—C7—H7	120.5 (8)	C21—C20—C19	123.54 (13)
C8—C7—H7	119.5 (8)	C21—C20—H20	119.9 (7)
C9—C8—C7	116.10 (12)	C19—C20—H20	116.5 (7)
C9—C8—C21	123.76 (12)	C20—C21—C8	127.09 (12)
C7—C8—C21	120.12 (12)	C20—C21—H21	118.8 (7)
C10—C9—C8	119.73 (12)	C8—C21—H21	114.1 (8)

supplementary materials

C10—C9—H9	119.7 (7)	C23—C22—C13	126.71 (12)
C8—C9—H9	120.5 (7)	C23—C22—H22	117.8 (7)
N2—C10—C9	124.44 (13)	C13—C22—H22	115.3 (7)
N2—C10—H10	115.9 (7)	C22—C23—C24	124.00 (13)
C9—C10—H10	119.6 (7)	C22—C23—H23	119.9 (7)
N3—C11—C12	123.79 (13)	C24—C23—H23	116.1 (7)
N3—C11—H11	116.4 (8)	C24 ⁱ —C24—C23	124.67 (17)
C12—C11—H11	119.8 (8)	C24 ⁱ —C24—H24	119.5 (8)
C11—C12—C13	120.21 (13)	C23—C24—H24	115.8 (8)
C11—C12—H12	120.5 (8)		
C5—N1—C1—C2	-0.5 (2)	C11—C12—C13—C22	-174.59 (11)
N1—C1—C2—C3	-0.7 (2)	C12—C13—C14—C15	-1.72 (18)
C1—C2—C3—C4	1.43 (18)	C22—C13—C14—C15	174.85 (12)
C1—C2—C3—C16	-176.92 (12)	C11—N3—C15—C14	1.1 (2)
C2—C3—C4—C5	-1.03 (18)	C13—C14—C15—N3	0.1 (2)
C16—C3—C4—C5	177.26 (11)	C2—C3—C16—C17	175.93 (12)
C1—N1—C5—C4	0.9 (2)	C4—C3—C16—C17	-2.3 (2)
C3—C4—C5—N1	-0.1 (2)	C3—C16—C17—C18	-177.16 (11)
C10—N2—C6—C7	0.60 (19)	C16—C17—C18—C19	-179.16 (12)
N2—C6—C7—C8	-0.2 (2)	C17—C18—C19—C20	177.79 (11)
C6—C7—C8—C9	-0.64 (17)	C18—C19—C20—C21	-176.06 (13)
C6—C7—C8—C21	177.48 (11)	C19—C20—C21—C8	175.54 (11)
C7—C8—C9—C10	1.12 (17)	C9—C8—C21—C20	-1.5 (2)
C21—C8—C9—C10	-176.93 (11)	C7—C8—C21—C20	-179.51 (12)
C6—N2—C10—C9	-0.08 (19)	C12—C13—C22—C23	175.18 (13)
C8—C9—C10—N2	-0.8 (2)	C14—C13—C22—C23	-1.3 (2)
C15—N3—C11—C12	-0.6 (2)	C13—C22—C23—C24	-173.53 (11)
N3—C11—C12—C13	-1.0 (2)	C22—C23—C24—C24 ⁱ	177.66 (15)
C11—C12—C13—C14	2.12 (18)		

Symmetry codes: (i) $-x+1, -y+2, -z$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

