

## Methyl 3-(4-bromophenyl)-1-methyl-1,2,3,3a,4,9b-hexahydrobenzo[f]-chromeno[4,3-b]pyrrole-3a-carboxylate

S. Nirmala,<sup>a</sup> E. Theboral Sugi Kamala,<sup>a</sup> L. Sudha,<sup>b\*</sup>  
S. Kathiravan<sup>c</sup> and R. Raghunathan<sup>c</sup>

<sup>a</sup>Department of Physics, Easwari Engineering College, Ramapuram, Chennai 600 089, India, <sup>b</sup>Department of Physics, SRM University, Ramapuram Campus, Chennai 600 089, India, and <sup>c</sup>Department of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India  
Correspondence e-mail: sudharose18@gmail.com


Received 17 July 2009; accepted 23 July 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.038;  $wR$  factor = 0.102; data-to-parameter ratio = 23.5.

In the title compound,  $\text{C}_{24}\text{H}_{22}\text{BrNO}_3$ , the dihydropyran ring adopts a half-chair conformation, whereas the pyrrolidine ring is in an envelope conformation. The bromophenyl group is oriented at an angle of  $66.44(4)^\circ$  with respect to the naphthalene ring system. In the crystal structure, molecules are linked into centrosymmetric dimers by  $\text{C}-\text{H}\cdots\pi$  interactions and the dimers are connected via  $\text{C}-\text{H}\cdots\text{Br}$  hydrogen bonds. The crystal structure is further stabilized by  $\pi-\pi$  interactions [centroid–centroid distance =  $3.453(1)\text{ \AA}$ ].

### Related literature

For the biological activity of pyrrole derivatives, see: Biava *et al.* (2005); Borthwick *et al.* (2000); Caine (1993); Carlson (1993); Fernandes *et al.* (2004); Jiang *et al.* (2004); Sokoloff *et al.* (1990); Tidey (1992); Wilner (1985). For a related structure, see: Nirmala *et al.* (2009). For ring-puckering parameters, see: Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_{24}\text{H}_{22}\text{BrNO}_3$   
 $M_r = 452.34$   
Monoclinic,  $P2_1/c$   
 $a = 12.7856(4)\text{ \AA}$   
 $b = 19.9348(6)\text{ \AA}$   
 $c = 8.0189(3)\text{ \AA}$   
 $\beta = 106.163(2)^\circ$

$V = 1963.06(11)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 2.12\text{ mm}^{-1}$   
 $T = 293\text{ K}$   
 $0.25 \times 0.20 \times 0.15\text{ mm}$

#### Data collection

Bruker Kappa APEXII area-detector diffractometer  
Absorption correction: multi-scan (Blessing, 1995)  
 $T_{\min} = 0.619$ ,  $T_{\max} = 0.742$

49847 measured reflections  
6162 independent reflections  
4068 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.102$   
 $S = 1.02$   
6162 reflections

262 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.58\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.51\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C24—H24A $\cdots$ BrI <sup>i</sup>	0.96	2.84	3.789 (3)	172
C20—H20 $\cdots$ Cg1 <sup>ii</sup>	0.93	2.77	3.653 (2)	160

Symmetry codes: (i)  $x + 1, -y + \frac{3}{2}, z + \frac{1}{2}$ ; (ii)  $-x + 1, -y, -z + 2$ . Cg1 is the centroid of the C3–C8 ring.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *PLATON* (Spek, 2009).

SN thanks Dr Babu Vargheese, SAIF, IIT Madras, India, for his help with the data collection and SRM management, India, for their support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2858).

### References

- Biava, M., Porretta, G. C., Poce, G., Deidda, D., Pompei, R., Tafi, A. & Manetti, F. (2005). *Bioorg. Med. Chem.* **13**, 1221–1230.  
Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.  
Borthwick, A. D., Angier, S. J., Crame, A. J., Exall, A. M., Haley, T. M., Hart, G. J., Mason, A. M., Pennell, A. M. K. & Weingarten, G. G. (2000). *J. Med. Chem.* **43**, 4452–4464.  
Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Caine, B. (1993). *Science*, **260**, 1814–1816.  
Carlson, J. (1993). *Neur. Transm.* **94**, 11–19.  
Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Fernandes, E., Costa, D., Toste, S. A., Lima, J. L. & Reis, S. (2004). *Free Radical Biol. Med.* **37**, 1895–1905.  
Jiang, S., Lu, H., Liu, S., Zhao, Q., He, Y. & Debnath, A. K. (2004). *Antimicrob. Agents Chemother.* **48**, 4349–4359.

- Nirmala, S., Kamala, E. T. S., Sudha, L., Kathiravan, S. & Raghunathan, R. (2009). *Acta Cryst. E*65, o1811.
- Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
- Sokoloff, P., Giros, B., Martres, M. P., Bouthenet, M. L. & Schwartz, J. C. (1990). *Nature (London)*, 347, 147–151.
- Spek, A. L. (2009). *Acta Cryst. D*65, 148–155.
- Tidey, J. W. (1992). *Behav. Pharm.* 3, 553–566.
- Wilner, P. (1985). *Clin. Neuropharmac.* 18, suppl.1, 549–556.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o2028-o2029 [doi:10.1107/S1600536809029389]

## Methyl 3-(4-bromophenyl)-1-methyl-1,2,3,3a,4,9b-hexahydrobenzo[f]chromeno[4,3-*b*]pyrrole-3a-carboxylate

S. Nirmala, E. T. S. Kamala, L. Sudha, S. Kathiravan and R. Raghunathan

### Comment

Chromenopyrrole compounds are used in the treatment of impulsive disorders (Caine, 1993), aggressiveness (Tidey, 1992), parkinson's disease (Carlson, 1993), psychoses, memory disorders (Sokoloff *et al.*, 1990), anxiety and depression (Wilner, 1985). Pyrrole derivatives have good *in vitro* activities against mycobacteria and candidae (Biava *et al.*, 2005). These derivatives also possess anti-inflammatory (Fernandes *et al.*, 2004) and antiviral (Borthwick *et al.*, 2000) activities. It has also been shown that N-substituted pyrrole derivatives inhibit human immuno deficiency virus type-I (HIV-I) (Jiang *et al.*, 2004). In view of its medicinal importance, the crystal structure determination of the title compound was undertaken.

The geometric parameters of the title molecule (Fig. 1) agree well with those reported for a similar structure (Nirmala *et al.*, 2009). The sum of bond angles around atom N1 [332.0 (8) $^{\circ}$ ] is in accordance with  $sp^3$  hybridization. The naphthalene ring system (C2-C11) and the bromophenyl group (Br1/C16-C21) are oriented at an angle of 66.44 (4) $^{\circ}$  with respect to each other. The heterocyclic ring (O1/C1/C2/C11-C13) of the chromenopyrrole unit adopts a half chair conformation with puckering parameters  $Q$  = 0.468 (2) Å,  $\theta$  = 132.5 (2) $^{\circ}$  and  $\varphi$  = 83.1 (3) $^{\circ}$  (Cremer & Pople, 1975). The pyrrolidine ring (N1/C1/C13-C15) adopts an envelope conformation with puckering parameters  $q_2$  = 0.433 (2) Å and  $\varphi$  = 214.5 (3) $^{\circ}$  (Cremer & Pople, 1975). Atom C1 deviates by -0.654 Å from the least-square plane through the remaining four atoms.

The crystal packing is stabilized by intermolecular C—H $\cdots$ Br hydrogen bonds. The molecules are linked into centrosymmetric dimers by C—H $\cdots$  $\pi$  (C20—H20 $\cdots$ Cg1; Cg1 is the centroid of the C3—C8 ring) interactions (Table 1). In addition,  $\pi$ — $\pi$  interactions between C3—C8 rings at ( $x$ ,  $y$ ,  $z$ ) and (2 - $x$ , 1 - $y$ , 2 - $z$ ) stabilize the structure, with a centroid-to-centroid distance of 3.453 (1) Å.

### Experimental

A mixture of (*Z*)-methyl 2-((1-formylnaphthalen-2-yloxy)methyl)-3-(4-bromophenyl)acrylate (20 mmol) and sarcosine (30 mmol) were refluxed in benzene for 20 h and the solvent was removed under reduced pressure. The crude product was subjected to column chromatography to get the pure product. Chloroform and methanol (1:1) solvent mixture was used for the crystallization under slow evaporation method.

### Refinement

H atoms were placed in idealized positions and allowed to ride on their parent atoms, with C-H = 0.93, 0.98, 0.97 and 0.96 Å for aromatic, methine, methylene and methyl H respectively, and  $U_{\text{iso}}(\text{H})$  = 1.5 $U_{\text{eq}}(\text{C})$  for methyl H and  $U_{\text{iso}}(\text{H})$  = 1.2 $U_{\text{eq}}(\text{C})$  for all other H atoms. Reflections 110 and 100 were omitted during the final cycles of refinement since they were seriously effected by the beamstop.

# supplementary materials

---

## Figures




Fig. 1. The molecular structure of the title compound, with 30% probability displacement ellipsoids.




Fig. 2. The packing of the molecules viewed down  $a$  axis. H atoms not involved in C—H···Br and C—H··· $\pi$  interactions have been omitted.

## Methyl 3-(4-bromophenyl)-1-methyl-1,2,3,3a,4,9b-hexahydrobenzo[*f*]chromeno[4,3-*b*]pyrrole-3a-carboxylate

### Crystal data

$C_{24}H_{22}BrNO_3$	$F_{000} = 928$
$M_r = 452.34$	$D_x = 1.531 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 14860 reflections
$a = 12.7856 (4) \text{ \AA}$	$\theta = 2.6\text{--}25.2^\circ$
$b = 19.9348 (6) \text{ \AA}$	$\mu = 2.12 \text{ mm}^{-1}$
$c = 8.0189 (3) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 106.163 (2)^\circ$	Prism, colourless
$V = 1963.06 (11) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.15 \text{ mm}$
$Z = 4$	

### Data collection

Bruker Kappa APEXII area-detector diffractometer	6162 independent reflections
Radiation source: fine-focus sealed tube	4068 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.031$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 30.9^\circ$
$\omega$ and $\varphi$ scans	$\theta_{\text{min}} = 2.0^\circ$
Absorption correction: multi-scan (Blessing, 1995)	$h = -18 \rightarrow 18$
$T_{\text{min}} = 0.619$ , $T_{\text{max}} = 0.742$	$k = -16 \rightarrow 28$
49847 measured reflections	$l = -11 \rightarrow 11$

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.038$	H-atom parameters constrained

$$wR(F^2) = 0.102 \quad w = 1/[\sigma^2(F_o)^2 + (0.0439P)^2 + 0.7192P] \\ \text{where } P = (E_x^2 + 2E_y^2)/3$$

$S = 1.02$	$(\Delta/\sigma)_{\text{max}} = 0.001$
6162 reflections	$\Delta\rho_{\text{max}} = 0.58 \text{ e \AA}^{-3}$
262 parameters	$\Delta\rho_{\text{min}} = -0.51 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

### *Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.68758 (13)	0.58289 (8)	0.6893 (2)	0.0281 (3)
H1	0.7433	0.6171	0.6940	0.034*
C2	0.73957 (13)	0.52048 (8)	0.7836 (2)	0.0287 (3)
C3	0.84557 (13)	0.49812 (8)	0.7827 (2)	0.0306 (3)
C4	0.91361 (15)	0.53431 (10)	0.7029 (2)	0.0381 (4)
H4	0.8889	0.5745	0.6470	0.046*
C5	1.01452 (16)	0.51164 (11)	0.7061 (2)	0.0455 (5)
H5	1.0576	0.5367	0.6530	0.055*
C6	1.05416 (17)	0.45149 (12)	0.7878 (3)	0.0510 (5)
H6	1.1226	0.4360	0.7869	0.061*
C7	0.99246 (17)	0.41579 (11)	0.8685 (3)	0.0474 (5)
H7	1.0195	0.3759	0.9241	0.057*
C8	0.88813 (15)	0.43783 (9)	0.8699 (2)	0.0357 (4)
C9	0.82649 (16)	0.40202 (9)	0.9600 (3)	0.0425 (4)
H9	0.8535	0.3620	1.0149	0.051*
C10	0.72848 (16)	0.42484 (9)	0.9681 (2)	0.0417 (4)
H10	0.6893	0.4013	1.0309	0.050*
C11	0.68588 (14)	0.48451 (8)	0.8811 (2)	0.0329 (3)
C12	0.52824 (14)	0.55294 (8)	0.7923 (2)	0.0338 (4)
H12A	0.4714	0.5700	0.8396	0.041*
H12B	0.4935	0.5324	0.6810	0.041*
C13	0.59903 (13)	0.61089 (8)	0.7656 (2)	0.0289 (3)
C14	0.53395 (14)	0.65915 (8)	0.6186 (2)	0.0333 (4)
H14	0.5695	0.7031	0.6403	0.040*
C15	0.55720 (16)	0.63062 (10)	0.4547 (2)	0.0424 (4)

## supplementary materials

---

H15A	0.4896	0.6189	0.3692	0.051*
H15B	0.5952	0.6635	0.4040	0.051*
C16	0.41597 (13)	0.66967 (8)	0.6127 (2)	0.0308 (3)
C17	0.38794 (16)	0.72456 (9)	0.6967 (3)	0.0449 (5)
H17	0.4420	0.7544	0.7536	0.054*
C18	0.28173 (17)	0.73619 (10)	0.6981 (3)	0.0491 (5)
H18	0.2645	0.7732	0.7558	0.059*
C19	0.20242 (15)	0.69243 (9)	0.6136 (2)	0.0379 (4)
C20	0.22678 (14)	0.63726 (9)	0.5282 (2)	0.0362 (4)
H20	0.1721	0.6077	0.4715	0.043*
C21	0.33319 (14)	0.62635 (8)	0.5280 (2)	0.0337 (4)
H21	0.3499	0.5892	0.4699	0.040*
C22	0.68903 (16)	0.55698 (9)	0.3883 (2)	0.0385 (4)
H22A	0.7326	0.5177	0.4259	0.058*
H22B	0.7357	0.5945	0.3858	0.058*
H22C	0.6413	0.5496	0.2740	0.058*
C23	0.63964 (14)	0.65052 (8)	0.9323 (2)	0.0338 (4)
C24	0.7836 (2)	0.71536 (12)	1.1053 (3)	0.0631 (7)
H24A	0.8555	0.7299	1.1072	0.095*
H24B	0.7869	0.6896	1.2078	0.095*
H24C	0.7378	0.7538	1.1018	0.095*
N1	0.62481 (12)	0.57086 (7)	0.50776 (18)	0.0322 (3)
O1	0.58933 (10)	0.50339 (6)	0.90663 (17)	0.0412 (3)
O2	0.73947 (11)	0.67466 (8)	0.9542 (2)	0.0529 (4)
O3	0.58771 (13)	0.66160 (9)	1.03113 (19)	0.0594 (4)
Br1	0.056104 (18)	0.708461 (13)	0.61041 (4)	0.06427 (10)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0292 (8)	0.0257 (7)	0.0314 (8)	-0.0020 (6)	0.0121 (6)	0.0029 (6)
C2	0.0324 (8)	0.0275 (7)	0.0275 (8)	0.0002 (6)	0.0103 (7)	0.0006 (6)
C3	0.0332 (8)	0.0333 (8)	0.0255 (8)	0.0009 (6)	0.0085 (6)	-0.0059 (6)
C4	0.0382 (9)	0.0452 (10)	0.0322 (9)	0.0007 (7)	0.0118 (7)	-0.0013 (7)
C5	0.0370 (10)	0.0638 (13)	0.0382 (10)	-0.0007 (9)	0.0144 (8)	-0.0083 (9)
C6	0.0379 (10)	0.0685 (14)	0.0469 (12)	0.0140 (10)	0.0121 (9)	-0.0119 (10)
C7	0.0477 (11)	0.0486 (11)	0.0429 (11)	0.0169 (9)	0.0076 (9)	-0.0056 (9)
C8	0.0382 (9)	0.0375 (9)	0.0297 (9)	0.0063 (7)	0.0064 (7)	-0.0052 (7)
C9	0.0523 (11)	0.0324 (9)	0.0405 (10)	0.0103 (8)	0.0092 (9)	0.0053 (8)
C10	0.0512 (11)	0.0344 (9)	0.0426 (10)	0.0008 (8)	0.0181 (9)	0.0105 (8)
C11	0.0362 (9)	0.0315 (8)	0.0326 (9)	0.0004 (7)	0.0124 (7)	0.0030 (7)
C12	0.0304 (8)	0.0343 (8)	0.0397 (9)	0.0005 (6)	0.0146 (7)	0.0089 (7)
C13	0.0278 (8)	0.0288 (7)	0.0324 (8)	0.0006 (6)	0.0122 (6)	0.0043 (6)
C14	0.0322 (8)	0.0302 (8)	0.0375 (9)	0.0009 (6)	0.0100 (7)	0.0072 (7)
C15	0.0399 (10)	0.0523 (11)	0.0375 (10)	0.0099 (8)	0.0150 (8)	0.0143 (8)
C16	0.0321 (8)	0.0260 (7)	0.0334 (9)	0.0029 (6)	0.0077 (7)	0.0028 (6)
C17	0.0396 (10)	0.0328 (9)	0.0559 (12)	0.0008 (7)	0.0024 (9)	-0.0141 (8)
C18	0.0480 (11)	0.0385 (10)	0.0585 (13)	0.0110 (8)	0.0106 (10)	-0.0155 (9)

C19	0.0348 (9)	0.0392 (9)	0.0401 (10)	0.0095 (7)	0.0110 (8)	0.0041 (7)
C20	0.0348 (9)	0.0346 (9)	0.0373 (9)	-0.0027 (7)	0.0068 (7)	-0.0015 (7)
C21	0.0377 (9)	0.0293 (8)	0.0346 (9)	0.0016 (7)	0.0110 (7)	-0.0045 (7)
C22	0.0453 (10)	0.0421 (10)	0.0315 (9)	-0.0002 (8)	0.0164 (8)	0.0016 (7)
C23	0.0333 (9)	0.0317 (8)	0.0367 (9)	0.0059 (7)	0.0103 (7)	0.0042 (7)
C24	0.0480 (13)	0.0636 (14)	0.0739 (16)	-0.0068 (10)	0.0107 (11)	-0.0321 (12)
N1	0.0346 (7)	0.0358 (7)	0.0280 (7)	-0.0006 (6)	0.0118 (6)	0.0051 (6)
O1	0.0393 (7)	0.0416 (7)	0.0498 (8)	0.0059 (5)	0.0238 (6)	0.0190 (6)
O2	0.0394 (7)	0.0582 (9)	0.0646 (9)	-0.0116 (6)	0.0203 (7)	-0.0283 (7)
O3	0.0555 (9)	0.0822 (11)	0.0476 (8)	-0.0085 (8)	0.0263 (7)	-0.0169 (8)
Br1	0.04256 (13)	0.07000 (17)	0.0859 (2)	0.01747 (10)	0.02724 (12)	0.00945 (12)

*Geometric parameters (Å, °)*

C1—N1	1.473 (2)	C14—C16	1.511 (2)
C1—C2	1.510 (2)	C14—C15	1.536 (3)
C1—C13	1.534 (2)	C14—H14	0.98
C1—H1	0.98	C15—N1	1.464 (2)
C2—C11	1.377 (2)	C15—H15A	0.97
C2—C3	1.429 (2)	C15—H15B	0.97
C3—C4	1.413 (2)	C16—C17	1.383 (2)
C3—C8	1.421 (2)	C16—C21	1.388 (2)
C4—C5	1.361 (3)	C17—C18	1.381 (3)
C4—H4	0.93	C17—H17	0.93
C5—C6	1.392 (3)	C18—C19	1.366 (3)
C5—H5	0.93	C18—H18	0.93
C6—C7	1.353 (3)	C19—C20	1.376 (3)
C6—H6	0.93	C19—Br1	1.8910 (18)
C7—C8	1.408 (3)	C20—C21	1.378 (2)
C7—H7	0.93	C20—H20	0.93
C8—C9	1.403 (3)	C21—H21	0.93
C9—C10	1.352 (3)	C22—N1	1.452 (2)
C9—H9	0.93	C22—H22A	0.96
C10—C11	1.410 (2)	C22—H22B	0.96
C10—H10	0.93	C22—H22C	0.96
C11—O1	1.360 (2)	C23—O3	1.188 (2)
C12—O1	1.424 (2)	C23—O2	1.329 (2)
C12—C13	1.519 (2)	C24—O2	1.437 (3)
C12—H12A	0.97	C24—H24A	0.96
C12—H12B	0.97	C24—H24B	0.96
C13—C23	1.514 (2)	C24—H24C	0.96
C13—C14	1.569 (2)		
N1—C1—C2	113.78 (13)	C16—C14—C13	115.15 (13)
N1—C1—C13	101.17 (13)	C15—C14—C13	103.15 (13)
C2—C1—C13	111.80 (13)	C16—C14—H14	107.0
N1—C1—H1	109.9	C15—C14—H14	107.0
C2—C1—H1	109.9	C13—C14—H14	107.0
C13—C1—H1	109.9	N1—C15—C14	106.96 (14)
C11—C2—C3	117.65 (15)	N1—C15—H15A	110.3

## supplementary materials

---

C11—C2—C1	119.57 (14)	C14—C15—H15A	110.3
C3—C2—C1	122.72 (14)	N1—C15—H15B	110.3
C4—C3—C8	117.08 (16)	C14—C15—H15B	110.3
C4—C3—C2	123.20 (15)	H15A—C15—H15B	108.6
C8—C3—C2	119.69 (15)	C17—C16—C21	117.69 (16)
C5—C4—C3	121.46 (18)	C17—C16—C14	119.11 (15)
C5—C4—H4	119.3	C21—C16—C14	123.20 (15)
C3—C4—H4	119.3	C18—C17—C16	121.72 (17)
C4—C5—C6	120.93 (19)	C18—C17—H17	119.1
C4—C5—H5	119.5	C16—C17—H17	119.1
C6—C5—H5	119.5	C19—C18—C17	118.97 (17)
C7—C6—C5	119.57 (18)	C19—C18—H18	120.5
C7—C6—H6	120.2	C17—C18—H18	120.5
C5—C6—H6	120.2	C18—C19—C20	121.18 (17)
C6—C7—C8	121.30 (19)	C18—C19—Br1	119.58 (14)
C6—C7—H7	119.4	C20—C19—Br1	119.23 (14)
C8—C7—H7	119.4	C19—C20—C21	119.14 (16)
C9—C8—C7	120.97 (17)	C19—C20—H20	120.4
C9—C8—C3	119.39 (16)	C21—C20—H20	120.4
C7—C8—C3	119.61 (18)	C20—C21—C16	121.29 (15)
C10—C9—C8	120.98 (17)	C20—C21—H21	119.4
C10—C9—H9	119.5	C16—C21—H21	119.4
C8—C9—H9	119.5	N1—C22—H22A	109.5
C9—C10—C11	119.71 (17)	N1—C22—H22B	109.5
C9—C10—H10	120.1	H22A—C22—H22B	109.5
C11—C10—H10	120.1	N1—C22—H22C	109.5
O1—C11—C2	123.97 (15)	H22A—C22—H22C	109.5
O1—C11—C10	113.62 (15)	H22B—C22—H22C	109.5
C2—C11—C10	122.39 (16)	O3—C23—O2	122.60 (17)
O1—C12—C13	112.14 (14)	O3—C23—C13	124.61 (16)
O1—C12—H12A	109.2	O2—C23—C13	112.73 (14)
C13—C12—H12A	109.2	O2—C24—H24A	109.5
O1—C12—H12B	109.2	O2—C24—H24B	109.5
C13—C12—H12B	109.2	H24A—C24—H24B	109.5
H12A—C12—H12B	107.9	O2—C24—H24C	109.5
C23—C13—C12	110.09 (14)	H24A—C24—H24C	109.5
C23—C13—C1	115.59 (13)	H24B—C24—H24C	109.5
C12—C13—C1	108.33 (13)	C22—N1—C15	111.10 (13)
C23—C13—C14	108.83 (13)	C22—N1—C1	115.52 (14)
C12—C13—C14	111.11 (13)	C15—N1—C1	105.67 (13)
C1—C13—C14	102.70 (13)	C11—O1—C12	116.96 (12)
C16—C14—C15	117.01 (15)	C23—O2—C24	116.95 (16)
N1—C1—C2—C11	-94.11 (18)	C12—C13—C14—C16	37.3 (2)
C13—C1—C2—C11	19.7 (2)	C1—C13—C14—C16	152.89 (14)
N1—C1—C2—C3	88.72 (18)	C23—C13—C14—C15	147.21 (14)
C13—C1—C2—C3	-157.44 (15)	C12—C13—C14—C15	-91.41 (16)
C11—C2—C3—C4	-173.26 (16)	C1—C13—C14—C15	24.21 (16)
C1—C2—C3—C4	4.0 (2)	C16—C14—C15—N1	-125.38 (15)
C11—C2—C3—C8	4.7 (2)	C13—C14—C15—N1	2.14 (18)

C1—C2—C3—C8	-178.13 (15)	C15—C14—C16—C17	-143.63 (18)
C8—C3—C4—C5	1.5 (3)	C13—C14—C16—C17	94.9 (2)
C2—C3—C4—C5	179.42 (16)	C15—C14—C16—C21	37.0 (2)
C3—C4—C5—C6	0.3 (3)	C13—C14—C16—C21	-84.4 (2)
C4—C5—C6—C7	-1.5 (3)	C21—C16—C17—C18	0.5 (3)
C5—C6—C7—C8	0.7 (3)	C14—C16—C17—C18	-178.94 (19)
C6—C7—C8—C9	-177.30 (18)	C16—C17—C18—C19	-0.3 (3)
C6—C7—C8—C3	1.1 (3)	C17—C18—C19—C20	0.2 (3)
C4—C3—C8—C9	176.27 (16)	C17—C18—C19—Br1	-178.80 (16)
C2—C3—C8—C9	-1.8 (2)	C18—C19—C20—C21	-0.2 (3)
C4—C3—C8—C7	-2.1 (2)	Br1—C19—C20—C21	178.80 (13)
C2—C3—C8—C7	179.83 (16)	C19—C20—C21—C16	0.3 (3)
C7—C8—C9—C10	176.89 (18)	C17—C16—C21—C20	-0.5 (3)
C3—C8—C9—C10	-1.5 (3)	C14—C16—C21—C20	178.91 (16)
C8—C9—C10—C11	1.8 (3)	C12—C13—C23—O3	-37.5 (2)
C3—C2—C11—O1	173.77 (15)	C1—C13—C23—O3	-160.58 (17)
C1—C2—C11—O1	-3.5 (3)	C14—C13—C23—O3	84.5 (2)
C3—C2—C11—C10	-4.5 (3)	C12—C13—C23—O2	145.27 (15)
C1—C2—C11—C10	178.16 (16)	C1—C13—C23—O2	22.1 (2)
C9—C10—C11—O1	-177.08 (17)	C14—C13—C23—O2	-92.74 (17)
C9—C10—C11—C2	1.4 (3)	C14—C15—N1—C22	-155.30 (15)
O1—C12—C13—C23	-67.90 (18)	C14—C15—N1—C1	-29.30 (18)
O1—C12—C13—C1	59.38 (18)	C2—C1—N1—C22	-72.26 (17)
O1—C12—C13—C14	171.46 (13)	C13—C1—N1—C22	167.70 (13)
N1—C1—C13—C23	-160.07 (13)	C2—C1—N1—C15	164.50 (13)
C2—C1—C13—C23	78.49 (17)	C13—C1—N1—C15	44.46 (15)
N1—C1—C13—C12	75.89 (15)	C2—C11—O1—C12	16.4 (2)
C2—C1—C13—C12	-45.55 (18)	C10—C11—O1—C12	-165.22 (16)
N1—C1—C13—C14	-41.73 (14)	C13—C12—O1—C11	-45.0 (2)
C2—C1—C13—C14	-163.17 (13)	O3—C23—O2—C24	0.2 (3)
C23—C13—C14—C16	-84.11 (17)	C13—C23—O2—C24	177.54 (17)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
C24—H24A···Br1 <sup>i</sup>	0.96	2.84	3.789 (3)	172
C20—H20···Cg1 <sup>ii</sup>	0.93	2.77	3.653 (2)	160

Symmetry codes: (i)  $x+1, -y+3/2, z+1/2$ ; (ii)  $-x+1, -y, -z+2$ .

## supplementary materials

---

Fig. 1




Fig. 2

