

trans-Di- μ -iodido-bis[(3*H*-1,2-benzodithiole-3-thione)iodidomercury(II)]

El Adoui Laifa,^a Lamia Bendjeddou,^{a*} Naouel Boudraa,^a Slimane Dahaoui^b and Claude Lecomte^b

^aLaboratoire de Chimie Moléculaire, du Contrôle, de l'Environnement et des Mesures Physico-Chimiques, Faculté des Sciences, Département de Chimie, Université Mentouri de Constantine, 25000 Constantine, Algeria, and ^bCristallographie, Résonance Magnétique et Modélisation (CRM2), Université Henri Poincaré, Nancy 1, Faculté des Sciences, BP 70239, 54506 Vandoeuvre lès Nancy CEDEX, France
Correspondence e-mail: Lamiabendjeddou@yahoo.fr


Received 27 July 2009; accepted 13 August 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.015; wR factor = 0.037; data-to-parameter ratio = 42.0.

The complete molecule of the dinuclear title compound, $[\text{Hg}_2\text{I}_4(\text{C}_7\text{H}_4\text{S}_3)_2]$, is generated by crystallographic inversion symmetry. The complex has a dimeric structure in which each Hg^{II} ion adopts a tetrahedral geometry and is coordinated by two bridging I atoms, one terminal iodide ion and one thiocarbonyl S atom ($\text{C}=\text{S}$) of the ligand. The square plane formed by the Hg and I atoms and their symmetry counterparts makes a dihedral angle of $89.66(3)^\circ$ with the DDT plane. There is no classical hydrogen bonding, but weak $\text{S}\cdots\text{S}$ interactions of $3.4452(7)$ and $3.6859(7)$ Å maintain the cohesion of the crystal structure.

Related literature

For $\text{S}\cdots\text{S}$ interactions in sulfur-rich organic donor-acceptor compounds or radical salts, see: Cassoux *et al.* (1991); Klinsberg & Schrabber (1962). For the effects on the molecular packing and $\text{S}\cdots\text{S}$ contacts of modifying the structure of the organic molecule or changing the counter-ion or the co-crystallized solvent, see: Pullen & Olk (1999); Schlueter *et al.* (1996). In this context, a series of polymeric complexes has been reported with Ag^+ (Dai, Munakata, Kuroda-Sowa *et al.*, 1997; Dai, Kudora-Sowa *et al.*, 1997) and a tetra-nuclear CuI_4 cluster (Dai, Munakata, Wu *et al.*, 1997). For comparison bond lengths and angles in related chloride-bridged dimeric $\text{Hg}(\text{II})$ complexes, see: Brodersen & Hummel (1987); Dean (1978).



Experimental

Crystal data

$[\text{Hg}_2\text{I}_4(\text{C}_7\text{H}_4\text{S}_3)_2]$
 $M_r = 1277.40$
 Triclinic, $P\bar{1}$
 $a = 7.86285(10)$ Å
 $b = 8.19304(11)$ Å
 $c = 10.46506(13)$ Å
 $\alpha = 105.2917(11)^\circ$
 $\beta = 98.3031(10)^\circ$
 $\gamma = 105.6957(11)^\circ$
 $V = 608.92(2)$ Å³
 $Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 18.18$ mm⁻¹
 $T = 100$ K
 $0.16 \times 0.1 \times 0.08$ mm

Data collection

Oxford Diffraction SuperNova diffractometer with an Atlas detector
 Absorption correction: analytical (Clark *et al.*, 1995)
 $T_{\min} = 0.167$, $T_{\max} = 0.347$
 44929 measured reflections
 4951 independent reflections
 4698 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.051$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.015$
 $wR(F^2) = 0.037$
 $S = 1.05$
 4951 reflections
 118 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.20$ e Å⁻³
 $\Delta\rho_{\min} = -1.55$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

Hg1—S1	2.5169 (7)	Hg1—I1	2.8148 (4)
Hg1—I2	2.6599 (2)	Hg1—I1 ⁱ	3.1002 (4)
S1—Hg1—I2	138.842 (15)	I2—Hg1—I1 ⁱ	103.851 (13)
S1—Hg1—I1	98.374 (17)	I1—Hg1—I1 ⁱ	95.210 (10)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *CrysAlis CCD* (Oxford Diffraction, 2009); cell refinement: *CrysAlis RED* (Oxford Diffraction, 2009); data reduction: *CrysAlis RED*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3I* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Technical support (X-ray measurements at SCDRX) from Université Henry Poincaré, Nancy 1 is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2156).

References

- Altomare, A., Cascarano, G., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343–350.
- Brodersen, K. & Hummel, H.-U. (1987). In *Comprehensive Coordination Chemistry*, Vol. 5, edited by G. Wilkinson, R. D. Gillard & J. A. McCleverty, pp. 1047–1170. New York: Pergamon.
- Cassoux, P., Valade, L., Kobayashi, H., Kobayashi, A., Clark, R. A. & Underhill, A. E. (1991). *Coord. Chem. Rev.* pp. 110–115.
- Clark, R. C. & Reid, J. S. (1995). *Acta Cryst.* **A51**, 887–897.
- Dai, J., Kudora-Sowa, T., Munakata, M., Maekawa, M., Suenaga, Y. & Ohno, Y. (1997). *J. Chem. Soc. Dalton Trans.* pp. 2363–2368.
- Dai, J., Munakata, M., Kuroda-Sowa, T., Suenaga, Y., Wu, L. P. & Yamamoto, M. (1997). *Inorg. Chim. Acta*, **255**, 163–166.
- Dai, J., Munakata, M., Wu, L. P., Kuroda-Sowa, T. & Suenaga, Y. (1997). *Inorg. Chim. Acta*, **258**, 65–69.
- Dean, P. A. W. (1978). *Prog. Inorg. Chem.* **24**, 109–178.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Klinsberg, E. & Schraber, A. M. (1962). *J. Am. Chem. Soc.* **84**, 2941–2944.
- Oxford Diffraction (2009). *CrysAlis CCD* and *CrysAlis RED*. Oxford Diffraction, Wrocław, Poland.
- Pullen, A. E. & Oik, R.-M. (1999). *Coord. Chem. Rev.* pp. 118–211.
- Schlueter, J. A., Williams, J. M., Geiser, U., Wang, H. H., Kini, A. M., Kelly, M. E. & Dudek, J. D. (1996). *Mol. Cryst. Liq. Cryst.* **285**, 43–50.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m1080-m1081 [doi:10.1107/S1600536809032152]

***trans*-Di- μ -iodido-bis[(3*H*-1,2-benzodithiole-3-thione)iodidomercury(II)]**

E. A. Laifa, L. Bendjeddou, N. Boudraa, S. Dahaoui and C. Lecomte

Comment

Sulfur-rich organic donor-acceptor compounds or radical salts are widely used as bricks in the crystal architecture of molecular materials. The S...S interaction or contact in this compounds is one of the most important contributors to their unique electronic properties (Klinsberg *et al.*, 1962, Cassoux *et al.*, 1991). It has been understood that intermolecular S...S interactions are providing the pass way of the electrons in the molecular conductor. Up to now great efforts have been made to design the molecular packing and to strengthen the S...S contacts by modifying the structure of the organic molecule itself, by changing the size of the counter ion and even the co-crystallized solvent molecules (Pullen *et al.*, 1999; Schlueter *et al.*, 1996). In this context a series of polymeric complexes have been reported with (Ag⁺) metal (Dai, Munakata, Kuroda-Sowa *et al.*, 1997; Dai, Kudora-Sowa *et al.*, 1997) and a tetra-nuclear cluster CuI₄ (Dai, Munakata, Wu *et al.*, 1997). In this paper an organic-inorganic hybrid compound is reported with the formula: [Hg₂I₄(DTT)₂](1), DTT=C₇H₄S₃, 4,5-benzo-1,2-dithiole-3-thione.

The complex [Hg₂(C₁₄H₈S₆)I₄] is found to be a halogen-bridged dimer related by an inversion centre. An *ORTEP* view of the complex together with the atomic labeling scheme is given in Fig.1. The mercury atom is four-coordinated with significant distortion from tetrahedral. Two of these bonds are formed by two asymmetric iodide bridges, while the remaining two bonds are formed by a terminal iodide ion and thiocarbonyl sulfur atom of the ligand. The square-like plane formed by mercury and iodide atoms with their symmetry counterparts makes 89.66 (3)° dihedral angle with DDT plane.

The selected bond lengths and angles for the complex are listed in table (1) in which the data of the dimeric unit are comparable to those reported in the literature for related chloride bridged dimeric Hg(II) complexes (Brodersen *et al.*, 1987; Dean, 1978).

Coordination modes of the C₇H₄S₃ ligand have been classified into four types (Dai, Kudora-Sowa *et al.*, 1997). They are monodentate coordination by the thiocarbonyl group (type I), bridge formation by the sulfur of the thiocarbonyl group (type II), bidentate coordination by both thiocarbonyl group and thioether group (type III) and tridentate coordination by the thiocarbonyl sulfur (acting as a bridge) and the thioether sulfur (type IV).

Although the coordination of the ligand in type I has been found in complex: [Cu₄I₄(C₅H₄S₅)₄][∞] and {[Ag(C₅H₄S₅)₃]ClO₄CH₃CN}₂ (Dai, Kudora-Sowa *et al.*, 1997; Dai, Munakata, Wu *et al.*, 1997), other coordination types co-exist in these complexes.

[Hg₂(C₇H₄S₃)₂I₄] is the complex found in the coordination of the ligands only in type I. The S(2)—C(1) distance of 1.703 (2)Å in this compound is longer than that for the free ligand 1.645Å (Dai, Munakata, Kuroda-Sowa *et al.*, 1997). This longer S(2)—C(1) distance attributes to the strong coordination of the thiocarbonyl sulfur to soft mercury(II) ion that weakens the S(2)—C(1) bond.

supplementary materials

The absence of intermolecular hydrogen bonding shows that the molecules are retained to each other by Van der Waals interaction only. The second type of bonds arises from an S(2)—I(1) and S(3)—I(1) interaction with two different bond lengths, 3.4452 (7) Å and 3.6859 (7) Å respectively (Fig.2). These bonds maintain the cohesion of the crystalline structure.

Experimental

The reagent C₇H₄S₃ was prepared using a literature method (Klinsberg *et al.*, 1962) and characterized. The solvent was dried and distilled by a standard method before use. All other chemicals were obtained from commercial sources and used without further purification. Infrared spectra were measured as KBr disc on a Nicolet 205 FT—IR spectrometer.

A solution of HgI₂ (2.5 mmol, 1.130 g) in acetone (15 ml) was added to a solution of C₇H₄S₃ (2.5 mmol, 0.460 g) in acetone at room temperature under argon atmosphere. An orange precipitate was formed immediately and the mixture was stirred for 50 min. The precipitate was filtered, washed with petroleum ether (yield: 60%). Orange crystals for x-ray measurement were obtained by recrystallizing the solid in THF. IR (KBr, cm⁻¹): ν(C=C) 1447 ms, ν(C=S) 991,3vs, ν(Hg—S) 256,5 ms, ν(Hg—I) 167,8vs.

Refinement

H atoms were positioned geometrically and refined in the riding-model approximation, with C—H = 0.93 Å and with $U_{iso}(H) = 1.2 U_{eq}(C)$

Figures




Fig. 1. : An ORTEP view of the dimeric structure of (I) with the atom-labeling scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level. [Symmetry code: (a) -x, 1-y, 1-z]




Fig. 2. : A View of the structure compound showing the interactions between two adjacent dimer. Atoms marked with a hash symbol (#) and dollar sign (\$), are at the symmetry positions (x, 1 + y, z), (x, -1 + y, z) respectively.

trans-Di-μ-iodido-bis[(3*H*-1,2-benzodithiole-3- thione)iodidomercury(II)]

Crystal data

[Hg₂I₄(C₇H₄S₃)₂]

$M_r = 1277.40$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 7.86285 (10) \text{ \AA}$

$b = 8.19304 (11) \text{ \AA}$

$c = 10.46506 (13) \text{ \AA}$

$\alpha = 105.2917 (11)^\circ$

$\beta = 98.3031 (10)^\circ$

$Z = 1$

$F_{000} = 560$

$D_x = 3.483 \text{ Mg m}^{-3}$

Melting point: 192 K

Mo $K\alpha$ radiation, $\lambda = 0.7107 \text{ \AA}$

Cell parameters from 4951 reflections

$\theta = 2.8\text{--}35.0^\circ$

$\mu = 18.18 \text{ mm}^{-1}$

$T = 100 \text{ K}$

$\gamma = 105.6957$ (11) $^\circ$ Needle, orange
 $V = 608.917$ (15) \AA^3 $0.16 \times 0.1 \times 0.08$ mm

Data collection

Super Nova diffractometer (Dual, Cu at zero, Mo active) with an Atlas detector 4951 independent reflections
 Radiation source: Enhance (Mo) X-ray Source 4698 reflections with $I > 2\sigma(I)$
 Monochromator: graphite $R_{\text{int}} = 0.051$
 Detector resolution: 10.4508 pixels mm^{-1} $\theta_{\text{max}} = 34.0^\circ$
 $T = 100$ K $\theta_{\text{min}} = 3.2^\circ$
 ω scans $h = -12 \rightarrow 12$
 Absorption correction: analytical (Clark *et al.*, 1995) $k = -13 \rightarrow 13$
 $T_{\text{min}} = 0.167$, $T_{\text{max}} = 0.347$ $l = -16 \rightarrow 16$
 44929 measured reflections

Refinement

Refinement on F^2 H-atom parameters constrained
 Least-squares matrix: full $w = 1/[\sigma^2(F_o^2) + (0.0169P)^2 + 0.3949P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $R[F^2 > 2\sigma(F^2)] = 0.015$ $(\Delta/\sigma)_{\text{max}} = 0.002$
 $wR(F^2) = 0.037$ $\Delta\rho_{\text{max}} = 1.20 \text{ e \AA}^{-3}$
 $S = 1.05$ $\Delta\rho_{\text{min}} = -1.54 \text{ e \AA}^{-3}$
 4951 reflections Extinction correction: none
 118 parameters

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Hg1	0.176661 (11)	0.378520 (10)	0.439759 (7)	0.01756 (2)
I1	0.127512 (16)	0.692780 (15)	0.404010 (12)	0.01240 (3)
I2	0.072031 (19)	0.088861 (17)	0.221273 (13)	0.01654 (3)
S1	0.41650 (7)	0.51906 (6)	0.65918 (5)	0.01343 (8)

supplementary materials

C1	0.4481 (3)	0.3455 (3)	0.70715 (19)	0.0124 (3)
S2	0.33319 (7)	0.13144 (6)	0.60634 (5)	0.01541 (9)
S3	0.44554 (7)	0.00480 (7)	0.72353 (5)	0.01732 (9)
C7	0.6773 (3)	0.5278 (3)	0.92676 (19)	0.0145 (3)
H7	0.6733	0.6349	0.9138	0.017*
C2	0.5686 (3)	0.3630 (3)	0.82988 (19)	0.0117 (3)
C3	0.5762 (3)	0.2021 (3)	0.85055 (19)	0.0142 (3)
C5	0.7939 (3)	0.3662 (3)	1.0612 (2)	0.0185 (4)
H5	0.8692	0.369	1.1394	0.022*
C6	0.7895 (3)	0.5283 (3)	1.0409 (2)	0.0173 (4)
H6	0.8629	0.6363	1.1049	0.021*
C4	0.6894 (3)	0.2039 (3)	0.9680 (2)	0.0176 (4)
H4	0.6937	0.0976	0.9825	0.021*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Hg1	0.02112 (4)	0.01604 (4)	0.01403 (4)	0.00738 (3)	-0.00021 (3)	0.00324 (3)
I1	0.01305 (5)	0.01087 (5)	0.01336 (5)	0.00370 (4)	0.00296 (4)	0.00420 (4)
I2	0.01919 (6)	0.01330 (5)	0.01626 (6)	0.00601 (5)	0.00372 (4)	0.00271 (4)
S1	0.0141 (2)	0.01235 (18)	0.01343 (19)	0.00436 (16)	0.00122 (15)	0.00450 (15)
C1	0.0114 (7)	0.0132 (7)	0.0130 (7)	0.0045 (6)	0.0037 (6)	0.0038 (6)
S2	0.0157 (2)	0.01270 (18)	0.0152 (2)	0.00376 (17)	-0.00177 (16)	0.00373 (16)
S3	0.0200 (2)	0.01284 (19)	0.0179 (2)	0.00602 (18)	-0.00073 (17)	0.00482 (17)
C7	0.0142 (8)	0.0160 (8)	0.0127 (8)	0.0052 (7)	0.0022 (6)	0.0037 (7)
C2	0.0104 (7)	0.0145 (7)	0.0110 (7)	0.0053 (6)	0.0026 (6)	0.0039 (6)
C3	0.0155 (8)	0.0166 (8)	0.0123 (7)	0.0076 (7)	0.0038 (6)	0.0050 (6)
C5	0.0187 (9)	0.0244 (9)	0.0135 (8)	0.0110 (8)	0.0014 (7)	0.0049 (7)
C6	0.0155 (9)	0.0198 (9)	0.0138 (8)	0.0054 (7)	0.0007 (7)	0.0026 (7)
C4	0.0200 (9)	0.0201 (9)	0.0143 (8)	0.0098 (8)	0.0025 (7)	0.0055 (7)

Geometric parameters (\AA , $^\circ$)

Hg1—S1	2.5169 (7)	C7—C6	1.377 (3)
Hg1—I2	2.6599 (2)	C7—C2	1.409 (3)
Hg1—I1	2.8148 (4)	C7—H7	0.93
Hg1—I1 ⁱ	3.1002 (4)	C2—C3	1.406 (3)
I1—Hg1 ⁱ	3.1002 (4)	C3—C4	1.404 (3)
S1—C1	1.6935 (19)	C5—C4	1.373 (3)
C1—C2	1.430 (3)	C5—C6	1.408 (3)
C1—S2	1.702 (2)	C5—H5	0.93
S2—S3	2.0532 (7)	C6—H6	0.93
S3—C3	1.734 (2)	C4—H4	0.93
S1—Hg1—I2	138.842 (15)	C3—C2—C7	119.89 (17)
S1—Hg1—I1	98.374 (17)	C3—C2—C1	115.91 (17)
I2—Hg1—I1	117.574 (11)	C7—C2—C1	124.19 (17)
S1—Hg1—I1 ⁱ	91.326 (17)	C4—C3—C2	120.60 (18)
I2—Hg1—I1 ⁱ	103.851 (13)	C4—C3—S3	122.18 (16)

I1—Hg1—I1 ⁱ	95.210 (10)	C2—C3—S3	117.21 (14)
Hg1—I1—Hg1 ⁱ	84.790 (10)	C4—C5—C6	121.54 (18)
C1—S1—Hg1	105.26 (7)	C4—C5—H5	119.2
C2—C1—S1	124.72 (15)	C6—C5—H5	119.2
C2—C1—S2	115.08 (14)	C7—C6—C5	120.41 (19)
S1—C1—S2	120.20 (11)	C7—C6—H6	119.8
C1—S2—S3	97.66 (7)	C5—C6—H6	119.8
C3—S3—S2	94.06 (7)	C5—C4—C3	118.47 (19)
C6—C7—C2	119.08 (18)	C5—C4—H4	120.8
C6—C7—H7	120.5	C3—C4—H4	120.8
C2—C7—H7	120.5		

Symmetry codes: (i) $-x, -y+1, -z+1$.

Fig. 1

