

catena-Poly[[[tetraaquazinc(II)]- μ -4,4'-bipyridine- κ^2 N:N'] naphthalene-1,5-disulfonate]

Jing Lin^a and Wen-Lian Cai^{b*}

^aDepartment of Pharmacy, Xiamen University, Xiamen, Fujian 363105, People's Republic of China, and ^bDepartment of Chemistry and Environmental Science, Zhangzhou Normal University, Zhangzhou, Fujian 363000, People's Republic of China

Correspondence e-mail: ghx919@yahoo.com.cn


Received 12 August 2009; accepted 13 August 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.034; wR factor = 0.107; data-to-parameter ratio = 13.3.

In the title complex, $\{[Zn(C_{10}H_8N_2)(H_2O)_4](C_{10}H_6O_6S_2)\}_n$, the $[Zn(4,4'\text{-bipy})(H_2O)_4]^{2+}$ ($4,4'$ -bipy is $4,4'$ -bipyridine) cations are linked into linear chains along [001] by the $4,4'$ -bipy ligands. The Zn^{II} ion exhibits a slightly distorted octahedral coordination geometry in which the four water molecules are in the equatorial positions. The anions are hydrogen bonded to the polycationic chains by $O-H\cdots O$ hydrogen bonds, forming a three-dimensional network. The Zn^{II} ion, $4,4'$ -bipy ligand and anion lie on special positions of $2/m$ site symmetry.

Related literature

For the design, preparation and applications of metal-organic hybrid materials, see: Batten & Robson (1998); Hagrman *et al.* (1999); Cui *et al.* (2003). For the structural and photoluminescent properties of d^{10} metal (such as Zn) complexes, see: Li *et al.* (2003); Sattarzadeh *et al.* (2009). $4,4'$ -Bipyridine can be used to assemble many transition metal coordination polymers through covalent or hydrogen bonds, see: Yaghi & Li (1995, 1996).



Experimental

Crystal data

$[Zn(C_{10}H_8N_2)(H_2O)_4](C_{10}H_6O_6S_2)$

$M_r = 579.89$

Monoclinic, $C2/m$
 $a = 14.584$ (3) Å
 $b = 7.3948$ (15) Å
 $c = 11.380$ (2) Å
 $\beta = 108.38$ (3) $^\circ$
 $V = 1164.7$ (4) Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.29$ mm⁻¹
 $T = 293$ K
 $0.38 \times 0.29 \times 0.19$ mm

Data collection

Siemens SMART CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{min} = 0.658$, $T_{max} = 0.794$

5711 measured reflections
 1421 independent reflections
 1302 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.107$
 $S = 1.03$
 1421 reflections
 107 parameters
 15 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.46$ e Å⁻³
 $\Delta\rho_{\min} = -0.52$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1w-H1wa-O3	0.85 (2)	1.92 (2)	2.763 (3)	175 (3)
O1w-H1wb-O2 ⁱ	0.83 (2)	1.95 (2)	2.768 (3)	166 (3)

Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + 1$.

Data collection: *SMART* (Siemens, 1994); cell refinement: *SAINT* (Siemens, 1994); data reduction: *SAINT* program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*

This work was supported by the Natural Science Foundation of Fujian Province (No. 2008 J0172).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2626).

References

- Batten, S. R. & Robson, R. (1998). *Angew. Chem. Int. Ed.* **37**, 1460–1494.
- Cui, Y., Ngo, H. L., White, P. S. & Lin, W. (2003). *Inorg. Chem.* **42**, 652–654.
- Hagrman, P. J., Hagrman, D. & Zubietta, J. (1999). *Angew. Chem. Int. Ed.* **38**, 2638–2684.
- Li, R. Z., Li, D., Huang, X. C., Qi, Z. Y. & Chen, X. M. (2003). *Inorg. Chem. Commun.* **6**, 1017–1019.
- Sattarzadeh, E., Mohammadnezhad, G., Amini, M. M. & Ng, S. W. (2009). *Acta Cryst. E* **65**, m712–m713.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1994). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Yaghi, O. M. & Li, H. (1995). *J. Am. Chem. Soc.* **117**, 10401–10402.
- Yaghi, O. M. & Li, H. (1996). *J. Am. Chem. Soc.* **118**, 295–296.

supplementary materials

Acta Cryst. (2009). E65, m1088 [doi:10.1107/S1600536809032127]

[*catena-Poly[[[tetraaquazinc(II)]- μ -4,4'-bipyridine- κ^2 N:N'] naphthalene-1,5-disulfonate*]

J. Lin and W.-L. Cai

Comment

The design and preparation of metal-organic hybrid materials have been studied widely during the past decade owing to their intriguing structures and potential practical applications. (Hagrman *et al.*, 1999; Batten & Robson, 1998; Cui *et al.*, 2003). It has been demonstrated that many d¹⁰ metal(such as Zn) complexes present intriguing structural and photoluminescent properties (Li *et al.*, 2003; Sattarzadeh *et al.*, 2009). On the other hand, bridged ligand,4,4'-bipyridine, can be efficiently used to assembly many interesting transition metal coordination polymers through covalent or hydrogen bonds (Yaghi & Li, 1995, 1996). In this work, we use 4,4'-bpy, 1, 5-naphthalenedisulphonic acid(NDS) and Zn(OAc)₂ to synthesize a novel 1-D chain polymer through hydrothermal synthesis.

Complex (I) consists of one-dimensional chains formed by 4,4'-bipy ligands through connecting Zn atoms, uncoordinated NDS²⁻ anions, as shown in Fig. 1. The [Zn(4,4'-bipy)(H₂O)₄]²⁺ cation is located on a twofold rotation axis that passes through atoms Zn1, N1 and C3. In the cation, the Zn1 atom exhibits slightly distorted octahedral coordination geometry, completed by four O atoms from four water molecules in the equatorial positions and two N donors from two 4,4'-bipy ligands in the apical positions. These Zn–O and Zn–N distances are 2.127 (2) and 2.131 (2) Å, respectively. The 4,4'-bipy ligand acts as bis-monodentate linkers and bridge adjacent Zn centers with the Zn···Zn separation of 11.380 (2) Å into an infinite one-dimensional chain.

In complex (I), the NDS²⁻ anions are not involved in coordination but hydrogen-bonded to the [Zn(4,4'-bipy)(H₂O)₄]²⁺ cations through the sulfonate oxygen atoms and the coordinated oxygen atoms, forming a three-dimensional hydrogen-bonding network, as shown in Fig. 2.

Experimental

The hydrothermal reaction of Zn(OAc)₂·2H₂O (0.5707 g, 2.6 mmol), 1,5-Naphthalenedisulphonic acid(0.5405 g, 1.5 mmol), 4,4'-bipyridine(0.4681 g, 3.0 mmol) and water (15 ml) was carried out at 443 K for 3 d. After cooling to room temperature at 5 K h⁻¹, the colorless block crystalline complex, (I), was isolated in 49% yield (based on Zn).

Refinement

H atoms attached to C atoms were positioned geometrically and refined using a riding model, with C–H = 0.93 Å, and U_{iso}(H)= 1.2U_{eq}(C); Water H atoms were located in a difference map and refined with O–H and H···H distance restraints of 0.85 (2) and 1.39 (2) Å, respectively, and with U_{iso}(H)= 1.5U_{eq}(O). During the refinement, the displacement parameters of the C1 and C2 atoms were restrained to an approximately isotropic behaviour.

supplementary materials

Figures




Fig. 1. View of the structure of compound (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 35% probability level; H atoms have been omitted for clarity.




Fig. 2. View of the 3D hydrogen-bonded network in the packing of the title compound. The packing is viewed along the b axis; O-H \cdots O interactions are shown as dashed lines.

catena-Poly[[[tetraaquazinc(II)]- μ -4,4'-bipyridine- κ^2 N:N'] naphthalene-1,5-disulfonate]

Crystal data

[Zn(C ₁₀ H ₈ N ₂)(H ₂ O) ₄](C ₁₀ H ₆ O ₆ S ₂)	$F_{000} = 596$
$M_r = 579.89$	$D_x = 1.654 \text{ Mg m}^{-3}$
Monoclinic, $C2/m$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -C 2y	Cell parameters from 5711 reflections
$a = 14.584 (3) \text{ \AA}$	$\theta = 3.1\text{--}27.4^\circ$
$b = 7.3948 (15) \text{ \AA}$	$\mu = 1.29 \text{ mm}^{-1}$
$c = 11.380 (2) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 108.38 (3)^\circ$	Block, colorless
$V = 1164.7 (4) \text{ \AA}^3$	$0.38 \times 0.29 \times 0.19 \text{ mm}$
$Z = 2$	

Data collection

Siemens SMART CCD area-detector diffractometer	1421 independent reflections
Radiation source: fine-focus sealed tube	1302 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.033$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 27.4^\circ$
ω scans	$\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -18 \rightarrow 18$
$T_{\text{min}} = 0.658$, $T_{\text{max}} = 0.794$	$k = -9 \rightarrow 9$
5711 measured reflections	$l = -13 \rightarrow 14$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0663P)^2 + 1.4457P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\max} = 0.001$
1421 reflections	$\Delta\rho_{\max} = 0.46 \text{ e \AA}^{-3}$
107 parameters	$\Delta\rho_{\min} = -0.52 \text{ e \AA}^{-3}$
15 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.5000	0.5000	0.0328 (2)
S1	0.84039 (6)	0.5000	0.70300 (8)	0.0323 (3)
O1W	0.60582 (14)	0.7084 (3)	0.54834 (19)	0.0443 (5)
H1WA	0.6631 (16)	0.698 (5)	0.596 (2)	0.053*
H1WB	0.605 (2)	0.790 (4)	0.498 (3)	0.053*
O2	0.8672 (2)	0.5000	0.5896 (2)	0.0451 (7)
O3	0.78840 (14)	0.6629 (3)	0.71577 (18)	0.0425 (5)
C1	0.5820 (3)	0.5000	0.7822 (4)	0.0669 (15)
H1A	0.6403	0.5000	0.7651	0.080*
C2	0.5848 (3)	0.5000	0.9042 (4)	0.0647 (14)
H2A	0.6440	0.5000	0.9667	0.078*
C3	0.4997 (3)	0.5000	0.9346 (3)	0.0324 (8)
C4	0.4162 (3)	0.5000	0.8359 (3)	0.0364 (8)
H4A	0.3568	0.5000	0.8501	0.044*
C5	0.4189 (3)	0.5000	0.7155 (3)	0.0337 (8)
H5A	0.3607	0.5000	0.6511	0.040*
C6	0.8693 (3)	0.5000	0.9887 (4)	0.0580 (14)
H6A	0.8090	0.5000	0.9283	0.070*
C7	0.9543 (2)	0.5000	0.9528 (3)	0.0329 (8)
C8	0.9520 (3)	0.5000	0.8266 (3)	0.0338 (8)

supplementary materials

C9	1.0349 (3)	0.5000	0.7957 (4)	0.0549 (13)
H9A	1.0322	0.5000	0.7130	0.066*
C10	1.1247 (3)	0.5000	0.8898 (5)	0.080 (2)
H10A	1.1812	0.5000	0.8684	0.096*
N1	0.5005 (2)	0.5000	0.6874 (3)	0.0371 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.0270 (3)	0.0560 (4)	0.0140 (3)	0.000	0.0046 (2)	0.000
S1	0.0278 (5)	0.0414 (5)	0.0202 (4)	0.000	-0.0030 (3)	0.000
O1W	0.0347 (10)	0.0589 (13)	0.0301 (10)	-0.0062 (9)	-0.0031 (8)	0.0086 (9)
O2	0.0468 (16)	0.0607 (18)	0.0213 (13)	0.000	0.0014 (12)	0.000
O3	0.0360 (10)	0.0440 (11)	0.0372 (10)	0.0048 (8)	-0.0033 (8)	-0.0028 (8)
C1	0.036 (2)	0.137 (4)	0.030 (2)	0.000	0.0134 (18)	0.000
C2	0.032 (2)	0.135 (4)	0.027 (2)	0.000	0.0088 (17)	0.000
C3	0.0311 (18)	0.048 (2)	0.0190 (17)	0.000	0.0089 (14)	0.000
C4	0.0294 (17)	0.058 (2)	0.0230 (17)	0.000	0.0106 (14)	0.000
C5	0.0302 (17)	0.050 (2)	0.0186 (16)	0.000	0.0044 (13)	0.000
C6	0.0194 (17)	0.120 (5)	0.029 (2)	0.000	-0.0003 (15)	0.000
C7	0.0241 (17)	0.046 (2)	0.0247 (17)	0.000	0.0027 (15)	0.000
C8	0.0244 (16)	0.049 (2)	0.0224 (16)	0.000	-0.0008 (13)	0.000
C9	0.034 (2)	0.106 (4)	0.0219 (18)	0.000	0.0048 (16)	0.000
C10	0.025 (2)	0.178 (7)	0.036 (3)	0.000	0.0093 (19)	0.000
N1	0.0307 (15)	0.065 (2)	0.0157 (13)	0.000	0.0072 (12)	0.000

Geometric parameters (\AA , $^\circ$)

Zn1—O1W ⁱ	2.127 (2)	C3—C4	1.372 (5)
Zn1—O1W	2.127 (2)	C3—C3 ^{iv}	1.485 (6)
Zn1—O1W ⁱⁱ	2.127 (2)	C4—C5	1.383 (5)
Zn1—O1W ⁱⁱⁱ	2.127 (2)	C4—H4A	0.9300
Zn1—N1 ⁱ	2.131 (3)	C5—N1	1.326 (5)
Zn1—N1	2.131 (3)	C5—H5A	0.9300
S1—O3 ⁱⁱ	1.455 (2)	C6—C10 ^v	1.358 (6)
S1—O3	1.455 (2)	C6—C7	1.421 (5)
S1—O2	1.461 (3)	C6—H6A	0.9300
S1—C8	1.784 (4)	C7—C7 ^v	1.424 (7)
O1W—H1WA	0.846 (19)	C7—C8	1.426 (5)
O1W—H1WB	0.83 (2)	C8—C9	1.362 (6)
C1—N1	1.330 (6)	C9—C10	1.406 (6)
C1—C2	1.376 (6)	C9—H9A	0.9300
C1—H1A	0.9300	C10—C6 ^v	1.358 (6)
C2—C3	1.390 (6)	C10—H10A	0.9300
C2—H2A	0.9300		
O1W ⁱ —Zn1—O1W	180.00 (9)	C3—C2—H2A	119.8
O1W ⁱ —Zn1—O1W ⁱⁱ	87.13 (12)	C4—C3—C2	115.3 (3)

O1W—Zn1—O1W ⁱⁱ	92.87 (12)	C4—C3—C3 ^{iv}	123.0 (4)
O1W ⁱ —Zn1—O1W ⁱⁱⁱ	92.87 (12)	C2—C3—C3 ^{iv}	121.7 (4)
O1W—Zn1—O1W ⁱⁱⁱ	87.13 (12)	C3—C4—C5	121.1 (3)
O1W ⁱⁱ —Zn1—O1W ⁱⁱⁱ	180.000 (1)	C3—C4—H4A	119.4
O1W ⁱ —Zn1—N1 ⁱ	88.18 (8)	C5—C4—H4A	119.4
O1W—Zn1—N1 ⁱ	91.82 (8)	N1—C5—C4	123.1 (3)
O1W ⁱⁱ —Zn1—N1 ⁱ	91.82 (8)	N1—C5—H5A	118.4
O1W ⁱⁱⁱ —Zn1—N1 ⁱ	88.18 (8)	C4—C5—H5A	118.4
O1W ⁱ —Zn1—N1	91.82 (8)	C10 ^v —C6—C7	120.7 (4)
O1W—Zn1—N1	88.18 (8)	C10 ^v —C6—H6A	119.7
O1W ⁱⁱ —Zn1—N1	88.18 (8)	C7—C6—H6A	119.7
O1W ⁱⁱⁱ —Zn1—N1	91.82 (8)	C6—C7—C7 ^v	118.6 (4)
N1 ⁱ —Zn1—N1	180.0	C6—C7—C8	122.9 (3)
O3 ⁱⁱ —S1—O3	111.78 (18)	C7 ^v —C7—C8	118.5 (4)
O3 ⁱⁱ —S1—O2	112.40 (11)	C9—C8—C7	121.3 (3)
O3—S1—O2	112.40 (11)	C9—C8—S1	117.4 (3)
O3 ⁱⁱ —S1—C8	107.20 (10)	C7—C8—S1	121.3 (3)
O3—S1—C8	107.20 (10)	C8—C9—C10	119.6 (4)
O2—S1—C8	105.36 (17)	C8—C9—H9A	120.2
Zn1—O1W—H1WA	126 (3)	C10—C9—H9A	120.2
Zn1—O1W—H1WB	120 (2)	C6 ^v —C10—C9	121.3 (4)
H1WA—O1W—H1WB	108 (3)	C6 ^v —C10—H10A	119.3
N1—C1—C2	123.5 (4)	C9—C10—H10A	119.3
N1—C1—H1A	118.2	C5—N1—C1	116.5 (3)
C2—C1—H1A	118.2	C5—N1—Zn1	121.4 (2)
C1—C2—C3	120.5 (4)	C1—N1—Zn1	122.1 (3)
C1—C2—H2A	119.8		
N1—C1—C2—C3	0.000 (2)	C7—C8—C9—C10	0.000 (2)
C1—C2—C3—C4	0.000 (2)	S1—C8—C9—C10	180.000 (2)
C1—C2—C3—C3 ^{iv}	180.000 (2)	C8—C9—C10—C6 ^v	0.000 (2)
C2—C3—C4—C5	0.000 (1)	C4—C5—N1—C1	0.000 (2)
C3 ^{iv} —C3—C4—C5	180.000 (1)	C4—C5—N1—Zn1	180.000 (1)
C3—C4—C5—N1	0.000 (2)	C2—C1—N1—C5	0.000 (1)
C10 ^v —C6—C7—C7 ^v	0.000 (2)	C2—C1—N1—Zn1	180.000 (1)
C10 ^v —C6—C7—C8	180.000 (2)	O1W ⁱ —Zn1—N1—C5	-46.47 (6)
C6—C7—C8—C9	180.000 (2)	O1W—Zn1—N1—C5	133.53 (6)
C7 ^v —C7—C8—C9	0.000 (2)	O1W ⁱⁱ —Zn1—N1—C5	-133.53 (6)
C6—C7—C8—S1	0.000 (1)	O1W ⁱⁱⁱ —Zn1—N1—C5	46.47 (6)
C7 ^v —C7—C8—S1	180.000 (1)	N1 ⁱ —Zn1—N1—C5	0(100)
O3 ⁱⁱ —S1—C8—C9	119.92 (10)	O1W ⁱ —Zn1—N1—C1	133.53 (6)
O3—S1—C8—C9	-119.92 (10)	O1W—Zn1—N1—C1	-46.47 (6)
O2—S1—C8—C9	0.0	O1W ⁱⁱ —Zn1—N1—C1	46.47 (6)
O3 ⁱⁱ —S1—C8—C7	-60.08 (10)	O1W ⁱⁱⁱ —Zn1—N1—C1	-133.53 (6)

supplementary materials

O3—S1—C8—C7	60.08 (10)	N1 ⁱ —Zn1—N1—C1	180 (100)
O2—S1—C8—C7	180.000 (1)		


Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x, -y+1, z$; (iii) $-x+1, y, -z+1$; (iv) $-x+1, -y+1, -z+2$; (v) $-x+2, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1w—H1wa···O3	0.85 (2)	1.92 (2)	2.763 (3)	175 (3)
O1w—H1wb···O2 ^{vi}	0.83 (2)	1.95 (2)	2.768 (3)	166 (3)

Symmetry codes: (vi) $-x+3/2, y+1/2, -z+1$.

Fig. 1



supplementary materials

Fig. 2

