

***trans*-(Pyrimidine-2-thiolato- κ^2N,S)-
[tris(2-aminoethyl)amine-
 κ^4N,N',N'',N''']cobalt(III) chloride
hexafluoridophosphate**

Keisuke Fujihara and Toshiaki Yonemura*

Department of Applied Science, Faculty of Science, Kochi University, Akebono-cho, Kochi 780-8520, Japan

Correspondence e-mail: yonemura@kochi-u.ac.jp


Received 15 August 2009; accepted 15 August 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.006$ Å; R factor = 0.057; wR factor = 0.174; data-to-parameter ratio = 18.3.

In the title compound, $[Co(C_4H_3N_2S)(C_6H_{18}N_4)](Cl)PF_6$, the Co^{III} ion is coordinated by a tripod-like tetradentate ligand and a monoanionic N,S -bidentate ligand in an approximately octahedral CoN_4OS geometry. The anionic S atom of the pyrimidine-2-thiolate (pymt) ligand is coordinated in the *trans* position to the primary amine N atom (N_{prim}) of the tris(2-aminoethyl)amine (tren) ligand. The crystal structure exhibits short intermolecular $N-H\cdots N$ hydrogen bonds ($N\cdots N < 3.2$ Å), and intermolecular $N-H\cdots Cl$ and $C-H\cdots F$ contacts, leading to the formation of an infinite two-dimensional network.

Related literature

For the synthesis and chemistry of similar tren [tren = tris(2-aminoethyl)amine] complexes, see: Jackson & Sargeson (1978); Kojima *et al.* (1994); Mitsui *et al.* (1976); Ohba & Saito (1984); Okamoto *et al.* (1990); Yonemura *et al.* (1997).



Experimental

Crystal data

$[Co(C_4H_3N_2S)(C_6H_{18}N_4)](Cl)PF_6$ $V = 1888.7$ (5) Å³
 $M_r = 496.73$ $Z = 4$
 Monoclinic, $P2_1/c$ $Mo K\alpha$ radiation
 $a = 12.7106$ (17) Å $\mu = 1.31$ mm⁻¹
 $b = 11.326$ (2) Å $T = 296$ K
 $c = 14.205$ (2) Å $0.45 \times 0.35 \times 0.20$ mm
 $\beta = 112.549$ (10)°

Data collection

Rigaku AFC-7S diffractometer 3416 reflections with $F^2 > 2\sigma(F^2)$
 Absorption correction: ψ scan $R_{int} = 0.031$
 (North *et al.*, 1968) 3 standard reflections
 $T_{min} = 0.503$, $T_{max} = 0.770$ every 150 reflections
 5201 measured reflections intensity decay: 5.4%
 4342 independent reflections

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$ 237 parameters
 $wR(F^2) = 0.174$ H-atom parameters constrained
 $S = 1.05$ $\Delta\rho_{max} = 1.14$ e Å⁻³
 4342 reflections $\Delta\rho_{min} = -0.87$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N4-H8\cdots N2^i$	0.90	2.50	3.079 (4)	123
$N4-H9\cdots Cl1^{ii}$	0.90	2.52	3.278 (3)	142
$N5-H14\cdots Cl1^{iii}$	0.90	2.39	3.287 (3)	173
$N5-H15\cdots Cl1^{iii}$	0.90	2.42	3.273 (3)	160
$N6-H21\cdots Cl1$	0.90	2.29	3.187 (3)	173
$C6-H6\cdots F1$	0.97	2.39	3.210 (6)	143
$C7-H11\cdots F4^{iv}$	0.97	2.52	3.328 (6)	141
$C9-H17\cdots F6^v$	0.97	2.52	3.454 (6)	161

Symmetry codes: (i) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iv) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$; (v) $-x + 1, -y + 1, -z + 1$.

Data collection: *WinAFC* (Rigaku/MSK, 2000); cell refinement: *WinAFC*; data reduction: *CrystalStructure* (Rigaku/MSK, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *CrystalStructure*.

This work was partially supported by Grants-in-Aid for Scientific Research C (No. 20550138) from the Japanese Society for the Promotion of Science (JSPS). The authors are grateful to Kochi University for financial support (The Kochi University President's Discretionary Grant 2009).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2138).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Jackson, W. G. & Sargeson, A. M. (1978). *Inorg. Chem.* **17**, 2165–2169.
 Kojima, M., Shimizu, Y., Nakajima, K. & Yoshikawa, Y. (1994). *Bull. Chem. Soc. Jpn.* **67**, 869–872.
 Mitsui, Y., Watanabe, J., Harada, Y., Sakamaki, T., Kushi, Y. & Kimura, E. (1976). *J. Chem. Soc. Dalton Trans.* pp. 2095–2102.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.

metal-organic compounds

- Ohba, S. & Saito, Y. (1984). *Acta Cryst.* **C40**, 398–401.
- Okamoto, K., Konno, T. & Hidaka, J. (1990). *Acta Cryst.* **C46**, 216–218.
- Rigaku/MS (2000). *WinAFC*. Rigaku/MS, The Woodlands, Texas, USA.
- Rigaku/MS (2007). *CrystalStructure*. Rigaku/MS, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yonemura, T., Shibuya, K., Okamoto, K., Ama, T., Kawaguchi, H. & Yasui, T. (1997). *Inorg. Chim. Acta*, **260**, 119–128.

supplementary materials

Acta Cryst. (2009). E65, m1127-m1128 [doi:10.1107/S1600536809032395]

***trans*-(Pyrimidine-2-thiolato- κ^2N,S)[tris(2-aminoethyl)amine- κ^4N,N',N'',N''']cobalt(III) chloride hexafluoridophosphate**

K. Fujihara and T. Yonemura

Comment

Cobalt(III)-tren [tren = tris(2-aminoethyl)amine] complexes with thiolate and/or thioether ligands have been investigated in view of their stereochemistry and ligand substitution reactions (Mitsui & Kimura, *et al.*, 1976; Jackson & Sargeson *et al.*, 1978; Ohba & Saito, *et al.*, 1984; Okamoto, *et al.*, 1990; Kojima, *et al.*, 1994; Yonemura, *et al.*, 1997). The chemistry of the aliphatic and aromatic thiolato cobalt(III) complexes has led to many interesting results concerning the features of the coordinated sulfur atoms. Although aliphatic thiolato cobalt(III)-tren complexes have been extensively investigated, aromatic thiolato complexes have not been thus far because of the difficulties involved in their preparation.

In view of our interest in the stereochemistry and spectrochemical properties of aromatic and aliphatic cobalt(III) complexes, we synthesized the title compound using the tripod-like tetradentate tren ligand to fix the remaining two *cis* coordination sites in the cobalt(III) complexes, as two geometrical isomers, *p*-(*trans*(*Nprim,S*)) and *t*-(*trans*(*Ntert,S*)), are possible for [Co(bidentate-*N,S*)(tren)]-type complexes. The known cobalt(III)-tren complexes of 2-aminoethanethiolate-*N,S* (aet) (Sargeson *et al.*, 1978) formed *p*-isomers as the major products, and those of 2-mercaptoacetate-*O,S* (ma) or 3-mercaptopropionate-*O,S* (mp) selectively formed the *t*-isomers (Mitsui & Kimura, *et al.*, 1976; Jackson & Sargeson *et al.*, 1978; Ohba & Saito, *et al.*, 1984; Okamoto, *et al.*, 1990; Kojima, *et al.*, 1994; Yonemura, *et al.*, 1997). Herein we report on the structure of a cobalt(III)-tren complex coordinated with the aromatic thiolato ligand pyrimidine-2-thiolate (pymt).

In the title compound the cobalt(III) atom is coordinated by a tripod-like tetradentate ligand and a monoanionic *N,S*-bidentate ligand, producing an approximately octahedral CoN₄OS geometry (Fig. 1). In the present complex the anionic sulfur atom of the pymt ligand is coordinated *trans* to the primary amine N-atom (*Nprim*) of the tren ligand, with a nearly linear S1—Co1—N4 angle (170.71 (9)°) hence the title compound is the *p*-isomer (*trans*(*Nprim,S*)). The known cobalt(III)-tren complex of aet shows the same systematic trend, at least in the coordination manner of the *N,S*-bidentate ligand. The Co—S distance (2.3187 (12) Å) is significantly longer than those observed in the Co(III)-tren complexes with aliphatic thiolato ligands: 2.239 (1) Å in *t*-[Co{CH₃SCH(CH₃)COO}(tren)]²⁺ (Ohba & Saito, *et al.*, 1984), 2.236 (2) Å in *t*-[Co(mp)(tren)]⁺ and 2.232 (1) Å in *t*-[Co(ma)(tren)]⁺ (Yonemura, *et al.*, 1997). The Co—N4 distance (1.954 (3) Å), which is in the *trans* position relative to the sulfur atom, is shorter than the other Co-N(tren) distances (Co1—N5 = 1.966 (3) Å and Co1—N6 = 1.970 (3) Å). The S1-Co1-N1 angle of 72.31 (9)° is far from the ideal angle of 90°. This distortion is relaxed by the other angles in the same plane, that is, S1-Co1-N3 = 101.96 (9)°, and N1-Co1-N4 = 98.40 (12)°. The other bond distances and angles are similar to those in the cobalt(III)-tren complexes mentioned above.

The aromatic pymt *N,S*-chelate ring is almost planar and the three *N,N'*-chelate rings of the tren are in a *gauche* conformation. The *gauche* conformations of the two —N—CH₂—CH₂—NH₂ chelates in the tren ligand have unsymmetrical skew forms with the λ and δ conformations, and the *gauche* conformation of the central —N—CH₂—CH₂—NH₂ chelate in the tren ligand has an unsymmetrical skew form with the λ conformation.

supplementary materials

In the crystal structure each complex cation is linked to adjacent cations through N–H⋯N hydrogen bonds so constructing a one-dimensional chain structure (Fig. 2 and Table 1). These chains are further bridged by ordered Cl and PF₆ anions through N–H⋯Cl and C–H⋯F contacts, leading to the formation of an infinite two-dimensional network (Table 1 and Fig. 2).

Experimental

A methanolic solution of 2-pyrimidinethiol (1.12 g, 10 mmol) was adjusted to pH 8.0 with aqueous solutions of NaOH and HCl. The mixed solution was added to an aqueous solution of [CoCl₂(tren)]Cl (3.11 g, 10 mmol) and stirred at 40 °C for 1 h. The reaction mixture was poured onto an SP-Sepahadex C-25 column (Na⁺ form, 7 cm × 7 cm) and the adsorbed band was developed with a 0.2 mol dm⁻³ NaCl aqueous solution. The eluted red band was concentrated to a small volume, the precipitated NaCl was filtered off and the red filtrate was added to a solution of NH₄PF₆ (1.1 g, 6.7 mmol). The resulting red powder was collected by filtration and recrystallized from water. A few days later red crystals of the title compound were obtained with a yield of 1.50 g (35%). Anal. Calc. for [Co(pytm)(tren)]ClPF₆ = C₁₀H₂₁CoN₆SClPF₆: C 24.18, H 4.26, N 16.92%, Found: C 24.09, H 4.24, N 17.04%.

Refinement

The H-atoms were included in calculated positions and treated as riding atoms: N—H = 0.90 Å, C—H = 0.93–0.97 Å, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$.

Figures




Fig. 1. A view of the molecular structure of the title compound, showing the atom-labelling scheme and 50% probability displacement ellipsoids.




Fig. 2. A view along the b-axis of the crystal packing diagram of the title compound, showing the N–H⋯N hydrogen-bonds and N–H⋯Cl and C–H⋯F contacts as dashed lines (the H-atoms have been omitted for clarity).

trans-(Pyrimidine-2-thiolato- $\kappa^2 N, S$)[tris(2-aminoethyl)amine- $\kappa^4 N, N', N'', N'''$]cobalt(III) chloride hexafluoridophosphate

Crystal data

[Co(C₄H₃N₂S)(C₆H₁₈N₄)](Cl)PF₆

$F_{000} = 1008.0$

$M_r = 496.73$
 Monoclinic, $P2_1/c$
 Hall symbol: -P 2ybc
 $a = 12.7106$ (17) Å
 $b = 11.326$ (2) Å
 $c = 14.205$ (2) Å
 $\beta = 112.549$ (10)°
 $V = 1888.7$ (5) Å³
 $Z = 4$

$D_x = 1.747$ Mg m⁻³
 Mo $K\alpha$ radiation, $\lambda = 0.71069$ Å
 Cell parameters from 25 reflections
 $\theta = 15.4$ – 17.1 °
 $\mu = 1.31$ mm⁻¹
 $T = 296$ K
 Prismatic, red-brown
 $0.45 \times 0.35 \times 0.20$ mm

Data collection

Rigaku AFC-7S
 diffractometer
 $T = 296$ K
 ω - 2θ scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.503$, $T_{\max} = 0.770$
 5201 measured reflections
 4342 independent reflections
 3416 reflections with $F^2 > 2\sigma(F^2)$

$R_{\text{int}} = 0.031$
 $\theta_{\text{max}} = 27.5$ °
 $h = -16 \rightarrow 15$
 $k = -14 \rightarrow 0$
 $l = -10 \rightarrow 18$
 3 standard reflections
 every 150 reflections
 intensity decay: 5.4%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.174$
 $S = 1.05$
 4342 reflections
 237 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1106P)^2 + 1.816P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 1.14$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.87$ e Å⁻³
 Extinction correction: SHELXL97 (Sheldrick, 2008)
 Extinction coefficient: 0.0077 (13)

Special details

Geometry. Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell esds are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement was performed using all reflections. The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R -factor (gt) are based on F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	0.18778 (4)	0.70419 (4)	0.19018 (3)	0.0308 (1)

supplementary materials

S1	0.23158 (8)	0.80886 (9)	0.06968 (7)	0.0410 (3)
N1	0.0476 (2)	0.7685 (3)	0.0930 (2)	0.0372 (9)
N2	0.0184 (3)	0.8748 (3)	-0.0597 (2)	0.0470 (10)
N3	0.3357 (2)	0.6418 (3)	0.2771 (2)	0.0385 (8)
N4	0.1255 (2)	0.6241 (3)	0.2792 (2)	0.0406 (9)
N5	0.1706 (2)	0.5600 (3)	0.1086 (2)	0.0405 (9)
N6	0.2288 (2)	0.8465 (3)	0.2764 (2)	0.0410 (9)
C1	0.0848 (3)	0.8230 (3)	0.0267 (2)	0.0384 (10)
C2	-0.0930 (4)	0.8686 (4)	-0.0804 (3)	0.0548 (12)
C3	-0.1387 (3)	0.8134 (4)	-0.0174 (3)	0.0568 (14)
C4	-0.0639 (3)	0.7636 (4)	0.0716 (3)	0.0488 (12)
C5	0.3252 (3)	0.5685 (3)	0.3615 (2)	0.0452 (11)
C6	0.2038 (3)	0.5283 (3)	0.3347 (2)	0.0438 (11)
C7	0.3755 (3)	0.5708 (4)	0.2084 (3)	0.0508 (12)
C8	0.2801 (4)	0.4922 (4)	0.1429 (3)	0.0516 (12)
C9	0.4102 (3)	0.7457 (4)	0.3219 (3)	0.0484 (11)
C10	0.3430 (3)	0.8304 (3)	0.3597 (3)	0.0493 (12)
P1	0.35869 (9)	0.19313 (10)	0.36387 (9)	0.0445 (3)
F1	0.2435 (3)	0.2566 (4)	0.2930 (3)	0.1068 (16)
F2	0.4274 (5)	0.2804 (3)	0.3247 (6)	0.142 (3)
F3	0.3517 (3)	0.1030 (3)	0.2757 (2)	0.0873 (12)
F4	0.4707 (3)	0.1283 (5)	0.4302 (3)	0.1148 (16)
F5	0.2846 (5)	0.1060 (4)	0.3963 (5)	0.147 (3)
F6	0.3610 (4)	0.2851 (4)	0.4487 (3)	0.1020 (16)
Cl1	0.03840 (9)	0.91504 (11)	0.36135 (8)	0.0531 (3)
H1	-0.14280	0.90310	-0.14030	0.0660*
H2	-0.21700	0.81010	-0.03460	0.0680*
H3	-0.09110	0.72690	0.11630	0.0590*
H4	0.34970	0.61460	0.42370	0.0540*
H5	0.37450	0.50000	0.37360	0.0540*
H6	0.18900	0.45820	0.29230	0.0530*
H7	0.19180	0.50880	0.39630	0.0530*
H8	0.11730	0.67600	0.32400	0.0490*
H9	0.05650	0.59400	0.24190	0.0490*
H10	0.44060	0.52320	0.24880	0.0610*
H11	0.39870	0.62300	0.16570	0.0610*
H12	0.29470	0.46570	0.08420	0.0610*
H13	0.27480	0.42330	0.18150	0.0610*
H14	0.11560	0.51450	0.11480	0.0490*
H15	0.14950	0.57950	0.04250	0.0490*
H16	0.43190	0.78360	0.27070	0.0580*
H17	0.47890	0.72090	0.37780	0.0580*
H18	0.33480	0.79860	0.42000	0.0590*
H19	0.38220	0.90560	0.37730	0.0590*
H20	0.23020	0.90950	0.23830	0.0490*
H21	0.17640	0.85960	0.30340	0.0490*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.0335 (2)	0.0386 (2)	0.0257 (2)	-0.0010 (2)	0.0173 (2)	0.0001 (2)
S1	0.0416 (4)	0.0527 (5)	0.0362 (4)	-0.0043 (3)	0.0232 (3)	0.0054 (3)
N1	0.0375 (14)	0.0471 (17)	0.0312 (14)	0.0010 (12)	0.0177 (11)	0.0025 (12)
N2	0.057 (2)	0.0517 (19)	0.0315 (14)	0.0037 (15)	0.0162 (14)	0.0037 (13)
N3	0.0372 (14)	0.0458 (16)	0.0349 (14)	0.0015 (12)	0.0164 (12)	0.0003 (13)
N4	0.0485 (17)	0.0501 (17)	0.0316 (14)	-0.0057 (13)	0.0246 (13)	-0.0012 (12)
N5	0.0490 (17)	0.0472 (17)	0.0318 (14)	-0.0058 (13)	0.0226 (13)	-0.0046 (12)
N6	0.0473 (17)	0.0433 (16)	0.0369 (15)	-0.0022 (13)	0.0211 (13)	-0.0040 (13)
C1	0.0427 (18)	0.0446 (19)	0.0325 (16)	-0.0017 (15)	0.0194 (14)	-0.0001 (14)
C2	0.057 (2)	0.058 (2)	0.041 (2)	0.012 (2)	0.0096 (18)	-0.0013 (19)
C3	0.037 (2)	0.070 (3)	0.058 (2)	0.0057 (19)	0.0121 (18)	-0.006 (2)
C4	0.044 (2)	0.063 (2)	0.046 (2)	0.0001 (18)	0.0245 (17)	-0.0018 (19)
C5	0.054 (2)	0.046 (2)	0.0344 (17)	0.0047 (17)	0.0156 (16)	0.0055 (15)
C6	0.059 (2)	0.044 (2)	0.0335 (16)	-0.0009 (17)	0.0235 (16)	0.0045 (14)
C7	0.047 (2)	0.066 (2)	0.046 (2)	0.0145 (19)	0.0253 (17)	-0.0033 (19)
C8	0.069 (2)	0.049 (2)	0.045 (2)	0.0116 (19)	0.031 (2)	-0.0071 (17)
C9	0.0367 (18)	0.058 (2)	0.045 (2)	-0.0066 (17)	0.0094 (15)	0.0031 (19)
C10	0.057 (2)	0.043 (2)	0.041 (2)	-0.0059 (17)	0.0112 (17)	-0.0054 (16)
P1	0.0437 (5)	0.0468 (5)	0.0452 (5)	0.0057 (4)	0.0196 (4)	0.0026 (4)
F1	0.084 (2)	0.113 (3)	0.102 (3)	0.046 (2)	0.012 (2)	0.005 (2)
F2	0.170 (5)	0.072 (2)	0.264 (7)	-0.021 (2)	0.171 (5)	-0.006 (3)
F3	0.103 (2)	0.085 (2)	0.073 (2)	0.016 (2)	0.0329 (19)	-0.0226 (18)
F4	0.089 (2)	0.148 (4)	0.077 (2)	0.048 (2)	-0.002 (2)	0.002 (2)
F5	0.182 (5)	0.078 (2)	0.265 (7)	0.003 (3)	0.180 (5)	0.022 (3)
F6	0.120 (3)	0.112 (3)	0.073 (2)	0.015 (2)	0.036 (2)	-0.027 (2)
Cl1	0.0510 (5)	0.0707 (7)	0.0425 (5)	0.0165 (4)	0.0235 (4)	0.0068 (4)

Geometric parameters (\AA , $^\circ$)

Co1—S1	2.3187 (12)	N4—H9	0.9000
Co1—N1	1.930 (3)	N5—H15	0.9000
Co1—N3	1.944 (3)	N5—H14	0.9000
Co1—N4	1.954 (3)	N6—H21	0.9000
Co1—N5	1.966 (3)	N6—H20	0.9000
Co1—N6	1.970 (3)	C2—C3	1.389 (6)
Co1—C1	2.566 (3)	C3—C4	1.378 (6)
S1—C1	1.734 (4)	C5—C6	1.511 (5)
P1—F3	1.592 (3)	C7—C8	1.504 (6)
P1—F4	1.557 (5)	C9—C10	1.513 (6)
P1—F5	1.551 (6)	C2—H1	0.9300
P1—F6	1.585 (4)	C3—H2	0.9300
P1—F1	1.595 (4)	C4—H3	0.9300
P1—F2	1.556 (6)	C5—H5	0.9700
N1—C1	1.354 (4)	C5—H4	0.9700
N1—C4	1.332 (5)	C6—H6	0.9700

supplementary materials

N2—C2	1.333 (7)	C6—H7	0.9700
N2—C1	1.330 (4)	C7—H10	0.9700
N3—C5	1.506 (4)	C7—H11	0.9700
N3—C7	1.495 (5)	C8—H13	0.9700
N3—C9	1.491 (5)	C8—H12	0.9700
N4—C6	1.479 (5)	C9—H16	0.9700
N5—C8	1.498 (6)	C9—H17	0.9700
N6—C10	1.491 (5)	C10—H19	0.9700
N4—H8	0.9000	C10—H18	0.9700
S1—Co1—N1	72.31 (9)	H14—N5—H15	108.00
S1—Co1—N3	101.96 (9)	Co1—N5—H14	109.00
S1—Co1—N4	170.71 (9)	H20—N6—H21	108.00
S1—Co1—N5	89.62 (9)	Co1—N6—H20	110.00
S1—Co1—N6	87.67 (9)	C10—N6—H21	110.00
S1—Co1—C1	41.18 (9)	Co1—N6—H21	110.00
N1—Co1—N3	173.92 (12)	C10—N6—H20	110.00
N1—Co1—N4	98.40 (12)	Co1—C1—N1	47.51 (15)
N1—Co1—N5	91.63 (13)	Co1—C1—S1	61.73 (10)
N1—Co1—N6	95.02 (13)	Co1—C1—N2	171.7 (3)
N1—Co1—C1	31.15 (12)	N1—C1—N2	125.2 (4)
N3—Co1—N4	87.33 (12)	S1—C1—N1	109.2 (2)
N3—Co1—N5	86.23 (13)	S1—C1—N2	125.6 (3)
N3—Co1—N6	86.65 (13)	N2—C2—C3	123.6 (4)
N3—Co1—C1	143.02 (12)	C2—C3—C4	117.6 (4)
N4—Co1—N5	90.85 (13)	N1—C4—C3	119.5 (4)
N4—Co1—N6	93.09 (13)	N3—C5—C6	111.1 (2)
N4—Co1—C1	129.54 (12)	N4—C6—C5	109.1 (3)
N5—Co1—N6	171.69 (12)	N3—C7—C8	109.1 (3)
N5—Co1—C1	89.96 (11)	N5—C8—C7	109.0 (3)
N6—Co1—C1	93.16 (11)	N3—C9—C10	107.4 (3)
Co1—S1—C1	77.09 (11)	N6—C10—C9	107.8 (3)
F5—P1—F6	91.6 (3)	N2—C2—H1	118.00
F1—P1—F2	89.6 (3)	C3—C2—H1	118.00
F1—P1—F3	91.8 (2)	C2—C3—H2	121.00
F1—P1—F4	178.1 (2)	C4—C3—H2	121.00
F1—P1—F5	87.7 (3)	N1—C4—H3	120.00
F1—P1—F6	85.9 (2)	C3—C4—H3	120.00
F2—P1—F3	89.8 (3)	N3—C5—H5	109.00
F2—P1—F4	90.4 (3)	N3—C5—H4	109.00
F2—P1—F5	176.5 (4)	H4—C5—H5	108.00
F2—P1—F6	90.3 (3)	C6—C5—H4	109.00
F3—P1—F4	86.3 (2)	C6—C5—H5	109.00
F3—P1—F5	88.2 (3)	N4—C6—H6	110.00
F3—P1—F6	177.7 (2)	C5—C6—H7	110.00
F4—P1—F5	92.2 (3)	N4—C6—H7	110.00
F4—P1—F6	96.0 (2)	C5—C6—H6	110.00
Co1—N1—C1	101.4 (2)	H6—C6—H7	108.00
Co1—N1—C4	139.2 (3)	H10—C7—H11	108.00
C1—N1—C4	119.0 (3)	N3—C7—H10	110.00

C1—N2—C2	115.1 (3)	N3—C7—H11	110.00
Co1—N3—C5	110.3 (2)	C8—C7—H10	110.00
Co1—N3—C7	105.5 (2)	C8—C7—H11	110.00
Co1—N3—C9	106.6 (2)	N5—C8—H12	110.00
C5—N3—C7	112.3 (3)	N5—C8—H13	110.00
C5—N3—C9	109.3 (3)	C7—C8—H12	110.00
C7—N3—C9	112.6 (3)	C7—C8—H13	110.00
Co1—N4—C6	109.2 (2)	H12—C8—H13	108.00
Co1—N5—C8	110.8 (2)	N3—C9—H16	110.00
Co1—N6—C10	109.8 (2)	N3—C9—H17	110.00
C6—N4—H9	110.00	C10—C9—H16	110.00
H8—N4—H9	108.00	C10—C9—H17	110.00
Co1—N4—H8	110.00	H16—C9—H17	108.00
C6—N4—H8	110.00	N6—C10—H18	110.00
Co1—N4—H9	110.00	N6—C10—H19	110.00
C8—N5—H14	110.00	C9—C10—H18	110.00
C8—N5—H15	109.00	C9—C10—H19	110.00
Co1—N5—H15	109.00	H18—C10—H19	108.00
N1—Co1—S1—C1	-1.46 (15)	N4—Co1—N6—C10	85.4 (2)
N3—Co1—S1—C1	176.44 (15)	C1—Co1—N6—C10	-144.7 (2)
N5—Co1—S1—C1	90.38 (14)	S1—Co1—C1—N1	-177.3 (3)
N6—Co1—S1—C1	-97.48 (14)	N1—Co1—C1—S1	177.3 (3)
S1—Co1—N1—C1	1.86 (19)	N3—Co1—C1—S1	-5.8 (2)
S1—Co1—N1—C4	173.9 (4)	N3—Co1—C1—N1	176.9 (2)
N4—Co1—N1—C1	-178.3 (2)	N4—Co1—C1—S1	179.46 (13)
N4—Co1—N1—C4	-6.3 (4)	N4—Co1—C1—N1	2.2 (3)
N5—Co1—N1—C1	-87.2 (2)	N5—Co1—C1—S1	-89.47 (12)
N5—Co1—N1—C4	84.8 (4)	N5—Co1—C1—N1	93.2 (2)
N6—Co1—N1—C1	87.8 (2)	N6—Co1—C1—S1	82.83 (12)
N6—Co1—N1—C4	-100.1 (4)	N6—Co1—C1—N1	-94.5 (2)
C1—Co1—N1—C4	172.1 (6)	Co1—S1—C1—N1	2.1 (2)
S1—Co1—N3—C5	-179.45 (19)	Co1—S1—C1—N2	-175.4 (3)
S1—Co1—N3—C7	-58.0 (2)	Co1—N1—C1—S1	-2.5 (3)
S1—Co1—N3—C9	62.0 (2)	Co1—N1—C1—N2	175.0 (3)
N4—Co1—N3—C5	0.4 (2)	C4—N1—C1—Co1	-174.1 (4)
N4—Co1—N3—C7	121.9 (3)	C4—N1—C1—S1	-176.6 (3)
N4—Co1—N3—C9	-118.2 (2)	C4—N1—C1—N2	0.9 (6)
N5—Co1—N3—C5	-90.6 (2)	Co1—N1—C4—C3	-170.8 (3)
N5—Co1—N3—C7	30.8 (2)	C1—N1—C4—C3	0.3 (6)
N5—Co1—N3—C9	150.8 (2)	C2—N2—C1—S1	175.8 (3)
N6—Co1—N3—C5	93.6 (2)	C2—N2—C1—N1	-1.2 (5)
N6—Co1—N3—C7	-144.9 (2)	C1—N2—C2—C3	0.4 (6)
N6—Co1—N3—C9	-24.9 (2)	Co1—N3—C5—C6	20.3 (3)
C1—Co1—N3—C5	-175.55 (18)	C7—N3—C5—C6	-97.1 (3)
C1—Co1—N3—C7	-54.1 (3)	C9—N3—C5—C6	137.2 (3)
C1—Co1—N3—C9	65.9 (3)	Co1—N3—C7—C8	-47.1 (4)
N1—Co1—N4—C6	156.7 (2)	C5—N3—C7—C8	73.1 (4)
N3—Co1—N4—C6	-21.3 (2)	C9—N3—C7—C8	-163.0 (3)
N5—Co1—N4—C6	64.9 (2)	Co1—N3—C9—C10	46.4 (3)

supplementary materials

N6—Co1—N4—C6	-107.8 (2)	C5—N3—C9—C10	-72.8 (4)
C1—Co1—N4—C6	155.58 (18)	C7—N3—C9—C10	161.7 (3)
S1—Co1—N5—C8	93.0 (2)	Co1—N4—C6—C5	37.0 (3)
N1—Co1—N5—C8	165.3 (3)	Co1—N5—C8—C7	-15.2 (4)
N3—Co1—N5—C8	-9.0 (3)	Co1—N6—C10—C9	27.8 (3)
N4—Co1—N5—C8	-96.3 (3)	N2—C2—C3—C4	0.6 (7)
C1—Co1—N5—C8	134.2 (3)	C2—C3—C4—N1	-1.0 (6)
S1—Co1—N6—C10	-103.9 (2)	N3—C5—C6—N4	-37.6 (3)
N1—Co1—N6—C10	-175.9 (2)	N3—C7—C8—N5	40.9 (4)
N3—Co1—N6—C10	-1.8 (2)	N3—C9—C10—N6	-48.7 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N4—H8...N2 ⁱ	0.90	2.50	3.079 (4)	123
N4—H9...Cl1 ⁱⁱ	0.90	2.52	3.278 (3)	142
N5—H14...Cl1 ⁱⁱ	0.90	2.39	3.287 (3)	173
N5—H15...Cl1 ⁱⁱⁱ	0.90	2.42	3.273 (3)	160
N6—H21...Cl1	0.90	2.29	3.187 (3)	173
C6—H6...F1	0.97	2.39	3.210 (6)	143
C7—H11...F4 ^{iv}	0.97	2.52	3.328 (6)	141
C9—H17...F6 ^v	0.97	2.52	3.454 (6)	161

Symmetry codes: (i) $x, -y+3/2, z+1/2$; (ii) $-x, y-1/2, -z+1/2$; (iii) $x, -y+3/2, z-1/2$; (iv) $-x+1, y+1/2, -z+1/2$; (v) $-x+1, -y+1, -z+1$.

Fig. 1




Fig. 2

