

Poly[[diaqua-hemi- μ_4 -oxalato- μ_2 -oxalato-praseodymium(III)] monohydrate]

Ting-Hai Yang,* Qiang Chen, Wei Zhuang, Zhe Wang and Bang-Yi Yue

High-Tech Institute of Nanjing University, Changzhou 213164, People's Republic of China

Correspondence e-mail: tinghaiyang@gmail.com


Received 9 August 2009; accepted 25 August 2009

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.029; wR factor = 0.075; data-to-parameter ratio = 12.9.

In the title complex, $\{[\text{Pr}(\text{C}_2\text{O}_4)_{1.5}(\text{H}_2\text{O})_2]\cdot\text{H}_2\text{O}\}_n$, the Pr^{III} ion, which lies on a crystallographic inversion centre, is coordinated by seven O atoms from four oxalate ligands and two O atoms from two water ligands; further Pr–O coordination from tetradentate oxalate ligands forms a three-dimensional structure. The compound crystallized as a monohydrate, the water molecule occupying space in small voids and being secured by O–H...O hydrogen bonding as an acceptor from ligand water H atoms and as a donor to oxalate O-acceptor sites.

Related literature

For background to lanthanide oxalates and their preparation, see: Hansson (1970, 1972, 1973a, 1973b); Michaelides *et al.* (1988); Ollendorf & Weigel (1969); Steinfink & Brunton (1970); Trollet *et al.* (1998); Trombe (2003); Unaleroglu *et al.* (1997). For related structures, see: Trombe *et al.* (2004); Barrett Adams *et al.* (1998); Beagley *et al.* (1988).



Experimental

Crystal data

$[\text{Pr}(\text{C}_2\text{O}_4)_{1.5}(\text{H}_2\text{O})_2]\cdot\text{H}_2\text{O}$
 $M_r = 326.99$
 Triclinic, $P\bar{1}$
 $a = 6.0367$ (12) Å
 $b = 7.6222$ (15) Å
 $c = 8.9353$ (18) Å
 $\alpha = 98.330$ (4)°
 $\beta = 99.814$ (3)°

$\gamma = 96.734$ (4)°
 $V = 396.58$ (14) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 6.17$ mm⁻¹
 $T = 273$ K
 $0.18 \times 0.16 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2001)
 $T_{\text{min}} = 0.341$, $T_{\text{max}} = 0.542$

2140 measured reflections
 1521 independent reflections
 1450 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.020$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.075$
 $S = 1.00$
 1521 reflections

118 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.80$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.50$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2W–H2WB...O1W	0.84	2.55	2.858 (7)	103
O1W–H1WA...O2 ⁱ	0.84	1.96	2.694 (6)	146
O1W–H1WA...O3 ⁱⁱ	0.84	2.60	3.235 (6)	133
O1W–H1WB...O3W ⁱⁱⁱ	0.84	2.00	2.833 (6)	169
O2W–H2WA...O3W ^{iv}	0.84	1.98	2.807 (7)	166
O2W–H2WB...O3 ^v	0.84	2.20	2.919 (6)	144
O3W–H3WA...O6 ^{vi}	0.84	2.08	2.829 (7)	149
O3W–H3WB...O4 ^{vii}	0.84	2.03	2.833 (6)	159

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $x-1, y, z$; (iii) $-x+1, -y+1, -z$; (iv) $x+1, y+1, z$; (v) $-x+2, -y+2, -z$; (vi) $-x+1, -y+1, -z+1$; (vii) $-x+2, -y+1, -z$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg, 1999); software used to prepare material for publication: SHELXTL (Sheldrick, 2008).

The authors acknowledge the High-Tech Research Institute of Nanjing University for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NK2001).

References

- Barrett Adams, D. M. Y., Kahwa, I. A. & Magee, J. T. (1998). *New J. Chem.* **22**, 919–921.
 Beagley, B., Pritchard, R. G., Evmiridis, N. P., Michailides, A. & Skoulika, S. (1988). *Acta Cryst.* **C44**, 174–175.
 Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
 Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *SMART* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hansson, E. (1970). *Acta Chem. Scand.* **24**, 2969–2982.
 Hansson, E. (1972). *Acta Chem. Scand.* **26**, 1337–1350.
 Hansson, E. (1973a). *Acta Chem. Scand.* **27**, 823–834.

- Hansson, E. (1973*b*). *Acta Chem. Scand.* **27**, 2852–2860.
- Michaelides, A., Skoulika, S. & Aubry, A. (1988). *Mater. Res. Bull.* **23**, 579–585.
- Ollendorf, W. & Weigel, F. (1969). *Inorg. Nucl. Chem. Lett.* **5**, 263–269.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Steinfink, H. & Brunton, G. D. (1970). *Inorg. Chem.* **9**, 2112–2117.
- Trollet, D., Rome, S. & Mosset, A. (1998). *Polyhedron*, **17**, 3977–3978.
- Trombe, J. C. (2003). *J. Chem. Crystallogr.* **33**, 19–26.
- Trombe, J. C. & Mohanu, A. (2004). *Solid State Sci.* **6**, 1403–1419.
- Unaleroglu, C., Zumreoglu-Karan, B. & Zencir, Y. (1997). *Polyhedron*, **16**, 2155–2161.

supplementary materials

Acta Cryst. (2009). E65, m1152-m1153 [doi:10.1107/S1600536809033947]

Poly[[diaquahemi- μ_4 -oxalato- μ_2 -oxalato-praseodymium(III)] monohydrate]

T.-H. Yang, Q. Chen, W. Zhuang, Z. Wang and B.-Y. Yue

Comment

In the last decade considerable attention has been afforded to the structures and properties of lanthanide oxalates due to their ability to act as precursors of lanthanide oxides. Some single crystals of lanthanide oxalates, such as $[\text{Ln}(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_4] \cdot (\text{H}_2\text{O})$ (Ln = Sc or Yb), $[\text{Ln}_2(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_6] \cdot 4\text{H}_2\text{O}$ (Ln = La, Ce, Pr or Nd) and $[\text{Nd}_2(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_6]$ have been obtained either in silica gel (Ollendorff & Weigel, 1969), by hydrothermal reaction (Michaelides *et al.*, 1988) or by other methods (Hansson, 1970, 1972, 1973a,b; Unaleroglu *et al.*, 1997; Trollet *et al.*, 1998; Trombe, 2003). Crystals of $[\text{Ln}(\text{C}_2\text{O}_4)(\text{HC}_2\text{O}_4)] \cdot 4\text{H}_2\text{O}$ (Ln = Er or Tm) were prepared by saturating a boiling solution of oxalic acid in 3 M H_2SO_4 with the lanthanide oxide and then slowly cooling to 273 K (Steinfink & Brunton, 1970). However, a few praseodymium oxalate complex has been reported. The present paper is concerned with a new crystal structure of praseodymium oxalate complex with a three-dimensional network structure.

The asymmetric unit of the title compound, $[\text{Pr}(\text{C}_2\text{O}_4)_{1.5}(\text{H}_2\text{O})_2] \cdot (\text{H}_2\text{O})$, (I), is shown in Fig. 1. The Pr atom lies on an inversion centre and is coordinated by seven O atoms [O1, O1^{iv}, O2ⁱⁱ, O3, O4ⁱⁱⁱ, O5 and O6ⁱ] from four oxalate ligands, and two O atoms from two aqua ligands, thereby forming a slightly distorted PrO_9 polyhedral coordination geometry. The Pr—O bond distances range from 2.451 (4) Å to 2.608 (3) Å, in agreement with those in compounds (Trombe *et al.* (2004); Barrett Adams *et al.* (1998)).

In the complex, the equivalent Pr atom are connected is coordinated by seven O atoms from four oxalate ligands and two O atoms from water ligands. Further Pr—O coordination from the tetradentate oxalate ligands forms a three-dimensional structure (Fig.2). The compound crystallized as a monohydrate; this water molecule occupies space in small voids and is secured by O—H \cdots O hydrogen bonding as an acceptor from ligand water H-atoms and as a donor to oxalate O acceptor sites.

Experimental

All solvents and chemicals were of analytical grade and were used without further purification. $\text{Pr}(\text{NO}_3)_3 \cdot 6\text{H}_2\text{O}$ (0.05 mmol, 0.023 g), $\text{Na}_2\text{C}_2\text{O}_4$ (0.075 mmol, 0.011 g), and deionized water (10 ml) were mixed together. The mixture was sealed in a Teflon-line autoclave and then heated at 443 K for 5 d under autogenous pressure and then cooled to room temperature. Green crystals were obtained.

Refinement

All non-hydrogen atoms were refined anisotropically. The water H atoms were located in a difference Fourier map and refined with a distance restraint of O—H = 0.83–0.85 Å, and with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{iso}}(\text{O})$.

Figures




Fig. 1. Molecular structure of (I), showing 50% probability displacement ellipsoids. [symmetry codes:(i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 3, -y + 2, -z + 1$; (iii) $-x + 2, -y + 1, -z$; (iv) $-x + 2, -y + 2, -z + 1$].




Fig. 2. The unit cell packing diagram of (I).

Poly[[diaquaheptamethylenediamine- μ_4 -oxalato- μ_2 -oxalato-praseodymium(III)] monohydrate]

Crystal data

$[\text{Pr}(\text{C}_2\text{O}_4)_{1.5}(\text{H}_2\text{O})_2] \cdot \text{H}_2\text{O}$

$M_r = 326.99$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.0367\ (12)\ \text{\AA}$

$b = 7.6222\ (15)\ \text{\AA}$

$c = 8.9353\ (18)\ \text{\AA}$

$\alpha = 98.330\ (4)^\circ$

$\beta = 99.814\ (3)^\circ$

$\gamma = 96.734\ (4)^\circ$

$V = 396.58\ (14)\ \text{\AA}^3$

$Z = 2$

$F_{000} = 310$

$D_x = 2.738\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 905 reflections

$\theta = 3.3\text{--}28.3^\circ$

$\mu = 6.17\ \text{mm}^{-1}$

$T = 273\ \text{K}$

Block, green

$0.18 \times 0.16 \times 0.10\ \text{mm}$

Data collection

Bruker SMART APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 273\ \text{K}$

ϕ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2001)

$T_{\text{min}} = 0.341$, $T_{\text{max}} = 0.542$

2140 measured reflections

1521 independent reflections

1450 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.020$

$\theta_{\text{max}} = 26.0^\circ$

$\theta_{\text{min}} = 2.4^\circ$

$h = -7 \rightarrow 7$

$k = -9 \rightarrow 8$

$l = -10 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.029$	H-atom parameters constrained
$wR(F^2) = 0.075$	$w = 1/[\sigma^2(F_o^2) + (0.049P)^2 + 1.09P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
1521 reflections	$(\Delta/\sigma)_{\max} = 0.001$
118 parameters	$\Delta\rho_{\max} = 0.80 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -1.49 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Pr1	1.00780 (5)	0.80762 (4)	0.29489 (3)	0.01182 (13)
C1	1.4452 (9)	1.0278 (7)	0.5703 (6)	0.0155 (11)
C2	1.0808 (10)	0.5659 (7)	-0.0322 (7)	0.0163 (12)
C3	0.8789 (9)	0.5155 (7)	0.5109 (6)	0.0157 (11)
O1	1.2322 (6)	0.9964 (5)	0.5514 (4)	0.0156 (8)
O2	1.5785 (7)	1.1009 (6)	0.6910 (5)	0.0220 (9)
O3	1.1296 (7)	0.7232 (5)	0.0431 (5)	0.0195 (9)
O4	1.1455 (7)	0.5077 (5)	-0.1515 (5)	0.0217 (9)
O5	0.8004 (7)	0.6413 (5)	0.4552 (5)	0.0195 (9)
O6	0.7818 (7)	0.4116 (6)	0.5849 (5)	0.0230 (9)
O1W	0.6751 (7)	0.8292 (7)	0.0928 (5)	0.0310 (11)
H1WA	0.5581	0.8534	0.1260	0.046*
H1WB	0.6299	0.7771	0.0011	0.046*
O2W	1.0612 (8)	1.0993 (6)	0.2071 (5)	0.0288 (10)
H2WA	1.1653	1.1872	0.2227	0.043*
H2WB	0.9752	1.1029	0.1239	0.043*
O3W	0.4213 (9)	0.3694 (7)	0.2101 (6)	0.0420 (13)
H3WA	0.4103	0.4339	0.2922	0.063*

supplementary materials

H3WB 0.5590 0.3823 0.2031 0.063*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pr1	0.01064 (19)	0.01179 (18)	0.01243 (18)	0.00024 (12)	0.00360 (12)	-0.00045 (12)
C1	0.012 (3)	0.016 (3)	0.019 (3)	0.001 (2)	0.005 (2)	0.001 (2)
C2	0.015 (3)	0.016 (3)	0.018 (3)	0.002 (2)	0.005 (2)	0.002 (2)
C3	0.016 (3)	0.016 (3)	0.014 (3)	0.002 (2)	0.004 (2)	0.000 (2)
O1	0.0075 (19)	0.019 (2)	0.020 (2)	0.0013 (15)	0.0047 (15)	-0.0003 (16)
O2	0.013 (2)	0.032 (2)	0.016 (2)	-0.0027 (17)	0.0036 (16)	-0.0062 (18)
O3	0.023 (2)	0.016 (2)	0.017 (2)	-0.0029 (16)	0.0072 (17)	-0.0033 (16)
O4	0.025 (2)	0.018 (2)	0.020 (2)	-0.0062 (17)	0.0118 (17)	-0.0028 (17)
O5	0.021 (2)	0.019 (2)	0.023 (2)	0.0084 (17)	0.0089 (17)	0.0085 (17)
O6	0.017 (2)	0.026 (2)	0.031 (2)	0.0070 (18)	0.0095 (18)	0.0138 (19)
O1W	0.015 (2)	0.053 (3)	0.023 (2)	0.010 (2)	0.0049 (18)	-0.005 (2)
O2W	0.029 (3)	0.021 (2)	0.035 (3)	-0.0021 (19)	0.001 (2)	0.013 (2)
O3W	0.029 (3)	0.050 (3)	0.038 (3)	-0.013 (2)	0.013 (2)	-0.017 (2)

Geometric parameters (\AA , $^\circ$)

Pr1—O5	2.448 (4)	C2—C2 ⁱⁱⁱ	1.555 (11)
Pr1—O2W	2.466 (4)	C3—O5	1.248 (7)
Pr1—O6 ⁱ	2.472 (4)	C3—O6	1.258 (7)
Pr1—O2 ⁱⁱ	2.490 (4)	C3—C3 ⁱ	1.548 (11)
Pr1—O1W	2.500 (4)	O1—Pr1 ^{iv}	2.609 (4)
Pr1—O3	2.504 (4)	O2—Pr1 ⁱⁱ	2.490 (4)
Pr1—O4 ⁱⁱⁱ	2.541 (4)	O4—Pr1 ⁱⁱⁱ	2.541 (4)
Pr1—O1	2.586 (4)	O6—Pr1 ⁱ	2.472 (4)
Pr1—O1 ^{iv}	2.609 (4)	O1W—H1WA	0.8413
C1—O2	1.243 (7)	O1W—H1WB	0.8417
C1—O1	1.258 (7)	O2W—H2WA	0.8412
C1—C1 ⁱⁱ	1.546 (11)	O2W—H2WB	0.8372
C2—O4	1.238 (7)	O3W—H3WA	0.8386
C2—O3	1.263 (7)	O3W—H3WB	0.8400
O5—Pr1—O2W	142.99 (15)	O6 ⁱ —Pr1—O1 ^{iv}	120.77 (14)
O5—Pr1—O6 ⁱ	65.89 (14)	O2 ⁱⁱ —Pr1—O1 ^{iv}	121.99 (13)
O2W—Pr1—O6 ⁱ	142.54 (15)	O1W—Pr1—O1 ^{iv}	77.27 (13)
O5—Pr1—O2 ⁱⁱ	131.79 (14)	O3—Pr1—O1 ^{iv}	146.78 (13)
O2W—Pr1—O2 ⁱⁱ	71.54 (15)	O4 ⁱⁱⁱ —Pr1—O1 ^{iv}	122.88 (13)
O6 ⁱ —Pr1—O2 ⁱⁱ	71.37 (14)	O1—Pr1—O1 ^{iv}	65.40 (14)
O5—Pr1—O1W	97.65 (15)	O2—C1—O1	126.7 (5)
O2W—Pr1—O1W	70.26 (16)	O2—C1—C1 ⁱⁱ	116.0 (6)
O6 ⁱ —Pr1—O1W	142.27 (16)	O1—C1—C1 ⁱⁱ	117.3 (6)
O2 ⁱⁱ —Pr1—O1W	130.26 (15)	O4—C2—O3	126.8 (5)

O5—Pr1—O3	133.93 (13)	O4—C2—C2 ⁱⁱⁱ	117.5 (6)
O2W—Pr1—O3	78.07 (15)	O3—C2—C2 ⁱⁱⁱ	115.7 (6)
O6 ⁱ —Pr1—O3	92.40 (14)	O5—C3—O6	126.5 (5)
O2 ⁱⁱ —Pr1—O3	67.24 (13)	O5—C3—C3 ⁱ	117.1 (6)
O1W—Pr1—O3	74.66 (14)	O6—C3—C3 ⁱ	116.4 (6)
O5—Pr1—O4 ⁱⁱⁱ	70.30 (13)	C1—O1—Pr1	118.7 (3)
O2W—Pr1—O4 ⁱⁱⁱ	132.32 (15)	C1—O1—Pr1 ^{iv}	123.3 (3)
O6 ⁱ —Pr1—O4 ⁱⁱⁱ	69.96 (15)	Pr1—O1—Pr1 ^{iv}	114.60 (14)
O2 ⁱⁱ —Pr1—O4 ⁱⁱⁱ	114.58 (14)	C1—O2—Pr1 ⁱⁱ	123.5 (4)
O1W—Pr1—O4 ⁱⁱⁱ	72.53 (15)	C2—O3—Pr1	121.8 (4)
O3—Pr1—O4 ⁱⁱⁱ	64.03 (13)	C2—O4—Pr1 ⁱⁱⁱ	120.5 (4)
O5—Pr1—O1	85.86 (13)	C3—O5—Pr1	120.4 (4)
O2W—Pr1—O1	81.79 (14)	C3—O6—Pr1 ⁱ	119.7 (4)
O6 ⁱ —Pr1—O1	77.20 (14)	Pr1—O1W—H1WA	115.2
O2 ⁱⁱ —Pr1—O1	63.42 (12)	Pr1—O1W—H1WB	132.7
O1W—Pr1—O1	137.77 (14)	H1WA—O1W—H1WB	106.2
O3—Pr1—O1	130.36 (13)	Pr1—O2W—H2WA	136.4
O4 ⁱⁱⁱ —Pr1—O1	145.06 (13)	Pr1—O2W—H2WB	113.6
O5—Pr1—O1 ^{iv}	67.14 (13)	H2WA—O2W—H2WB	107.3
O2W—Pr1—O1 ^{iv}	76.00 (14)	H3WA—O3W—H3WB	107.4
O2—C1—O1—Pr1	-172.7 (5)	C2 ⁱⁱⁱ —C2—O3—Pr1	6.3 (8)
C1 ⁱⁱ —C1—O1—Pr1	8.0 (8)	O5—Pr1—O3—C2	1.9 (5)
O2—C1—O1—Pr1 ^{iv}	29.2 (8)	O2W—Pr1—O3—C2	-156.5 (4)
C1 ⁱⁱ —C1—O1—Pr1 ^{iv}	-150.1 (5)	O6 ⁱ —Pr1—O3—C2	60.1 (4)
O5—Pr1—O1—C1	133.4 (4)	O2 ⁱⁱ —Pr1—O3—C2	128.7 (5)
O2W—Pr1—O1—C1	-81.6 (4)	O1W—Pr1—O3—C2	-83.9 (4)
O6 ⁱ —Pr1—O1—C1	67.2 (4)	O4 ⁱⁱⁱ —Pr1—O3—C2	-6.2 (4)
O2 ⁱⁱ —Pr1—O1—C1	-8.2 (4)	O1—Pr1—O3—C2	135.3 (4)
O1W—Pr1—O1—C1	-129.8 (4)	O1 ^{iv} —Pr1—O3—C2	-117.2 (4)
O3—Pr1—O1—C1	-15.0 (5)	O3—C2—O4—Pr1 ⁱⁱⁱ	-174.2 (5)
O4 ⁱⁱⁱ —Pr1—O1—C1	87.3 (4)	C2 ⁱⁱⁱ —C2—O4—Pr1 ⁱⁱⁱ	4.8 (9)
O1 ^{iv} —Pr1—O1—C1	-159.9 (5)	O6—C3—O5—Pr1	-174.3 (5)
O5—Pr1—O1—Pr1 ^{iv}	-66.72 (16)	C3 ⁱ —C3—O5—Pr1	6.5 (8)
O2W—Pr1—O1—Pr1 ^{iv}	78.30 (17)	O2W—Pr1—O5—C3	-154.9 (4)
O6 ⁱ —Pr1—O1—Pr1 ^{iv}	-132.92 (18)	O6 ⁱ —Pr1—O5—C3	-6.5 (4)
O2 ⁱⁱ —Pr1—O1—Pr1 ^{iv}	151.7 (2)	O2 ⁱⁱ —Pr1—O5—C3	-36.2 (5)
O1W—Pr1—O1—Pr1 ^{iv}	30.1 (3)	O1W—Pr1—O5—C3	138.0 (4)
O3—Pr1—O1—Pr1 ^{iv}	144.93 (15)	O3—Pr1—O5—C3	61.9 (5)
O4 ⁱⁱⁱ —Pr1—O1—Pr1 ^{iv}	-112.8 (2)	O4 ⁱⁱⁱ —Pr1—O5—C3	69.7 (4)
O1 ^{iv} —Pr1—O1—Pr1 ^{iv}	0.0	O1—Pr1—O5—C3	-84.4 (4)
O1—C1—O2—Pr1 ⁱⁱ	-171.6 (4)	O1 ^{iv} —Pr1—O5—C3	-149.4 (4)

supplementary materials

C1 ⁱⁱ —C1—O2—Pr1 ⁱⁱ	7.8 (9)	O5—C3—O6—Pr1 ⁱ	-173.9 (4)
O4—C2—O3—Pr1	-174.7 (5)	C3 ⁱ —C3—O6—Pr1 ⁱ	5.3 (8)

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+3, -y+2, -z+1$; (iii) $-x+2, -y+1, -z$; (iv) $-x+2, -y+2, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2W—H2WB \cdots O1W	0.84	2.55	2.858 (7)	103
O1W—H1WA \cdots O2 ^{iv}	0.84	1.96	2.694 (6)	146
O1W—H1WA \cdots O3 ^v	0.84	2.60	3.235 (6)	133
O1W—H1WB \cdots O3W ^{vi}	0.84	2.00	2.833 (6)	169
O2W—H2WA \cdots O3W ^{vii}	0.84	1.98	2.807 (7)	166
O2W—H2WB \cdots O3 ^{viii}	0.84	2.20	2.919 (6)	144
O3W—H3WA \cdots O6 ^{ix}	0.84	2.08	2.829 (7)	149
O3W—H3WB \cdots O4 ⁱⁱⁱ	0.84	2.03	2.833 (6)	159

Symmetry codes: (iv) $-x+2, -y+2, -z+1$; (v) $x-1, y, z$; (vi) $-x+1, -y+1, -z$; (vii) $x+1, y+1, z$; (viii) $-x+2, -y+2, -z$; (ix) $-x+1, -y+1, -z+1$; (iii) $-x+2, -y+1, -z$.

Fig. 1




Fig. 2

