

2,2'-Bipyridine-5,5'-dicarboxylic acid**Chongchen Wang**

School of Environmental and Energy Engineering, Beijing University of Civil Engineering and Architecture, 100044 Beijing, People's Republic of China
Correspondence e-mail: chongchenwang@126.com


Received 15 July 2009; accepted 29 July 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$;
 R factor = 0.071; wR factor = 0.214; data-to-parameter ratio = 10.9.

The title molecule, $\text{C}_{12}\text{H}_8\text{N}_2\text{O}_4$, lies on an inversion center. In the crystal structure, intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds connect molecules into one-dimensional chains along $[1\bar{1}1]$.

Related literature

For synthetic applications of the title compound, see: Schokecht & Kempe (2004).

**Experimental***Crystal data*

$\text{C}_{12}\text{H}_8\text{N}_2\text{O}_4$	$\gamma = 90.000(1)^\circ$
$M_r = 244.20$	$V = 254.34(6)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 3.7384(5)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 6.3934(8)\text{ \AA}$	$\mu = 0.12\text{ mm}^{-1}$
$c = 10.7786(13)\text{ \AA}$	$T = 298\text{ K}$
$\alpha = 98.774(2)^\circ$	$0.15 \times 0.11 \times 0.08\text{ mm}$
$\beta = 92.567(1)^\circ$	

Data collection

Bruker SMART CCD diffractometer	1343 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	893 independent reflections
$T_{\min} = 0.982$, $T_{\max} = 0.990$	657 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.071$	82 parameters
$wR(F^2) = 0.214$	H-atom parameters constrained
$S = 1.14$	$\Delta\rho_{\max} = 0.33\text{ e \AA}^{-3}$
893 reflections	$\Delta\rho_{\min} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1 \cdots O2 ⁱ	0.82	1.82	2.625 (3)	168

Symmetry code: (i) $-x + 1, -y, -z + 2$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge the financial support of the Funding Project for Academic Human Resources Development in Institutions of Higher Learning under the Jurisdiction of Beijing Municipality and the Research Fund of Beijing University of Civil Engineering and Architecture..

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2868).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Schokecht, B. & Kempe, R. (2004). *Z. Anorg. Allg. Chem.* **630**, 1377–1379.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supplementary materials

Acta Cryst. (2009). E65, o2081 [doi:10.1107/S1600536809030207]

2,2'-Bipyridine-5,5'-dicarboxylic acid

C. Wang

Comment

2,2'-bipyridine-5,5'-dicarboxylate acid is a potential multi-dentate ligand with a versatile coordination mode, which has been used in self-assembled porous coordination synthesis (Schökecht & Kempe, 2004). The crystals of the title compound were obtained unintentionally as the harvested product of the hydrothermal reaction of 2,2'-bipyridine-5,5'-dicarboxylate acid, Eu₂O₃ and 1,10-phenanthroline.

The molecular structure of the title compound is shown in Fig. 1. In the crystal structure, intermolecular O—H···O hydrogen bonds connect molecules into one-dimensional chains along [1 -1 1] (Fig. 2).

Experimental

Yellow needle-like crystals of the title compound were obtained by hydrothermal reaction of 2,2'-bipyridine-5,5'-dicarboxylate acid (0.04884 g), 1,10-phenanthroline (0.0360 g), Eu₂O₃ (0.0702 g) and deionized water (15 ml) in a 23 ml teflon-lined reaction vessel at 433 K for 120 h, followed by slow cooling to room temperature.

Refinement

All H atoms were placed in calculated positions and included in a riding-model approximation, with C—H = 0.93 Å, O—H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of the title compound shown with 30% probability ellipsoids [symmetry code: (a) -x, -y+1, -z+1].




Fig. 2. Part of the crystal structure of the title compound with hydrogen bonds shown as dashed lines. The one-dimensional hydrogen-bonded chains propagate along [1 -1 1].

supplementary materials

2,2'-Bipyridine-5,5'-dicarboxylic acid

Crystal data

C ₁₂ H ₈ N ₂ O ₄	Z = 1
M _r = 244.20	F ₀₀₀ = 126
Triclinic, P $\bar{1}$	D _x = 1.594 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 3.7384 (5) Å	Cell parameters from 528 reflections
b = 6.3934 (8) Å	θ = 3.2–27.6°
c = 10.7786 (13) Å	μ = 0.12 mm ⁻¹
α = 98.774 (2)°	T = 298 K
β = 92.567 (1)°	Needle, yellow
γ = 90.000 (1)°	0.15 × 0.11 × 0.08 mm
V = 254.34 (6) Å ³	

Data collection

Bruker SMART CCD diffractometer	893 independent reflections
Radiation source: fine-focus sealed tube	657 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
T = 298 K	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -4 \rightarrow 4$
$T_{\text{min}} = 0.982$, $T_{\text{max}} = 0.990$	$k = -7 \rightarrow 7$
1343 measured reflections	$l = -11 \rightarrow 12$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.071$	H-atom parameters constrained
$wR(F^2) = 0.214$	$w = 1/[\sigma^2(F_o^2) + (0.1116P)^2 + 0.1159P]$ where $P = (F_o^2 + 2F_c^2)/3$
S = 1.14	$(\Delta/\sigma)_{\text{max}} < 0.001$
893 reflections	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
82 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.1979 (7)	0.2556 (4)	0.5341 (2)	0.0351 (8)
O1	0.5129 (7)	-0.0316 (4)	0.8383 (2)	0.0492 (8)
H1	0.5614	-0.0788	0.9036	0.074*
O2	0.2843 (7)	0.2274 (4)	0.9736 (2)	0.0536 (9)
C1	0.3541 (8)	0.1439 (5)	0.8636 (3)	0.0334 (8)
C2	0.2905 (8)	0.1631 (5)	0.6337 (3)	0.0347 (9)
H2	0.3922	0.0293	0.6197	0.042*
C3	0.2438 (7)	0.2552 (5)	0.7571 (3)	0.0307 (9)
C4	0.0964 (8)	0.4566 (5)	0.7784 (3)	0.0353 (9)
H4	0.0640	0.5238	0.8598	0.042*
C5	-0.0008 (8)	0.5548 (5)	0.6768 (3)	0.0332 (8)
H5	-0.1009	0.6892	0.6890	0.040*
C6	0.0520 (8)	0.4512 (5)	0.5562 (3)	0.0295 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0443 (16)	0.0333 (15)	0.0284 (15)	0.0064 (12)	-0.0011 (12)	0.0079 (12)
O1	0.0724 (18)	0.0463 (16)	0.0308 (14)	0.0208 (13)	-0.0006 (12)	0.0129 (11)
O2	0.083 (2)	0.0525 (17)	0.0267 (14)	0.0231 (14)	0.0056 (12)	0.0108 (11)
C1	0.0347 (17)	0.0366 (18)	0.0297 (17)	0.0043 (14)	-0.0014 (13)	0.0088 (14)
C2	0.0390 (18)	0.0339 (18)	0.0329 (18)	0.0076 (14)	-0.0009 (13)	0.0108 (14)
C3	0.0309 (16)	0.0350 (19)	0.0271 (18)	0.0003 (14)	-0.0020 (13)	0.0083 (14)
C4	0.0446 (19)	0.0375 (19)	0.0240 (16)	0.0068 (15)	0.0024 (13)	0.0047 (13)
C5	0.0393 (18)	0.0321 (17)	0.0296 (17)	0.0078 (14)	0.0009 (13)	0.0092 (14)
C6	0.0288 (15)	0.0318 (18)	0.0289 (17)	-0.0009 (13)	-0.0021 (12)	0.0088 (14)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.335 (4)	C2—H2	0.9300
N1—C6	1.357 (4)	C3—C4	1.391 (4)
O1—C1	1.267 (4)	C4—C5	1.378 (4)
O1—H1	0.8200	C4—H4	0.9300

supplementary materials

O2—C1	1.263 (4)	C5—C6	1.388 (4)
C1—C3	1.484 (4)	C5—H5	0.9300
C2—C3	1.388 (4)	C6—C6 ⁱ	1.482 (6)
C2—N1—C6	117.4 (3)	C4—C3—C1	120.8 (3)
C1—O1—H1	109.5	C5—C4—C3	118.9 (3)
O2—C1—O1	123.7 (3)	C5—C4—H4	120.5
O2—C1—C3	118.7 (3)	C3—C4—H4	120.5
O1—C1—C3	117.6 (3)	C4—C5—C6	119.3 (3)
N1—C2—C3	123.8 (3)	C4—C5—H5	120.3
N1—C2—H2	118.1	C6—C5—H5	120.3
C3—C2—H2	118.1	N1—C6—C5	122.4 (3)
C2—C3—C4	118.2 (3)	N1—C6—C6 ⁱ	116.1 (3)
C2—C3—C1	121.0 (3)	C5—C6—C6 ⁱ	121.5 (4)
C6—N1—C2—C3	0.4 (5)	C2—C3—C4—C5	0.8 (5)
N1—C2—C3—C4	-0.9 (5)	C1—C3—C4—C5	179.7 (3)
N1—C2—C3—C1	-179.8 (3)	C3—C4—C5—C6	-0.3 (5)
O2—C1—C3—C2	-175.3 (3)	C2—N1—C6—C5	0.2 (5)
O1—C1—C3—C2	4.4 (5)	C2—N1—C6—C6 ⁱ	-179.3 (3)
O2—C1—C3—C4	5.9 (5)	C4—C5—C6—N1	-0.2 (5)
O1—C1—C3—C4	-174.4 (3)	C4—C5—C6—C6 ⁱ	179.2 (3)


Symmetry codes: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O1—H1···O2 ⁱⁱ	0.82	1.82	2.625 (3)	168

Symmetry codes: (ii) $-x+1, -y, -z+2$.

Fig. 1



supplementary materials

Fig. 2

