

N,N'-(Methylenedi-*p*-phenylene)-dibenzamide

Sohail Saeed,^{a*} Naghmana Rashid,^a Rizwan Hussain^b and Peter G. Jones^c

^aDepartment of Chemistry, Research Complex, Allama Iqbal Open University, Islamabad, Pakistan, ^bDirectorate of Chemical & Power Sources, National Development Complex, PO Box 2216, Islamabad, Pakistan, and ^cInstitut für Anorganische und Analytische Chemie, Technische Universität Braunschweig, Postfach 3329, 38023 Braunschweig, Germany
Correspondence e-mail: sohail262001@yahoo.com


Received 30 July 2009; accepted 7 August 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.038; wR factor = 0.102; data-to-parameter ratio = 15.1.

The title compound, $\text{C}_{27}\text{H}_{22}\text{N}_2\text{O}_2$, consists of two chemically equivalent halves. However, it displays no crystallographic symmetry, only an approximate local twofold symmetry (r.m.s. deviation = 0.15 Å between the two halves of the molecule) is observed. In the crystal, molecules are connected by two antiparallel classical $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds, forming broad chains parallel to $(10\bar{1})$. A series of weak $\text{C}-\text{H}\cdots\text{N}/\text{O}$ hydrogen bonds is also present.

Related literature

For general background to the chemistry of polymers and polyamides, see Ataei *et al.* (2005); Yang *et al.* (2002). For related structures, see: Im & Jung (2000).



Experimental

Crystal data

$\text{C}_{27}\text{H}_{22}\text{N}_2\text{O}_2$ $a = 5.7296$ (7) Å
 $M_r = 406.47$ $b = 9.601$ (1) Å
 Triclinic, $P\bar{1}$ $c = 20.045$ (2) Å

$\alpha = 88.517$ (8)°
 $\beta = 82.293$ (8)°
 $\gamma = 75.67$ (1)°
 $V = 1058.7$ (2) Å³
 $Z = 2$

Cu $K\alpha$ radiation
 $\mu = 0.64$ mm⁻¹
 $T = 100$ K
 $0.20 \times 0.10 \times 0.04$ mm

Data collection

Oxford Diffraction Xcalibur Nova A diffractometer $T_{\min} = 0.781$, $T_{\max} = 1.000$
 (expected range = 0.762–0.975)
 Absorption correction: multi-scan 14884 measured reflections
 (CrysAlis Pro; Oxford 4355 independent reflections
 Diffraction, 2009) 3819 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.102$ $\Delta\rho_{\text{max}} = 0.16$ e Å⁻³
 $S = 1.07$ $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³
 4355 reflections
 288 parameters

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H01}\cdots\text{O2}^i$	0.875 (17)	2.017 (17)	2.8745 (13)	166.3 (14)
$\text{N2}-\text{H02}\cdots\text{O1}^{ii}$	0.876 (16)	2.088 (16)	2.9358 (12)	162.7 (14)
$\text{C16}-\text{H16}\cdots\text{O2}^i$	0.95	2.60	3.3090 (14)	132
$\text{C35}-\text{H35}\cdots\text{N1}^i$	0.95	2.72	3.5459 (15)	146
$\text{C33}-\text{H33}\cdots\text{O1}^{ii}$	0.95	2.43	3.1913 (15)	137
$\text{C42}-\text{H42}\cdots\text{O1}^{ii}$	0.95	2.58	3.3003 (14)	133

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y, -z + 1$.

Data collection: *CrysAlis Pro* (Oxford Diffraction, 2009); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis Pro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* (Siemens, 1994); software used to prepare material for publication: *SHELXL97*.

The authors are grateful to Allama Iqbal Open University and National Development complex, Islamabad, Pakistan for the allocation of research and analytical laboratory facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IM2132).

References

- Ataei, S. M., Sarrafi, Y., Hatami, M. & Faizi, L. A. (2005). *Eur. Polym. J.* **41**, 491–499.
 Im, J. K. & Jung, J. C. (2000). *Polymers*, **41**, 8709–8716.
 Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Siemens (1994). *XP*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Yang, C.-P., Chen, R.-S. & Hsu, M.-F. (2002). *J. Polym. Res.* **9**, 245–250.

supplementary materials

Acta Cryst. (2009). E65, o2140 [doi:10.1107/S1600536809031304]

N,N'-(Methylenedi-*p*-phenylene)dibenzamide

S. Saeed, N. Rashid, R. Hussain and P. G. Jones

Comment

High-temperature polymers have received much attention because of increasing demands for the replacement of ceramics and metals (Ataei *et al.*, 2005). However, in many cases they are insoluble and do not melt below their decomposition temperature, which restricts their applications (Im & Jung, 2000). Many studies have therefore focused on obtaining aromatic polymers that are processable by conventional techniques (Yang *et al.*, 2002). The title compound is a logical precursor for an attempt to synthesize polyamides and polyimides having excellent thermal and mechanical properties.

The molecule of the title compound is shown in Fig. 1. Molecular dimensions may be regarded as normal, as may the *trans* geometry at the amide groups. The molecule possesses no crystallographic symmetry, but displays approximate twofold symmetry with a r.m.s. deviation of 0.15 Å between the two halves of the molecule. There are however significant differences between torsion angles of chemically equivalent amide groups, *e.g.* C1—N1—C24—C25 - 31.5 (2) *versus*. C3—N2—C34—C35 - 16.2 (2)°. The outer pairs of rings are approximately parallel [interplanar angles: C11—16/C21—26 2.91 (7), C31—36/C41—46 10.72 (6)°] whereas the central pair of rings are approximately perpendicular [C21—26/C31—36 84.02 (4)°].

The main features of the molecular packing are the classical H bonds of the N—H...OC type, which are mutually antiparallel and link the molecules to form broad chains parallel to (10 $\bar{1}$) (Fig. 2, Table 1). A series of narrow-angled (C—H...N/O 132–146°) weak H bonds are probably of less structural significance (Table 1).

Experimental

All reagents and organic solvents were of analytical grade and commercially available. The title compound was accidentally generated during the reaction of 4,4'-diaminodiphenylmethane with 2-thiophene-carbonyl chloride; it was isolated from the reaction mixture by column chromatography in 30% yield and then purified by recrystallization from ethanol. Colourless single crystals suitable for X-ray analysis were obtained after one week by slow evaporation from an ethanolic solution. Crystals formed as thin plates or somewhat thicker laths; both proved to have the same cell constants.

Refinement

NH H atoms were refined freely. Other H atoms were placed in calculated positions and refined using a riding model with C—H_{arom} 0.95 Å, C—H_{methylene} 0.99 Å; these hydrogen *U* values were fixed at 1.2 × *U*(eq) of the parent atom. Data are 99.6% complete to 2θ 145°.

Figures




Fig. 1. Thermal ellipsoid plot (50% probability level) of the title compound.




Fig. 2. Packing diagram viewed perpendicular to $(10\bar{1})$. Classical H bonds are indicated by thick dashed lines. H atoms not involved in these H bonds are omitted for clarity.

N,N'-(Methylenedi-*p*-phenylene)dibenzamide

Crystal data

$C_{27}H_{22}N_2O_2$

$M_r = 406.47$

Triclinic, $P\bar{1}$

$a = 5.7296$ (7) Å

$b = 9.601$ (1) Å

$c = 20.045$ (2) Å

$\alpha = 88.517$ (8)°

$\beta = 82.293$ (8)°

$\gamma = 75.67$ (1)°

$V = 1058.7$ (2) Å³

$Z = 2$

$F_{000} = 428$

$D_x = 1.275$ Mg m⁻³

Melting point: 449 K

Cu $K\alpha$ radiation, $\lambda = 1.54184$ Å

Cell parameters from 10160 reflections

$\theta = 3.3$ – 75.8 °

$\mu = 0.64$ mm⁻¹

$T = 100$ K

Lath, colourless

$0.20 \times 0.10 \times 0.04$ mm

Data collection

Oxford Diffraction Xcalibur Nova A diffractometer

Radiation source: Nova (Cu) X-ray Source

Monochromator: mirror

Detector resolution: 10.3543 pixels mm⁻¹

$T = 100$ K

ω scans

Absorption correction: multi-scan (CrysAlis Pro; Oxford Diffraction, 2009)

$T_{\min} = 0.781$, $T_{\max} = 1.000$

14884 measured reflections

4355 independent reflections

3819 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\text{max}} = 76.0$ °

$\theta_{\text{min}} = 4.5$ °

$h = -7 \rightarrow 7$

$k = -12 \rightarrow 10$

$l = -25 \rightarrow 24$

Refinement

Refinement on F^2

Least-squares matrix: full

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$$R[F^2 > 2\sigma(F^2)] = 0.038$$

$$wR(F^2) = 0.102$$

$$S = 1.07$$

4355 reflections

288 parameters

Primary atom site location: structure-invariant direct methods

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0552P)^2 + 0.2168P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Least-squares planes (x,y,z in crystal coordinates) and deviations from them (* indicates atom used to define plane)

$$-2.5047(0.0026)x + 7.1432(0.0034)y + 2.4159(0.0097)z = 0.1119(0.0034)$$

$$* -0.0003(0.0008)C11 * -0.0057(0.0008)C12 * 0.0066(0.0009)C13 * -0.0015(0.0009)C14 * -0.0045(0.0008)C15 * 0.0054(0.0008)C16 - 0.0844(0.0017)C1 - 0.5968(0.0020)O1 0.4205(0.0020)N1$$

Rms deviation of fitted atoms = 0.0046

$$-2.3282(0.0026)x + 7.4133(0.0033)y + 1.8134(0.0088)z = 0.5814(0.0051)$$

Angle to previous plane (with approximate e.s.d.) = 2.91 (0.07)

$$* -0.0079(0.0008)C21 * 0.0076(0.0008)C22 * 0.0009(0.0008)C23 * -0.0090(0.0008)C24 * 0.0085(0.0008)C25 * -0.0001(0.0008)C26 0.0290(0.0016)N1 - 0.0695(0.0019)C2$$

Rms deviation of fitted atoms = 0.0067

$$3.4319(0.0022)x + 3.8138(0.0049)y + 16.6363(0.0061)z = 13.6255(0.0031)$$

Angle to previous plane (with approximate e.s.d.) = 84.02 (0.04)

$$* 0.0014(0.0008)C31 * -0.0008(0.0009)C32 * -0.0014(0.0009)C33 * 0.0030(0.0008)C34 * -0.0024(0.0008)C35 * 0.0002(0.0008)C36 - 0.0109(0.0016)N2 0.0150(0.0018)C2$$

Rms deviation of fitted atoms = 0.0018

$$2.8287(0.0024)x + 4.9963(0.0040)y + 16.4884(0.0060)z = 14.1826(0.0051)$$

Angle to previous plane (with approximate e.s.d.) = 10.72 (0.06)

supplementary materials

* -0.0120 (0.0008) C41 * 0.0057 (0.0008) C42 * 0.0049 (0.0008) C43 * -0.0092 (0.0008) C44 * 0.0029 (0.0008) C45 * 0.0078 (0.0008) C46 - 0.0326 (0.0017) C3 0.3984 (0.0019) O2 - 0.5360 (0.0020) N2

Rms deviation of fitted atoms = 0.0077

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.76324 (16)	0.11312 (8)	0.25609 (4)	0.0293 (2)
O2	-0.05166 (15)	0.43352 (8)	0.76182 (4)	0.02817 (19)
N1	0.91641 (18)	0.30294 (10)	0.27475 (5)	0.0241 (2)
H01	0.958 (3)	0.3787 (17)	0.2567 (8)	0.034 (4)*
N2	0.15925 (18)	0.20072 (10)	0.73950 (5)	0.0237 (2)
H02	0.156 (3)	0.1111 (18)	0.7476 (8)	0.033 (4)*
C1	0.8376 (2)	0.21803 (11)	0.23510 (6)	0.0234 (2)
C2	0.9989 (2)	0.24643 (14)	0.55736 (6)	0.0294 (3)
H2A	1.1296	0.1591	0.5625	0.035*
H2B	1.0546	0.3298	0.5712	0.035*
C3	-0.0255 (2)	0.30332 (11)	0.77064 (5)	0.0231 (2)
C11	0.8434 (2)	0.25660 (11)	0.16191 (6)	0.0239 (2)
C12	0.6816 (2)	0.21104 (13)	0.12666 (6)	0.0291 (3)
H12	0.5695	0.1623	0.1497	0.035*
C13	0.6834 (2)	0.23651 (14)	0.05825 (6)	0.0340 (3)
H13	0.5705	0.2068	0.0347	0.041*
C14	0.8497 (2)	0.30526 (13)	0.02406 (6)	0.0330 (3)
H14	0.8522	0.3216	-0.0229	0.040*
C15	1.0122 (2)	0.35010 (12)	0.05876 (6)	0.0297 (3)
H15	1.1265	0.3968	0.0354	0.036*
C16	1.0087 (2)	0.32696 (12)	0.12763 (6)	0.0255 (2)
H16	1.1189	0.3591	0.1513	0.031*
C21	0.9656 (2)	0.26234 (12)	0.48346 (6)	0.0245 (2)
C22	1.1219 (2)	0.32417 (13)	0.44001 (6)	0.0275 (2)
H22	1.2421	0.3597	0.4577	0.033*
C23	1.1061 (2)	0.33505 (12)	0.37156 (6)	0.0270 (2)
H23	1.2157	0.3768	0.3428	0.032*
C24	0.9296 (2)	0.28478 (11)	0.34494 (5)	0.0228 (2)
C25	0.7677 (2)	0.22582 (12)	0.38770 (6)	0.0249 (2)
H25	0.6438	0.1936	0.3703	0.030*
C26	0.7884 (2)	0.21443 (12)	0.45618 (6)	0.0260 (2)
H26	0.6787	0.1729	0.4850	0.031*
C31	0.7762 (2)	0.23612 (13)	0.60485 (5)	0.0257 (2)
C32	0.7603 (2)	0.10857 (14)	0.63724 (7)	0.0333 (3)
H32	0.8919	0.0259	0.6290	0.040*
C33	0.5569 (2)	0.09899 (13)	0.68137 (6)	0.0314 (3)
H33	0.5504	0.0104	0.7028	0.038*
C34	0.3625 (2)	0.21848 (12)	0.69433 (5)	0.0227 (2)
C35	0.3748 (2)	0.34678 (12)	0.66207 (6)	0.0279 (2)
H35	0.2428	0.4293	0.6701	0.033*

C36	0.5802 (2)	0.35433 (12)	0.61812 (6)	0.0282 (3)
H36	0.5867	0.4428	0.5966	0.034*
C41	-0.2063 (2)	0.24956 (11)	0.81921 (5)	0.0230 (2)
C42	-0.1489 (2)	0.11654 (12)	0.85073 (6)	0.0258 (2)
H42	0.0112	0.0567	0.8422	0.031*
C43	-0.3239 (2)	0.07097 (12)	0.89452 (6)	0.0285 (3)
H43	-0.2834	-0.0199	0.9157	0.034*
C44	-0.5577 (2)	0.15796 (13)	0.90741 (6)	0.0289 (3)
H44	-0.6783	0.1261	0.9367	0.035*
C45	-0.6148 (2)	0.29195 (13)	0.87733 (6)	0.0292 (3)
H45	-0.7741	0.3524	0.8867	0.035*
C46	-0.4400 (2)	0.33797 (12)	0.83371 (6)	0.0266 (2)
H46	-0.4799	0.4301	0.8136	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0409 (5)	0.0230 (4)	0.0258 (4)	-0.0147 (3)	0.0015 (3)	0.0001 (3)
O2	0.0323 (4)	0.0205 (4)	0.0313 (4)	-0.0085 (3)	0.0007 (3)	0.0014 (3)
N1	0.0306 (5)	0.0204 (4)	0.0223 (5)	-0.0097 (4)	-0.0008 (4)	0.0027 (3)
N2	0.0295 (5)	0.0189 (4)	0.0236 (5)	-0.0094 (4)	-0.0005 (4)	0.0019 (3)
C1	0.0245 (5)	0.0195 (5)	0.0247 (5)	-0.0049 (4)	0.0012 (4)	-0.0004 (4)
C2	0.0259 (6)	0.0382 (6)	0.0256 (6)	-0.0111 (5)	-0.0035 (4)	0.0054 (5)
C3	0.0277 (6)	0.0213 (5)	0.0219 (5)	-0.0082 (4)	-0.0046 (4)	0.0006 (4)
C11	0.0269 (5)	0.0187 (5)	0.0242 (5)	-0.0038 (4)	-0.0001 (4)	-0.0004 (4)
C12	0.0298 (6)	0.0295 (6)	0.0285 (6)	-0.0095 (5)	-0.0011 (5)	-0.0003 (4)
C13	0.0358 (7)	0.0386 (7)	0.0290 (6)	-0.0099 (5)	-0.0076 (5)	-0.0012 (5)
C14	0.0412 (7)	0.0324 (6)	0.0230 (6)	-0.0058 (5)	-0.0026 (5)	0.0014 (5)
C15	0.0350 (6)	0.0254 (5)	0.0267 (6)	-0.0071 (5)	0.0024 (5)	0.0027 (4)
C16	0.0278 (6)	0.0219 (5)	0.0261 (6)	-0.0058 (4)	-0.0016 (4)	0.0003 (4)
C21	0.0235 (5)	0.0238 (5)	0.0245 (5)	-0.0038 (4)	-0.0010 (4)	0.0020 (4)
C22	0.0258 (6)	0.0308 (6)	0.0283 (6)	-0.0114 (5)	-0.0043 (4)	0.0042 (4)
C23	0.0275 (6)	0.0281 (6)	0.0266 (6)	-0.0114 (5)	-0.0001 (4)	0.0052 (4)
C24	0.0258 (5)	0.0177 (5)	0.0231 (5)	-0.0032 (4)	-0.0011 (4)	0.0013 (4)
C25	0.0242 (5)	0.0258 (5)	0.0251 (5)	-0.0081 (4)	-0.0005 (4)	-0.0007 (4)
C26	0.0251 (5)	0.0274 (5)	0.0251 (5)	-0.0091 (4)	0.0028 (4)	0.0017 (4)
C31	0.0283 (6)	0.0318 (6)	0.0196 (5)	-0.0119 (5)	-0.0039 (4)	0.0028 (4)
C32	0.0309 (6)	0.0291 (6)	0.0348 (6)	-0.0025 (5)	0.0030 (5)	0.0057 (5)
C33	0.0356 (7)	0.0234 (5)	0.0332 (6)	-0.0075 (5)	0.0016 (5)	0.0071 (4)
C34	0.0267 (5)	0.0235 (5)	0.0197 (5)	-0.0100 (4)	-0.0019 (4)	-0.0006 (4)
C35	0.0335 (6)	0.0230 (5)	0.0259 (6)	-0.0074 (4)	0.0010 (5)	0.0006 (4)
C36	0.0376 (6)	0.0238 (5)	0.0246 (5)	-0.0125 (5)	-0.0009 (5)	0.0027 (4)
C41	0.0273 (6)	0.0222 (5)	0.0211 (5)	-0.0090 (4)	-0.0032 (4)	-0.0015 (4)
C42	0.0286 (6)	0.0231 (5)	0.0248 (5)	-0.0065 (4)	-0.0005 (4)	-0.0005 (4)
C43	0.0354 (6)	0.0236 (5)	0.0264 (6)	-0.0095 (5)	0.0000 (5)	0.0022 (4)
C44	0.0305 (6)	0.0332 (6)	0.0251 (6)	-0.0142 (5)	0.0018 (5)	-0.0017 (5)
C45	0.0254 (6)	0.0319 (6)	0.0293 (6)	-0.0060 (5)	-0.0016 (5)	-0.0020 (5)
C46	0.0299 (6)	0.0244 (5)	0.0253 (5)	-0.0065 (4)	-0.0037 (4)	0.0009 (4)

supplementary materials

Geometric parameters (Å, °)

O1—C1	1.2310 (14)	C41—C46	1.3941 (16)
O2—C3	1.2325 (14)	C41—C42	1.3945 (16)
N1—C1	1.3462 (15)	C42—C43	1.3888 (16)
N1—C24	1.4230 (14)	C43—C44	1.3868 (17)
N2—C3	1.3485 (15)	C44—C45	1.3891 (17)
N2—C34	1.4188 (14)	C45—C46	1.3880 (17)
C1—C11	1.5018 (15)	N1—H01	0.875 (17)
C2—C31	1.5086 (16)	N2—H02	0.876 (16)
C2—C21	1.5180 (15)	C2—H2A	0.9900
C3—C41	1.5021 (15)	C2—H2B	0.9900
C11—C12	1.3937 (17)	C12—H12	0.9500
C11—C16	1.3943 (16)	C13—H13	0.9500
C12—C13	1.3858 (17)	C14—H14	0.9500
C13—C14	1.3878 (19)	C15—H15	0.9500
C14—C15	1.3872 (18)	C16—H16	0.9500
C15—C16	1.3908 (16)	C22—H22	0.9500
C21—C26	1.3907 (16)	C23—H23	0.9500
C21—C22	1.3933 (16)	C25—H25	0.9500
C22—C23	1.3866 (16)	C26—H26	0.9500
C23—C24	1.3932 (16)	C32—H32	0.9500
C24—C25	1.3918 (15)	C33—H33	0.9500
C25—C26	1.3928 (16)	C35—H35	0.9500
C31—C36	1.3874 (17)	C36—H36	0.9500
C31—C32	1.3884 (17)	C42—H42	0.9500
C32—C33	1.3861 (17)	C43—H43	0.9500
C33—C34	1.3889 (17)	C44—H44	0.9500
C34—C35	1.3892 (16)	C45—H45	0.9500
C35—C36	1.3882 (17)	C46—H46	0.9500
C1—N1—C24	125.97 (9)	C1—N1—H01	118.0 (10)
C3—N2—C34	128.27 (9)	C24—N1—H01	116.0 (10)
O1—C1—N1	123.28 (11)	C3—N2—H02	117.2 (10)
O1—C1—C11	119.98 (10)	C34—N2—H02	114.5 (10)
N1—C1—C11	116.74 (9)	C31—C2—H2A	108.4
C31—C2—C21	115.57 (10)	C21—C2—H2A	108.4
O2—C3—N2	124.60 (10)	C31—C2—H2B	108.4
O2—C3—C41	119.96 (10)	C21—C2—H2B	108.4
N2—C3—C41	115.44 (9)	H2A—C2—H2B	107.4
C12—C11—C16	119.42 (11)	C13—C12—H12	119.8
C12—C11—C1	116.81 (10)	C11—C12—H12	119.8
C16—C11—C1	123.66 (10)	C12—C13—H13	119.9
C13—C12—C11	120.35 (11)	C14—C13—H13	119.9
C12—C13—C14	120.16 (12)	C15—C14—H14	120.1
C15—C14—C13	119.78 (11)	C13—C14—H14	120.1
C14—C15—C16	120.33 (11)	C14—C15—H15	119.8
C15—C16—C11	119.94 (11)	C16—C15—H15	119.8
C26—C21—C22	117.76 (10)	C15—C16—H16	120.0

C26—C21—C2	122.95 (10)	C11—C16—H16	120.0
C22—C21—C2	119.26 (10)	C23—C22—H22	119.3
C23—C22—C21	121.45 (11)	C21—C22—H22	119.3
C22—C23—C24	120.02 (10)	C22—C23—H23	120.0
C25—C24—C23	119.47 (10)	C24—C23—H23	120.0
C25—C24—N1	122.82 (10)	C24—C25—H25	120.2
C23—C24—N1	117.64 (10)	C26—C25—H25	120.2
C24—C25—C26	119.59 (10)	C21—C26—H26	119.2
C21—C26—C25	121.68 (10)	C25—C26—H26	119.2
C36—C31—C32	117.52 (11)	C33—C32—H32	119.2
C36—C31—C2	121.11 (11)	C31—C32—H32	119.2
C32—C31—C2	121.37 (11)	C32—C33—H33	119.9
C33—C32—C31	121.54 (11)	C34—C33—H33	119.9
C32—C33—C34	120.24 (11)	C36—C35—H35	120.0
C33—C34—C35	118.99 (11)	C34—C35—H35	120.0
C33—C34—N2	117.27 (10)	C31—C36—H36	119.1
C35—C34—N2	123.72 (10)	C35—C36—H36	119.1
C36—C35—C34	119.93 (11)	C43—C42—H42	119.8
C31—C36—C35	121.77 (11)	C41—C42—H42	119.8
C46—C41—C42	119.09 (10)	C44—C43—H43	119.9
C46—C41—C3	118.15 (10)	C42—C43—H43	119.9
C42—C41—C3	122.75 (10)	C43—C44—H44	120.1
C43—C42—C41	120.47 (11)	C45—C44—H44	120.1
C44—C43—C42	120.11 (11)	C46—C45—H45	119.9
C43—C44—C45	119.71 (11)	C44—C45—H45	119.9
C46—C45—C44	120.30 (11)	C45—C46—H46	119.9
C45—C46—C41	120.28 (11)	C41—C46—H46	119.9
C24—N1—C1—O1	0.93 (18)	C2—C21—C26—C25	-177.65 (10)
C24—N1—C1—C11	-179.06 (10)	C24—C25—C26—C21	0.86 (17)
C34—N2—C3—O2	3.43 (18)	C21—C2—C31—C36	-70.56 (14)
C34—N2—C3—C41	-176.26 (10)	C21—C2—C31—C32	109.90 (13)
O1—C1—C11—C12	24.81 (15)	C36—C31—C32—C33	-0.13 (19)
N1—C1—C11—C12	-155.20 (11)	C2—C31—C32—C33	179.42 (12)
O1—C1—C11—C16	-151.47 (11)	C31—C32—C33—C34	-0.1 (2)
N1—C1—C11—C16	28.53 (15)	C32—C33—C34—C35	0.48 (19)
C16—C11—C12—C13	-0.57 (17)	C32—C33—C34—N2	179.41 (11)
C1—C11—C12—C13	-177.01 (10)	C3—N2—C34—C33	164.96 (11)
C11—C12—C13—C14	1.23 (19)	C3—N2—C34—C35	-16.17 (18)
C12—C13—C14—C15	-0.82 (19)	C33—C34—C35—C36	-0.58 (17)
C13—C14—C15—C16	-0.24 (18)	N2—C34—C35—C36	-179.43 (10)
C14—C15—C16—C11	0.89 (17)	C32—C31—C36—C35	0.03 (18)
C12—C11—C16—C15	-0.49 (16)	C2—C31—C36—C35	-179.52 (11)
C1—C11—C16—C15	175.69 (10)	C34—C35—C36—C31	0.32 (18)
C31—C2—C21—C26	-25.19 (16)	O2—C3—C41—C46	23.81 (15)
C31—C2—C21—C22	156.51 (11)	N2—C3—C41—C46	-156.49 (10)
C26—C21—C22—C23	-1.41 (17)	O2—C3—C41—C42	-155.33 (11)
C2—C21—C22—C23	176.98 (11)	N2—C3—C41—C42	24.37 (15)
C21—C22—C23—C24	0.60 (18)	C46—C41—C42—C43	1.81 (16)
C22—C23—C24—C25	0.97 (17)	C3—C41—C42—C43	-179.06 (10)

supplementary materials

C22—C23—C24—N1	177.98 (10)	C41—C42—C43—C44	-0.21 (17)
C1—N1—C24—C25	-31.50 (17)	C42—C43—C44—C45	-1.20 (18)
C1—N1—C24—C23	151.60 (11)	C43—C44—C45—C46	1.01 (18)
C23—C24—C25—C26	-1.68 (16)	C44—C45—C46—C41	0.60 (17)
N1—C24—C25—C26	-178.53 (10)	C42—C41—C46—C45	-2.00 (16)
C22—C21—C26—C25	0.67 (17)	C3—C41—C46—C45	178.83 (10)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H01 \cdots O2 ⁱ	0.875 (17)	2.017 (17)	2.8745 (13)	166.3 (14)
N2—H02 \cdots O1 ⁱⁱ	0.876 (16)	2.088 (16)	2.9358 (12)	162.7 (14)
C16—H16 \cdots O2 ⁱ	0.95	2.60	3.3090 (14)	132
C35—H35 \cdots N1 ⁱ	0.95	2.72	3.5459 (15)	146
C33—H33 \cdots O1 ⁱⁱ	0.95	2.43	3.1913 (15)	137
C42—H42 \cdots O1 ⁱⁱ	0.95	2.58	3.3003 (14)	133

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y, -z+1$.

Fig. 1




Fig. 2

