

7,9-Diallyl-6-methyl-7*H*-1,2,4-triazolo-[4,3-*b*][1,2,4]triazepin-8(9*H*)-one

Redwan Mohamed Zemama,^a Ibtissam Amari,^a Rachid Bouhfid,^b El Mokhtar Essassi^a and Seik Weng Ng^{c*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bInstitute of Nanomaterials and Nanotechnology, Avenue de l'Armée Royale, Madinat El Irfane, 10100 Rabat, Morocco, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 7 August 2009; accepted 8 August 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.053; wR factor = 0.176; data-to-parameter ratio = 14.8.

The title compound, $C_{12}H_{15}N_5O$, features a triazolyl ring fused with a seven-membered triazepinyl ring; the latter ring adopts a boat conformation with the allyl-bearing C atom as the prow and the C and N fused-ring atoms as the stern.

Related literature

Triazepines are used in the treatment of neuronal disorders. They are also the reactants for the synthesis of other heterocyclic compounds; see, for example: Essassi *et al.* (1977); Richter & Sheefelot (1991).



Experimental

Crystal data

$C_{12}H_{15}N_5O$	$V = 1254.19(8)\text{ \AA}^3$
$M_r = 245.29$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 7.4674(3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 8.3398(3)\text{ \AA}$	$T = 293\text{ K}$
$c = 20.2214(6)\text{ \AA}$	$0.3 \times 0.3 \times 0.3\text{ mm}$
$\beta = 95.174(2)^\circ$	

Data collection

Bruker APEX2 diffractometer	2435 independent reflections
Absorption correction: none	1600 reflections with $I > 2\sigma(I)$
11394 measured reflections	$R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$	164 parameters
$wR(F^2) = 0.176$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.62\text{ e \AA}^{-3}$
2435 reflections	$\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2583).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2005). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Essassi, E. M., Lavergne, J. P. & Viallefont, P. (1977). *Tetrahedron*, **33**, 2807–2812.
Richter, P. & Sheefelot, U. (1991). *Pharmazie*, **46**, 701–705.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o2152 [doi:10.1107/S160053680903133X]

7,9-Diallyl-6-methyl-7*H*-1,2,4-triazolo[4,3-*b*][1,2,4]triazepin-8(*9H*)-one

R. M. Zemama, I. Amari, R. Bouhfid, E. M. Essassi and S. W. Ng

Experimental

To a solution of 6-methyl-7*H*-[1,2,4]triazolo[4,3-*b*][1,2,4]triazepin-8(*9H*)-one (1 g, 6 mmol) in *N,N*-dimethylformamide (20 ml), potassium carbonate (1.26 g, 9 mmol), allyl bromide (0.8 ml, 9 mmol) and a catalytic amount of tetrabutylammonium bromide were added. The mixture was stirred for 12 h. After the completion of the reaction (as monitored by TLC), the solid material was removed by filtration and the solvent evaporated under vacuum. Dichloromethane (20 ml) was added and the solution filtered. The solvent was removed and the product purified by column chromatography (30% ethyl acetate/hexane) to afford colorless crystals in 30% yield; m.p. 423 K. The formulation was established by proton and carbon-13 NMR spectroscopy in DMSO-d₆.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation, with *U*(H) set to 1.2*U*(C).

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of C₁₂H₁₅N₅O at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

7,9-Diallyl-6-methyl-7*H*-1,2,4-triazolo[4,3-*b*][1,2,4]triazepin-8(*9H*)-one

Crystal data

C ₁₂ H ₁₅ N ₅ O	<i>F</i> ₀₀₀ = 520
<i>M_r</i> = 245.29	<i>D_x</i> = 1.299 Mg m ⁻³
Monoclinic, <i>P</i> 2 ₁ / <i>n</i>	Mo <i>K</i> α radiation, λ = 0.71073 Å
Hall symbol: -P 2yn	Cell parameters from 2666 reflections
<i>a</i> = 7.4674 (3) Å	θ = 2.6–23.2°
<i>b</i> = 8.3398 (3) Å	μ = 0.09 mm ⁻¹
<i>c</i> = 20.2214 (6) Å	<i>T</i> = 293 K
β = 95.174 (2)°	Block, colorless
<i>V</i> = 1254.19 (8) Å ³	0.3 × 0.3 × 0.3 mm
<i>Z</i> = 4	

supplementary materials

Data collection

Bruker APEXII diffractometer	1600 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.041$
Monochromator: graphite	$\theta_{\text{max}} = 25.9^\circ$
$T = 293 \text{ K}$	$\theta_{\text{min}} = 2.0^\circ$
φ and ω scans	$h = -9 \rightarrow 9$
Absorption correction: None	$k = -10 \rightarrow 10$
11394 measured reflections	$l = -22 \rightarrow 24$
2435 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.053$	H-atom parameters constrained
$wR(F^2) = 0.176$	$w = 1/[\sigma^2(F_o^2) + (0.1011P)^2 + 0.1814P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2435 reflections	$\Delta\rho_{\text{max}} = 0.62 \text{ e \AA}^{-3}$
164 parameters	$\Delta\rho_{\text{min}} = -0.31 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6016 (3)	0.9072 (2)	0.62240 (9)	0.0504 (5)
N1	0.6575 (3)	0.4291 (2)	0.67475 (9)	0.0372 (5)
N2	0.6714 (3)	0.4126 (2)	0.60624 (9)	0.0346 (5)
N3	0.7253 (3)	0.2937 (3)	0.51391 (10)	0.0486 (6)
N4	0.7390 (3)	0.4593 (3)	0.50437 (9)	0.0424 (6)
N5	0.7149 (3)	0.6893 (2)	0.57403 (9)	0.0360 (5)
C1	0.5941 (3)	0.7634 (3)	0.61190 (10)	0.0353 (6)
C2	0.4588 (3)	0.6533 (3)	0.63974 (10)	0.0328 (6)
H2	0.4112	0.5834	0.6033	0.039*
C3	0.3002 (3)	0.7447 (3)	0.66310 (12)	0.0460 (7)
H3A	0.3402	0.8057	0.7025	0.055*
H3B	0.2560	0.8200	0.6289	0.055*
C4	0.1488 (4)	0.6348 (4)	0.67861 (16)	0.0600 (8)
H4	0.1127	0.5587	0.6465	0.072*
C5	0.0662 (5)	0.6348 (5)	0.73019 (19)	0.0877 (12)
H5A	0.0969	0.7084	0.7639	0.105*
H5B	-0.0256	0.5613	0.7346	0.105*
C6	0.5616 (3)	0.5471 (3)	0.69075 (10)	0.0338 (6)

C7	0.5614 (4)	0.5818 (3)	0.76328 (11)	0.0480 (7)
H7A	0.6349	0.5044	0.7882	0.072*
H7B	0.4406	0.5759	0.7758	0.072*
H7C	0.6086	0.6874	0.7724	0.072*
C8	0.6870 (3)	0.2708 (3)	0.57428 (13)	0.0433 (6)
H8	0.6722	0.1707	0.5933	0.052*
C9	0.7056 (3)	0.5264 (3)	0.56022 (11)	0.0334 (6)
C10	0.8431 (3)	0.7867 (3)	0.53946 (12)	0.0449 (7)
H10A	0.9421	0.7191	0.5286	0.054*
H10B	0.8920	0.8704	0.5692	0.054*
C11	0.7587 (4)	0.8618 (3)	0.47735 (13)	0.0519 (7)
H11	0.6702	0.9388	0.4813	0.062*
C12	0.8000 (5)	0.8273 (4)	0.41876 (16)	0.0779 (11)
H12A	0.8880	0.7509	0.4130	0.093*
H12B	0.7420	0.8788	0.3820	0.093*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0657 (13)	0.0282 (11)	0.0585 (11)	-0.0048 (9)	0.0122 (9)	-0.0007 (8)
N1	0.0448 (12)	0.0352 (12)	0.0313 (10)	-0.0009 (10)	0.0013 (8)	0.0048 (8)
N2	0.0407 (12)	0.0290 (11)	0.0341 (10)	0.0007 (9)	0.0037 (8)	0.0025 (8)
N3	0.0564 (15)	0.0419 (14)	0.0482 (13)	0.0010 (11)	0.0084 (11)	-0.0081 (10)
N4	0.0485 (13)	0.0417 (13)	0.0376 (11)	-0.0001 (10)	0.0073 (9)	-0.0018 (9)
N5	0.0391 (12)	0.0314 (12)	0.0381 (10)	-0.0050 (9)	0.0063 (9)	0.0049 (8)
C1	0.0424 (15)	0.0303 (14)	0.0327 (12)	0.0003 (11)	-0.0005 (10)	0.0010 (10)
C2	0.0363 (13)	0.0296 (13)	0.0324 (12)	-0.0009 (10)	0.0016 (10)	0.0014 (9)
C3	0.0504 (17)	0.0422 (15)	0.0461 (14)	0.0071 (13)	0.0087 (12)	0.0053 (11)
C4	0.0377 (16)	0.081 (2)	0.0619 (18)	0.0146 (15)	0.0103 (13)	0.0091 (16)
C5	0.066 (2)	0.106 (3)	0.093 (3)	0.009 (2)	0.018 (2)	0.018 (2)
C6	0.0360 (14)	0.0328 (13)	0.0323 (12)	-0.0049 (11)	0.0018 (10)	0.0028 (10)
C7	0.0541 (17)	0.0556 (18)	0.0340 (13)	0.0011 (14)	0.0022 (11)	0.0019 (11)
C8	0.0489 (16)	0.0299 (14)	0.0510 (15)	0.0011 (12)	0.0038 (12)	-0.0002 (11)
C9	0.0323 (13)	0.0331 (14)	0.0342 (12)	-0.0008 (10)	0.0004 (9)	0.0016 (10)
C10	0.0443 (16)	0.0436 (16)	0.0475 (14)	-0.0111 (13)	0.0080 (12)	0.0063 (11)
C11	0.0567 (18)	0.0493 (18)	0.0519 (16)	-0.0001 (14)	0.0166 (13)	0.0142 (13)
C12	0.100 (3)	0.079 (3)	0.0560 (19)	0.009 (2)	0.0132 (18)	0.0164 (17)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.218 (3)	C3—H3B	0.9700
N1—C6	1.276 (3)	C4—C5	1.259 (4)
N1—N2	1.405 (3)	C4—H4	0.9300
N2—C8	1.358 (3)	C5—H5A	0.9300
N2—C9	1.369 (3)	C5—H5B	0.9300
N3—C8	1.293 (3)	C6—C7	1.495 (3)
N3—N4	1.400 (3)	C7—H7A	0.9600
N4—C9	1.304 (3)	C7—H7B	0.9600
N5—C1	1.381 (3)	C7—H7C	0.9600

supplementary materials

N5—C9	1.388 (3)	C8—H8	0.9300
N5—C10	1.479 (3)	C10—C11	1.491 (4)
C1—C2	1.511 (3)	C10—H10A	0.9700
C2—C6	1.514 (3)	C10—H10B	0.9700
C2—C3	1.519 (3)	C11—C12	1.283 (4)
C2—H2	0.9800	C11—H11	0.9300
C3—C4	1.510 (4)	C12—H12A	0.9300
C3—H3A	0.9700	C12—H12B	0.9300
C6—N1—N2	114.74 (18)	H5A—C5—H5B	120.0
C8—N2—C9	104.5 (2)	N1—C6—C7	116.6 (2)
C8—N2—N1	124.87 (19)	N1—C6—C2	122.7 (2)
C9—N2—N1	129.60 (19)	C7—C6—C2	120.7 (2)
C8—N3—N4	107.47 (19)	C6—C7—H7A	109.5
C9—N4—N3	106.37 (19)	C6—C7—H7B	109.5
C1—N5—C9	121.80 (19)	H7A—C7—H7B	109.5
C1—N5—C10	119.9 (2)	C6—C7—H7C	109.5
C9—N5—C10	117.75 (19)	H7A—C7—H7C	109.5
O1—C1—N5	121.0 (2)	H7B—C7—H7C	109.5
O1—C1—C2	123.7 (2)	N3—C8—N2	110.9 (2)
N5—C1—C2	115.2 (2)	N3—C8—H8	124.5
C1—C2—C6	107.12 (18)	N2—C8—H8	124.5
C1—C2—C3	112.1 (2)	N4—C9—N2	110.7 (2)
C6—C2—C3	116.29 (18)	N4—C9—N5	125.7 (2)
C1—C2—H2	106.9	N2—C9—N5	123.41 (19)
C6—C2—H2	106.9	N5—C10—C11	112.7 (2)
C3—C2—H2	106.9	N5—C10—H10A	109.0
C4—C3—C2	112.3 (2)	C11—C10—H10A	109.0
C4—C3—H3A	109.1	N5—C10—H10B	109.0
C2—C3—H3A	109.1	C11—C10—H10B	109.0
C4—C3—H3B	109.1	H10A—C10—H10B	107.8
C2—C3—H3B	109.1	C12—C11—C10	124.4 (3)
H3A—C3—H3B	107.9	C12—C11—H11	117.8
C5—C4—C3	127.1 (4)	C10—C11—H11	117.8
C5—C4—H4	116.4	C11—C12—H12A	120.0
C3—C4—H4	116.4	C11—C12—H12B	120.0
C4—C5—H5A	120.0	H12A—C12—H12B	120.0
C4—C5—H5B	120.0		
C6—N1—N2—C8	−147.4 (2)	C1—C2—C6—C7	101.0 (2)
C6—N1—N2—C9	46.1 (3)	C3—C2—C6—C7	−25.3 (3)
C8—N3—N4—C9	−0.7 (3)	N4—N3—C8—N2	0.8 (3)
C9—N5—C1—O1	−178.4 (2)	C9—N2—C8—N3	−0.6 (3)
C10—N5—C1—O1	−6.7 (3)	N1—N2—C8—N3	−170.0 (2)
C9—N5—C1—C2	3.0 (3)	N3—N4—C9—N2	0.3 (3)
C10—N5—C1—C2	174.68 (19)	N3—N4—C9—N5	176.1 (2)
O1—C1—C2—C6	−110.7 (2)	C8—N2—C9—N4	0.2 (3)
N5—C1—C2—C6	67.9 (2)	N1—N2—C9—N4	168.8 (2)
O1—C1—C2—C3	18.0 (3)	C8—N2—C9—N5	−175.7 (2)
N5—C1—C2—C3	−163.40 (19)	N1—N2—C9—N5	−7.1 (4)

supplementary materials

C1—C2—C3—C4	169.1 (2)	C1—N5—C9—N4	142.8 (2)
C6—C2—C3—C4	−67.2 (3)	C10—N5—C9—N4	−29.0 (3)
C2—C3—C4—C5	130.9 (3)	C1—N5—C9—N2	−41.9 (3)
N2—N1—C6—C7	−172.6 (2)	C10—N5—C9—N2	146.2 (2)
N2—N1—C6—C2	4.6 (3)	C1—N5—C10—C11	−78.4 (3)
C1—C2—C6—N1	−76.1 (3)	C9—N5—C10—C11	93.6 (3)
C3—C2—C6—N1	157.6 (2)	N5—C10—C11—C12	−114.4 (3)

supplementary materials

Fig. 1

