

## 2-[2-(2-Pyridyl)ethyl]isoindolinium perchlorate

Ray J. Butcher,\* Yohannes T. Tesema, Teshome B. Yisgedu and Yilma Gultneh

Department of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA

Correspondence e-mail: rbutcher99@yahoo.com


Received 12 August 2009; accepted 12 August 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ; disorder in solvent or counterion;  $R$  factor = 0.044;  $wR$  factor = 0.120; data-to-parameter ratio = 6.1.

In the title salt,  $\text{C}_{15}\text{H}_{17}\text{N}_2^+\cdot\text{ClO}_4^-$ , the isoindoline N atom is protonated and an intramolecular  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bond occurs. In the crystal,  $\text{N}-\text{H}\cdots\text{O}$  and numerous weak  $\text{C}-\text{H}\cdots\text{O}$  interactions occur between the cation and anion. The O atoms of the perchlorate anion are disordered over four sets of sites with occupancies of 0.438 (4), 0.270 (9), 0.155 (8) and 0.138 (5).

### Related literature

For further information on the synthesis, see: Bonnett *et al.* (1983); Meyers & Santiago (1995).



### Experimental

#### Crystal data

$\text{C}_{15}\text{H}_{17}\text{N}_2^+\cdot\text{ClO}_4^-$	$V = 1537.0$ (4) $\text{\AA}^3$
$M_r = 324.76$	$Z = 4$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation
$a = 7.7941$ (9) $\text{\AA}$	$\mu = 0.27\text{ mm}^{-1}$
$b = 11.7032$ (19) $\text{\AA}$	$T = 293\text{ K}$
$c = 16.850$ (3) $\text{\AA}$	$0.5 \times 0.2 \times 0.08\text{ mm}$

### Data collection

Bruker P4 diffractometer	2001 independent reflections
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	1660 reflections with $I > 2\sigma(I)$
$T_{\min} = 0.241$ , $T_{\max} = 0.265$	$R_{\text{int}} = 0.025$
(expected range = 0.890–0.979)	3 standard reflections
2027 measured reflections	every 97 reflections

intensity decay: <2%

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$	245 restraints
$wR(F^2) = 0.120$	H-atom parameters constrained
$S = 1.03$	$\Delta\rho_{\max} = 0.21\text{ e \AA}^{-3}$
2001 reflections	$\Delta\rho_{\min} = -0.15\text{ e \AA}^{-3}$
329 parameters	

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N2—H2B···N1	0.91	2.11	2.769 (4)	129
N2—H2B···O4	0.91	2.39	3.197 (9)	148
N2—H2B···O2A	0.91	2.53	3.222 (19)	133
N2—H2B···O1C	0.91	2.59	3.390 (16)	147
C2—H2A···O1B <sup>i</sup>	0.93	2.30	3.062 (11)	139
C3—H3A···O1A <sup>ii</sup>	0.93	2.62	3.193 (10)	121
C7—H7B···O2A <sup>iii</sup>	0.97	2.50	3.232 (16)	132
C8—H8A···O3	0.97	2.55	3.383 (10)	144
C8—H8A···O2B	0.97	2.39	3.168 (19)	136
C8—H8A···O3C	0.97	2.29	3.189 (17)	154
C8—H8B···O4B <sup>iv</sup>	0.97	2.48	3.167 (17)	128
C11—H11A···O3A <sup>v</sup>	0.93	2.55	3.27 (2)	135
C11—H11A···O3B <sup>v</sup>	0.93	2.57	3.47 (2)	162
C13—H13A···O3 <sup>vi</sup>	0.93	2.48	3.299 (9)	148
C13—H13A···O4A <sup>vi</sup>	0.93	2.20	2.976 (8)	140
C13—H13A···O1B <sup>vi</sup>	0.93	2.56	3.44 (2)	158
C13—H13A···O3C <sup>vi</sup>	0.93	2.70	3.444 (17)	138
C15—H15A···O1 <sup>iii</sup>	0.97	2.55	3.423 (9)	150
C15—H15B···O2 <sup>vi</sup>	0.97	2.48	3.425 (10)	163
C15—H15B···O4A <sup>vi</sup>	0.97	2.55	3.194 (19)	124

Symmetry codes: (i)  $-x - \frac{1}{2}, -y + 2, z - \frac{1}{2}$ ; (ii)  $x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$ ; (iii)  $-x, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iv)  $-x - 1, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (v)  $-x - \frac{1}{2}, -y + 2, z + \frac{1}{2}$ ; (vi)  $x + 1, y, z$ .

Data collection: *XSCANS* (Bruker, 1997); cell refinement: *XSCANS*; data reduction: *SHELXTL* (Sheldrick, 2008); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

RJB wishes to acknowledge the Laboratory for the Structure of Matter at the Naval Research Laboratory for access to their diffractometers.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5043).

### References

- Bonnett, R., North, S. A., Newton, R. F. & Scopes, D. I. C. (1983). *Tetrahedron*, **39**, 1401–1405.
- Bruker (1997). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Meyers, A. I. & Santiago, B. (1995). *Tetrahedron Lett.* **36**, 5877–5880.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o2175 [doi:10.1107/S1600536809032024]

## 2-[2-(2-Pyridyl)ethyl]isoindolinium perchlorate

R. J. Butcher, Y. T. Tesema, T. B. Yisgedu and Y. Gultneh

### Comment

The structure of the title compound, (I), is shown below. Dimensions are available in the archived CIF.

The crystal structure shows that the N atom of 2,3-dihydro-1*H*-isoindoline is protonated and involved in hydrogen bonding to the nearest pyridyl N atom (Table 1). The perchlorate ion O atoms are also involved in the hydrogen bonding with the isoindoline H atom. The amine H atom is also involved in hydrogen bonding to the perchlorate O atoms and there are extensive but weak interionic C—H···O interactions between the cation and anion (Table 1).

### Experimental

The neutral ligand 2-(2-pyridylethyl)-2,3-dihydro-1*H*-isoindole, was synthesized by reacting 2-(2-aminoethyl)pyridine and  $\alpha,\alpha'$ -dibromo-*o*-xylene in THF in the presence of triethylamine as a base and purified by column chromatography using modified literature methods of Bonnett *et al.* (1983) & Meyers *et al.* (1995). The perchlorate salt of the protonated ligand (I) was formed by addition of one equivalent of an aqueous solution of  $\text{HClO}_4$  to a solution of the neutral ligand in methanol and allowing the resulting solution to stand yielded pale pink needles of (I).

### Refinement

The perchlorate anion is disordered over four conformations with occupancy factors of 0.438 (4), 0.270 (9), 0.155 (8), and 0.138 (5). These were constrained to adopt a tetrahedral geometry. This disorder and its modelling resulted in 245 restraints. The H atoms were idealized with an N—H distance of 0.91 Å and C—H distances were idealized at 0.93 (aromatic C—H), 0.96 ( $\text{CH}_3$ ), and 0.97 ( $\text{CH}_2$ ) Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  ( $1.5U_{\text{eq}}(\text{C})$  for the  $\text{CH}_3$  protons). Since a unique data set was collected there were no Friedel pairs and no attempt was made to establish the absolute structure.

### Figures




Fig. 1. The title compound showing the N—H···N bond (dashed bonds). Ellipsoids are drawn at the 20% probability level.




Fig. 2. The packing arrangement viewed down the  $a$  axis showing the N—H···O and intermolecular C—H···O interactions (dashed bonds).

# supplementary materials

---

## 2-[2-(2-Pyridyl)ethyl]isoindolinium perchlorate

### Crystal data

$C_{15}H_{17}N_2^+\cdot ClO_4^-$	$F_{000} = 680$
$M_r = 324.76$	$D_x = 1.403 \text{ Mg m}^{-3}$
Orthorhombic, $P2_12_12_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2ac 2ab	Cell parameters from 39 reflections
$a = 7.7941 (9) \text{ \AA}$	$\theta = 5.1\text{--}12.5^\circ$
$b = 11.7032 (19) \text{ \AA}$	$\mu = 0.27 \text{ mm}^{-1}$
$c = 16.850 (3) \text{ \AA}$	$T = 293 \text{ K}$
$V = 1537.0 (4) \text{ \AA}^3$	Needle, pale pink
$Z = 4$	$0.5 \times 0.2 \times 0.08 \text{ mm}$

### Data collection

Bruker P4	$R_{\text{int}} = 0.025$
diffractometer	
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 27.5^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.4^\circ$
$T = 293 \text{ K}$	$h = 0 \rightarrow 10$
$\omega$ scans	$k = 0 \rightarrow 15$
Absorption correction: $\psi$ scan (North <i>et al.</i> , 1968)	$l = 0 \rightarrow 21$
$T_{\text{min}} = 0.241$ , $T_{\text{max}} = 0.265$	3 standard reflections
2027 measured reflections	every 97 reflections
2001 independent reflections	intensity decay: <2%
1660 reflections with $I > 2\sigma(I)$	

### Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.044$	H-atom parameters constrained
$wR(F^2) = 0.120$	$w = 1/[\sigma^2(F_o^2) + (0.0626P)^2 + 0.2989P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} = 0.005$
2001 reflections	$\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
329 parameters	$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$
245 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cl	-0.45920 (11)	0.96522 (8)	0.73473 (5)	0.0643 (3)	
O1	-0.4598 (13)	1.0879 (4)	0.7432 (7)	0.093 (3)	0.438 (4)
O2	-0.5971 (10)	0.9347 (9)	0.6824 (5)	0.101 (3)	0.438 (4)
O3	-0.4861 (13)	0.9137 (8)	0.8101 (3)	0.074 (2)	0.438 (4)
O4	-0.3007 (9)	0.9305 (9)	0.7017 (6)	0.105 (3)	0.438 (4)
O1A	-0.497 (3)	0.8798 (16)	0.6772 (10)	0.099 (5)	0.155 (8)
O2A	-0.2811 (10)	0.9589 (19)	0.7554 (14)	0.100 (6)	0.155 (8)
O3A	-0.494 (3)	1.0759 (10)	0.7025 (15)	0.103 (5)	0.155 (8)
O4A	-0.561 (3)	0.948 (2)	0.8038 (8)	0.104 (5)	0.155 (8)
O1B	-0.466 (3)	0.9618 (18)	0.8191 (3)	0.100 (6)	0.138 (5)
O2B	-0.404 (3)	0.8561 (9)	0.7054 (11)	0.099 (5)	0.138 (5)
O3B	-0.341 (2)	1.0506 (14)	0.7094 (12)	0.100 (4)	0.138 (5)
O4B	-0.6256 (14)	0.9896 (18)	0.7036 (13)	0.104 (5)	0.138 (5)
O1C	-0.2804 (8)	0.9697 (14)	0.7176 (9)	0.088 (4)	0.270 (9)
O2C	-0.5525 (18)	0.9411 (15)	0.6637 (6)	0.111 (5)	0.270 (9)
O3C	-0.492 (2)	0.8776 (12)	0.7918 (8)	0.096 (4)	0.270 (9)
O4C	-0.5146 (19)	1.0728 (8)	0.7659 (10)	0.111 (4)	0.270 (9)
N1	-0.0054 (4)	0.8416 (2)	0.61178 (17)	0.0633 (7)	
N2	-0.0414 (3)	0.7346 (2)	0.75736 (15)	0.0546 (6)	
H2B	-0.0805	0.7936	0.7272	0.065 (10)*	
C1	-0.0438 (5)	0.9290 (3)	0.5639 (2)	0.0716 (9)	
H1A	-0.0860	0.9961	0.5861	0.084 (13)*	
C2	-0.0232 (5)	0.9233 (4)	0.4823 (2)	0.0762 (10)	
H2A	-0.0468	0.9865	0.4507	0.114 (16)*	
C3	0.0328 (5)	0.8230 (4)	0.4489 (2)	0.0798 (11)	
H3A	0.0460	0.8167	0.3943	0.077 (11)*	
C4	0.0691 (5)	0.7319 (4)	0.4976 (2)	0.0702 (9)	
H4A	0.1059	0.6629	0.4762	0.086 (13)*	
C5	0.0502 (5)	0.7442 (3)	0.57909 (19)	0.0609 (8)	
C6	0.0970 (5)	0.6493 (3)	0.6367 (2)	0.0677 (9)	
H6A	0.2089	0.6658	0.6591	0.083 (13)*	
H6B	0.1062	0.5782	0.6074	0.073 (11)*	
C7	-0.0297 (5)	0.6324 (3)	0.7044 (2)	0.0652 (8)	

## supplementary materials

---

H7A	-0.1422	0.6163	0.6825	0.069 (10)*
H7B	0.0053	0.5668	0.7356	0.080 (11)*
C8	-0.1673 (4)	0.7187 (3)	0.8247 (2)	0.0626 (8)
H8A	-0.2813	0.7439	0.8098	0.105 (15)*
H8B	-0.1724	0.6395	0.8414	0.086 (12)*
C9	-0.0935 (4)	0.7931 (3)	0.88894 (19)	0.0552 (7)
C10	-0.1673 (5)	0.8278 (3)	0.9602 (2)	0.0690 (9)
H10A	-0.2794	0.8076	0.9729	0.095 (14)*
C11	-0.0696 (6)	0.8932 (3)	1.0117 (2)	0.0743 (11)
H11A	-0.1162	0.9156	1.0600	0.069 (10)*
C12	0.0928 (6)	0.9252 (3)	0.9930 (2)	0.0721 (10)
H12A	0.1547	0.9704	1.0282	0.112 (16)*
C13	0.1681 (5)	0.8917 (3)	0.9219 (2)	0.0654 (8)
H13A	0.2787	0.9146	0.9088	0.124 (19)*
C14	0.0738 (4)	0.8232 (3)	0.87123 (18)	0.0546 (7)
C15	0.1273 (4)	0.7717 (3)	0.79305 (19)	0.0612 (8)
H15A	0.2032	0.7070	0.8009	0.064 (10)*
H15B	0.1841	0.8277	0.7597	0.079 (12)*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cl	0.0620 (5)	0.0772 (5)	0.0537 (4)	-0.0053 (5)	0.0037 (4)	-0.0008 (4)
O1	0.083 (6)	0.077 (4)	0.121 (7)	-0.012 (4)	-0.002 (6)	-0.003 (4)
O2	0.094 (6)	0.146 (8)	0.062 (4)	-0.043 (5)	-0.005 (4)	-0.008 (5)
O3	0.082 (5)	0.093 (6)	0.047 (3)	0.015 (5)	0.017 (3)	-0.004 (3)
O4	0.094 (5)	0.110 (7)	0.112 (6)	0.028 (5)	0.040 (5)	0.003 (6)
O1A	0.099 (10)	0.117 (10)	0.082 (8)	-0.002 (9)	-0.010 (8)	-0.028 (8)
O2A	0.084 (9)	0.092 (10)	0.126 (12)	-0.007 (9)	0.028 (10)	0.011 (11)
O3A	0.085 (9)	0.111 (9)	0.112 (11)	0.031 (8)	-0.016 (9)	0.023 (9)
O4A	0.096 (10)	0.133 (11)	0.085 (9)	-0.003 (10)	0.044 (8)	-0.022 (9)
O1B	0.100 (10)	0.121 (12)	0.079 (9)	0.001 (11)	0.015 (9)	-0.017 (9)
O2B	0.099 (10)	0.102 (9)	0.096 (9)	0.012 (9)	0.006 (9)	-0.013 (8)
O3B	0.087 (8)	0.095 (8)	0.120 (9)	-0.014 (8)	0.021 (8)	0.002 (8)
O4B	0.081 (9)	0.128 (10)	0.101 (10)	0.022 (9)	-0.013 (9)	0.000 (9)
O1C	0.078 (6)	0.079 (7)	0.107 (9)	-0.010 (5)	0.056 (6)	-0.025 (7)
O2C	0.123 (10)	0.150 (9)	0.059 (6)	-0.021 (9)	-0.013 (7)	-0.010 (7)
O3C	0.082 (7)	0.107 (9)	0.099 (8)	0.007 (7)	0.015 (7)	0.016 (7)
O4C	0.096 (8)	0.117 (7)	0.120 (8)	0.039 (6)	0.008 (7)	-0.025 (7)
N1	0.0600 (17)	0.0669 (17)	0.0630 (15)	0.0055 (13)	-0.0045 (14)	-0.0003 (13)
N2	0.0552 (14)	0.0559 (13)	0.0526 (13)	0.0009 (12)	-0.0042 (13)	0.0014 (11)
C1	0.066 (2)	0.069 (2)	0.080 (2)	-0.0005 (19)	-0.004 (2)	0.0048 (18)
C2	0.060 (2)	0.091 (3)	0.078 (2)	-0.004 (2)	0.0013 (19)	0.026 (2)
C3	0.0581 (19)	0.122 (3)	0.0588 (19)	0.006 (2)	0.0058 (18)	0.010 (2)
C4	0.061 (2)	0.090 (2)	0.0599 (18)	0.0087 (19)	-0.0008 (17)	-0.0076 (18)
C5	0.0543 (17)	0.0722 (19)	0.0561 (16)	0.0054 (17)	-0.0032 (16)	-0.0035 (15)
C6	0.075 (2)	0.070 (2)	0.0584 (18)	0.0152 (18)	-0.0067 (18)	-0.0095 (16)
C7	0.076 (2)	0.0574 (17)	0.0620 (17)	0.0011 (17)	-0.0066 (19)	-0.0045 (14)

C8	0.0541 (18)	0.0631 (19)	0.071 (2)	-0.0068 (16)	0.0066 (16)	0.0027 (16)
C9	0.0571 (18)	0.0510 (15)	0.0576 (17)	0.0052 (14)	0.0056 (14)	0.0081 (14)
C10	0.070 (2)	0.065 (2)	0.072 (2)	0.0034 (18)	0.0226 (19)	0.0092 (17)
C11	0.093 (3)	0.075 (2)	0.0546 (18)	0.013 (2)	0.015 (2)	-0.0035 (17)
C12	0.081 (3)	0.078 (2)	0.0577 (18)	0.0099 (19)	-0.0069 (19)	-0.0030 (18)
C13	0.0591 (19)	0.076 (2)	0.0605 (18)	0.0006 (18)	-0.0069 (16)	-0.0013 (17)
C14	0.0525 (17)	0.0614 (17)	0.0499 (15)	0.0047 (14)	0.0009 (13)	0.0068 (13)
C15	0.0502 (16)	0.078 (2)	0.0550 (16)	-0.0031 (17)	0.0003 (14)	-0.0035 (16)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

Cl—O4	1.415 (4)	C4—C5	1.389 (5)
Cl—O3	1.421 (4)	C4—H4A	0.9300
Cl—O4A	1.423 (5)	C5—C6	1.520 (5)
Cl—O1B	1.424 (5)	C6—C7	1.521 (5)
Cl—O1A	1.424 (5)	C6—H6A	0.9700
Cl—O1C	1.424 (5)	C6—H6B	0.9700
Cl—O3B	1.424 (5)	C7—H7A	0.9700
Cl—O4B	1.427 (5)	C7—H7B	0.9700
Cl—O3C	1.428 (5)	C8—C9	1.503 (5)
Cl—O2C	1.429 (5)	C8—H8A	0.9700
Cl—O3A	1.429 (5)	C8—H8B	0.9700
Cl—O4C	1.431 (5)	C9—C14	1.383 (5)
N1—C1	1.337 (4)	C9—C10	1.392 (4)
N1—C5	1.338 (4)	C10—C11	1.385 (5)
N2—C7	1.495 (4)	C10—H10A	0.9300
N2—C15	1.510 (4)	C11—C12	1.357 (6)
N2—C8	1.512 (4)	C11—H11A	0.9300
N2—H2B	0.9100	C12—C13	1.390 (5)
C1—C2	1.385 (6)	C12—H12A	0.9300
C1—H1A	0.9300	C13—C14	1.382 (5)
C2—C3	1.374 (6)	C13—H13A	0.9300
C2—H2A	0.9300	C14—C15	1.508 (4)
C3—C4	1.373 (6)	C15—H15A	0.9700
C3—H3A	0.9300	C15—H15B	0.9700
O4—Cl—O3	111.0 (4)	C7—C6—H6B	108.6
O4A—Cl—O1A	110.1 (5)	H6A—C6—H6B	107.6
O1B—Cl—O3B	110.0 (5)	N2—C7—C6	112.5 (3)
O1B—Cl—O4B	109.8 (5)	N2—C7—H7A	109.1
O3B—Cl—O4B	109.7 (5)	C6—C7—H7A	109.1
O1C—Cl—O3C	109.7 (4)	N2—C7—H7B	109.1
O1C—Cl—O2C	109.6 (4)	C6—C7—H7B	109.1
O3C—Cl—O2C	109.3 (4)	H7A—C7—H7B	107.8
O4A—Cl—O3A	109.5 (4)	C9—C8—N2	102.8 (2)
O1A—Cl—O3A	109.8 (4)	C9—C8—H8A	111.2
O1C—Cl—O4C	109.7 (4)	N2—C8—H8A	111.2
O3C—Cl—O4C	109.4 (4)	C9—C8—H8B	111.2
O2C—Cl—O4C	109.1 (4)	N2—C8—H8B	111.2
C1—N1—C5	118.4 (3)	H8A—C8—H8B	109.1

## supplementary materials

---

C7—N2—C15	114.5 (3)	C14—C9—C10	120.1 (3)
C7—N2—C8	112.9 (2)	C14—C9—C8	110.7 (3)
C15—N2—C8	107.5 (2)	C10—C9—C8	129.2 (3)
C7—N2—H2B	107.2	C11—C10—C9	118.3 (4)
C15—N2—H2B	107.2	C11—C10—H10A	120.9
C8—N2—H2B	107.2	C9—C10—H10A	120.9
N1—C1—C2	122.4 (4)	C12—C11—C10	121.3 (4)
N1—C1—H1A	118.8	C12—C11—H11A	119.3
C2—C1—H1A	118.8	C10—C11—H11A	119.3
C3—C2—C1	119.0 (4)	C11—C12—C13	121.0 (4)
C3—C2—H2A	120.5	C11—C12—H12A	119.5
C1—C2—H2A	120.5	C13—C12—H12A	119.5
C4—C3—C2	119.0 (4)	C14—C13—C12	118.1 (4)
C4—C3—H3A	120.5	C14—C13—H13A	120.9
C2—C3—H3A	120.5	C12—C13—H13A	120.9
C3—C4—C5	119.2 (4)	C13—C14—C9	121.0 (3)
C3—C4—H4A	120.4	C13—C14—C15	128.6 (3)
C5—C4—H4A	120.4	C9—C14—C15	110.3 (3)
N1—C5—C4	122.0 (3)	C14—C15—N2	102.8 (3)
N1—C5—C6	115.9 (3)	C14—C15—H15A	111.2
C4—C5—C6	122.1 (3)	N2—C15—H15A	111.2
C5—C6—C7	114.7 (3)	C14—C15—H15B	111.2
C5—C6—H6A	108.6	N2—C15—H15B	111.2
C7—C6—H6A	108.6	H15A—C15—H15B	109.1
C5—C6—H6B	108.6		
C5—N1—C1—C2	-2.1 (6)	N2—C8—C9—C10	168.3 (3)
N1—C1—C2—C3	2.5 (6)	C14—C9—C10—C11	0.5 (5)
C1—C2—C3—C4	-1.0 (6)	C8—C9—C10—C11	177.3 (3)
C2—C3—C4—C5	-0.8 (6)	C9—C10—C11—C12	1.4 (6)
C1—N1—C5—C4	0.2 (6)	C10—C11—C12—C13	-1.2 (6)
C1—N1—C5—C6	178.2 (3)	C11—C12—C13—C14	-0.9 (5)
C3—C4—C5—N1	1.2 (6)	C12—C13—C14—C9	2.8 (5)
C3—C4—C5—C6	-176.7 (4)	C12—C13—C14—C15	-177.3 (3)
N1—C5—C6—C7	43.7 (5)	C10—C9—C14—C13	-2.6 (5)
C4—C5—C6—C7	-138.3 (4)	C8—C9—C14—C13	-180.0 (3)
C15—N2—C7—C6	-56.6 (4)	C10—C9—C14—C15	177.5 (3)
C8—N2—C7—C6	179.9 (3)	C8—C9—C14—C15	0.1 (4)
C5—C6—C7—N2	-63.2 (4)	C13—C14—C15—N2	-165.4 (3)
C7—N2—C8—C9	150.7 (3)	C9—C14—C15—N2	14.4 (4)
C15—N2—C8—C9	23.4 (3)	C7—N2—C15—C14	-149.7 (3)
N2—C8—C9—C14	-14.6 (3)	C8—N2—C15—C14	-23.4 (3)

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N2—H2B···N1	0.91	2.11	2.769 (4)	129
N2—H2B···O4	0.91	2.39	3.197 (9)	148
N2—H2B···O2A	0.91	2.53	3.222 (19)	133
N2—H2B···O1C	0.91	2.59	3.390 (16)	147

C2—H2A···O1B <sup>i</sup>	0.93	2.30	3.062 (11)	139
C3—H3A···O1A <sup>ii</sup>	0.93	2.62	3.193 (10)	121
C7—H7B···O2A <sup>iii</sup>	0.97	2.50	3.232 (16)	132
C8—H8A···O3	0.97	2.55	3.383 (10)	144
C8—H8A···O2B	0.97	2.39	3.168 (19)	136
C8—H8A···O3C	0.97	2.29	3.189 (17)	154
C8—H8B···O4B <sup>iv</sup>	0.97	2.48	3.167 (17)	128
C11—H11A···O3A <sup>v</sup>	0.93	2.55	3.27 (2)	135
C11—H11A···O3B <sup>v</sup>	0.93	2.57	3.47 (2)	162
C13—H13A···O3 <sup>vi</sup>	0.93	2.48	3.299 (9)	148
C13—H13A···O4A <sup>vi</sup>	0.93	2.20	2.976 (8)	140
C13—H13A···O1B <sup>vi</sup>	0.93	2.56	3.44 (2)	158
C13—H13A···O3C <sup>vi</sup>	0.93	2.70	3.444 (17)	138
C15—H15A···O1 <sup>iii</sup>	0.97	2.55	3.423 (9)	150
C15—H15B···O2 <sup>vi</sup>	0.97	2.48	3.425 (10)	163
C15—H15B···O4A <sup>vi</sup>	0.97	2.55	3.194 (19)	124

Symmetry codes: (i)  $-x-1/2, -y+2, z-1/2$ ; (ii)  $x+1/2, -y+3/2, -z+1$ ; (iii)  $-x, y-1/2, -z+3/2$ ; (iv)  $-x-1, y-1/2, -z+3/2$ ; (v)  $-x-1/2, -y+2, z+1/2$ ; (vi)  $x+1, y, z$ .

## supplementary materials

---

Fig. 1




Fig. 2

