

1-Benzyl-3-phenylimidazolium hexafluorophosphate

Ping Jiang

School of Chemistry and Chemical Engineering, China West Normal University,
Nanchong 637002, People's Republic of China
Correspondence e-mail: jphdp868@126.com


Received 29 July 2009; accepted 11 August 2009

Key indicators: single-crystal X-ray study; $T = 93\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.035; wR factor = 0.084; data-to-parameter ratio = 12.8.

in the title compound, $\text{C}_{16}\text{H}_{15}\text{N}_2^+\cdot\text{PF}_6^-$, a precursor of *N*-heterocyclic carbene, the phenyl and benzyl rings are twisted away from the central imidazolium ring system, making dihedral angles of $70.30(8)$ and $32.03(10)^\circ$, respectively. The crystal structure is stabilized by C—H \cdots F hydrogen bonds. Furthermore, P—F \cdots π interactions involving imidazolium rings are observed [$\text{F}\cdots\pi = 2.9857(16)$, $\text{P}\cdots\pi = 4.1630(16)\text{ \AA}$, $\text{P}\cdots\pi = 127.92(6)^\circ$].

Related literature

The first stable *N*-heterocyclic carbene was isolated by Arduengo *et al.* (1991). For the synthesis, see: Liu *et al.* (2003). For related structures, see: Wan *et al.* (2008). For related structures, see: Newman *et al.* (2007); Herrmann (2002); Yang *et al.* (2009).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{15}\text{N}_2^+\cdot\text{PF}_6^-$
 $M_r = 380.27$
Triclinic, $P\bar{1}$

$a = 9.221(2)\text{ \AA}$
 $b = 10.046(3)\text{ \AA}$
 $c = 10.108(2)\text{ \AA}$

$\alpha = 110.733(2)^\circ$
 $\beta = 91.969(2)^\circ$
 $\gamma = 110.315(2)^\circ$
 $V = 807.9(3)\text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.24\text{ mm}^{-1}$
 $T = 93\text{ K}$
 $0.43 \times 0.40 \times 0.37\text{ mm}$

Data collection

Rigaku SPIDER diffractometer
Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.905$, $T_{\max} = 0.919$

4739 measured reflections
2902 independent reflections
2361 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.035$
 $wR(F^2) = 0.084$
 $S = 1.00$
2902 reflections

226 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.22\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C7—H7 \cdots F1 ⁱ	0.95	2.50	3.099 (2)	121
C8—H8 \cdots F6 ⁱ	0.95	2.50	3.392 (2)	156
C9—H9 \cdots F5 ⁱⁱ	0.95	2.34	3.247 (2)	159
C10—H10A \cdots F4 ⁱⁱ	0.99	2.49	3.444 (2)	161
C10—H10B \cdots F3 ⁱⁱⁱ	0.99	2.49	3.455 (3)	164

Symmetry codes: (i) $x - 1, y - 1, z$; (ii) $x - 1, y - 1, z - 1$; (iii) $-x + 1, -y + 1, -z + 1$.

Data collection: *RAPID-AUTO* (Rigaku/MSC, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Fund Projects of Sichuan Educational department (grant No. 2005 A104).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2854).

References

- Arduengo, A. J., Harlow, R. L. & Kline, M. (1991). *J. Am. Chem. Soc.* **113**, 361–363.
- Herrmann, W. A. (2002). *Angew. Chem. Int. Ed.* **41**, 1290–1309.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Liu, J. P., Chen, J. B., Zhao, J. F., Zhao, Y. H., Li, L. & Zhang, H. B. (2003). *Synthesis*, pp. 2661–2666.
- Newman, C. P., Clarkson, G. J. & Rourke, J. P. (2007). *J. Organomet. Chem.* **692**, 4962–4968.
- Rigaku/MSC (2004). *RAPID-AUTO* and *CrystalStructure*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wan, Y., Xin, H., Chen, X., Xu, H. & Wu, H. (2008). *Acta Cryst. E* **64**, o2159.
- Yang, X. D., Zeng, X. H., Zhang, Y. L., Qing, C., Song, W. J., Li, L. & Zhang, H. B. (2009). *Bioorg. Med. Chem. Lett.* **19**, 1892–1895.

supplementary materials

Acta Cryst. (2009). E65, o2177 [doi:10.1107/S1600536809031584]

1-Benzyl-3-phenylimidazolium hexafluorophosphate

P. Jiang

Comment

Initiated by the isolation of the first stable N-heterocyclic carbene (NHC) by Arduengo *et al.* (1991), numerous stable NHC ligands have been prepared. 1,3-Disubstitutedimidazolium salts are considerable good precursor for synthesis of transition metal NHCs. In addition, the study of biological activities of imidazolium salts have been reported during these years. We report herein the synthesis and crystal structure of the title compound (I).

In (I), bond lengths and angles in title molecule (Fig. 1) are normal. The phenyl ring make dihedral angles with the benzyl ring and the imidazolium ring of 70.30 (8) $^{\circ}$ and 32.03 (10) $^{\circ}$, respectively.

The crystal structure is stabilized by C—H \cdots F hydrogen bonds (Table 1). Furthermore, P—F \cdots π interactions involving imidazolium rings [$F1\cdots Cg1^{iv} = 2.9857$ (16) Å, $P1\cdots Cg1^{iv} = 4.1630$ (16) Å, $P1—F1\cdots Cg1^{iv} = 127.92$ (6) $^{\circ}$, where $Cg1$ is a centroid of the N1/N2/C7—C9 ring; symmetry code: (iv) $-x+2, -y+1, -z+1$] are observed.

Experimental

The title compound was prepared according to the reported procedure of Liu *et al.* (2003). Colourless single crystals suitable for X-ray diffraction were obtained by recrystallization from dichloromethane and pPetroleum ether.

Refinement

H atoms were placed in calculated positions with C—H = 0.95–0.9900 Å, and refined in riding mode with $U_{iso}(H) = 1.2U_{eq}(C)$.

Figures





Fig. 1. The molecular structure of the title compound, showing 30% probability displacement ellipsoids and the atomic numbering.

1-Benzyl-3-phenylimidazolium hexafluorophosphate

Crystal data



$Z = 2$

$M_r = 380.27$

$F_{000} = 388$

Triclinic, $P\bar{1}$

$D_x = 1.563 \text{ Mg m}^{-3}$

Hall symbol: -P 1

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

supplementary materials

$a = 9.221 (2) \text{ \AA}$	Cell parameters from 2516 reflections
$b = 10.046 (3) \text{ \AA}$	$\theta = 3.1\text{--}27.5^\circ$
$c = 10.108 (2) \text{ \AA}$	$\mu = 0.24 \text{ mm}^{-1}$
$\alpha = 110.733 (2)^\circ$	$T = 93 \text{ K}$
$\beta = 91.969 (2)^\circ$	Block, colourless
$\gamma = 110.315 (2)^\circ$	$0.43 \times 0.40 \times 0.37 \text{ mm}$
$V = 807.9 (3) \text{ \AA}^3$	

Data collection

Rigaku SPIDER diffractometer	2902 independent reflections
Radiation source: Rotating Anode	2361 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.017$
$T = 93 \text{ K}$	$\theta_{\text{max}} = 25.5^\circ$
ω scans	$\theta_{\text{min}} = 3.1^\circ$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$h = -11 \rightarrow 7$
$T_{\text{min}} = 0.905, T_{\text{max}} = 0.919$	$k = -11 \rightarrow 12$
4739 measured reflections	$l = -12 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.035$	H-atom parameters constrained
$wR(F^2) = 0.084$	$w = 1/[\sigma^2(F_o^2) + (0.043P)^2 + 0.066P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.00$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2902 reflections	$\Delta\rho_{\text{max}} = 0.22 \text{ e \AA}^{-3}$
226 parameters	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
P1	1.23197 (5)	1.00569 (5)	0.79232 (5)	0.02127 (14)
F1	1.38948 (11)	1.06413 (12)	0.73171 (11)	0.0297 (3)
F2	1.07593 (13)	0.94747 (15)	0.85307 (13)	0.0444 (3)
F3	1.16399 (13)	1.10464 (14)	0.73467 (12)	0.0380 (3)
F4	1.30246 (13)	0.90834 (14)	0.85235 (12)	0.0381 (3)
F5	1.30920 (14)	1.15038 (13)	0.94289 (11)	0.0427 (3)
F6	1.15883 (14)	0.86186 (13)	0.64242 (12)	0.0433 (3)
N1	0.47043 (15)	0.20046 (16)	0.35708 (14)	0.0188 (3)
N2	0.28468 (15)	-0.02584 (16)	0.25378 (14)	0.0192 (3)
C1	0.5504 (2)	0.4242 (2)	0.29382 (18)	0.0235 (4)
H1	0.4593	0.3740	0.2211	0.028*
C2	0.6550 (2)	0.5713 (2)	0.31694 (19)	0.0255 (4)
H2	0.6367	0.6216	0.2581	0.031*
C3	0.7858 (2)	0.6454 (2)	0.42503 (18)	0.0241 (4)
H3	0.8553	0.7475	0.4422	0.029*
C4	0.8153 (2)	0.5709 (2)	0.50786 (19)	0.0270 (4)
H4	0.9061	0.6213	0.5810	0.032*
C5	0.7131 (2)	0.4229 (2)	0.48495 (19)	0.0243 (4)
H5	0.7337	0.3714	0.5416	0.029*
C6	0.58132 (19)	0.3514 (2)	0.37897 (18)	0.0191 (4)
C7	0.4319 (2)	0.1417 (2)	0.46130 (18)	0.0218 (4)
H7	0.4787	0.1912	0.5602	0.026*
C8	0.3160 (2)	0.0014 (2)	0.39716 (18)	0.0223 (4)
H8	0.2652	-0.0662	0.4423	0.027*
C9	0.37939 (19)	0.0950 (2)	0.23205 (18)	0.0194 (4)
H9	0.3822	0.1051	0.1421	0.023*
C10	0.1627 (2)	-0.1631 (2)	0.14214 (19)	0.0233 (4)
H10A	0.1772	-0.1595	0.0467	0.028*
H10B	0.0580	-0.1612	0.1581	0.028*
C11	0.16757 (19)	-0.3114 (2)	0.14213 (17)	0.0197 (4)
C12	0.28553 (19)	-0.3616 (2)	0.08983 (18)	0.0224 (4)
H12	0.3691	-0.2983	0.0598	0.027*
C13	0.2811 (2)	-0.5030 (2)	0.08160 (18)	0.0254 (4)
H13	0.3614	-0.5368	0.0453	0.030*
C14	0.1600 (2)	-0.5965 (2)	0.12598 (18)	0.0255 (4)
H14	0.1569	-0.6941	0.1195	0.031*
C15	0.0439 (2)	-0.5460 (2)	0.17967 (18)	0.0248 (4)
H15	-0.0385	-0.6087	0.2113	0.030*
C16	0.04776 (19)	-0.4049 (2)	0.18733 (17)	0.0219 (4)
H16	-0.0325	-0.3713	0.2240	0.026*

Atomic displacement parameters (\AA^2)

$$U^{11} \quad U^{22} \quad U^{33} \quad U^{12} \quad U^{13} \quad U^{23}$$

supplementary materials

P1	0.0232 (3)	0.0215 (3)	0.0217 (3)	0.0089 (2)	0.0073 (2)	0.0108 (2)
F1	0.0233 (5)	0.0358 (7)	0.0297 (6)	0.0075 (5)	0.0084 (5)	0.0159 (5)
F2	0.0366 (7)	0.0571 (9)	0.0629 (8)	0.0237 (6)	0.0301 (6)	0.0422 (7)
F3	0.0390 (7)	0.0481 (8)	0.0470 (7)	0.0250 (6)	0.0148 (6)	0.0323 (7)
F4	0.0498 (7)	0.0469 (8)	0.0437 (7)	0.0330 (6)	0.0230 (6)	0.0315 (6)
F5	0.0644 (8)	0.0339 (7)	0.0229 (6)	0.0196 (6)	0.0073 (6)	0.0029 (6)
F6	0.0445 (7)	0.0274 (7)	0.0348 (7)	-0.0020 (6)	-0.0008 (5)	0.0025 (6)
N1	0.0206 (7)	0.0181 (8)	0.0174 (8)	0.0075 (6)	0.0029 (6)	0.0066 (7)
N2	0.0197 (7)	0.0187 (8)	0.0184 (8)	0.0076 (6)	0.0021 (6)	0.0063 (7)
C1	0.0236 (9)	0.0251 (10)	0.0214 (10)	0.0088 (8)	0.0026 (8)	0.0094 (8)
C2	0.0305 (10)	0.0247 (10)	0.0259 (10)	0.0122 (8)	0.0077 (8)	0.0134 (9)
C3	0.0251 (9)	0.0209 (10)	0.0239 (10)	0.0065 (8)	0.0083 (8)	0.0081 (8)
C4	0.0244 (9)	0.0275 (11)	0.0228 (10)	0.0061 (8)	-0.0002 (8)	0.0070 (9)
C5	0.0264 (9)	0.0238 (10)	0.0227 (10)	0.0086 (8)	0.0015 (8)	0.0103 (9)
C6	0.0211 (9)	0.0178 (9)	0.0177 (9)	0.0081 (7)	0.0058 (7)	0.0053 (8)
C7	0.0308 (10)	0.0232 (10)	0.0148 (9)	0.0128 (8)	0.0059 (7)	0.0086 (8)
C8	0.0292 (10)	0.0234 (10)	0.0174 (9)	0.0121 (8)	0.0070 (8)	0.0093 (8)
C9	0.0203 (8)	0.0216 (10)	0.0164 (9)	0.0088 (7)	0.0023 (7)	0.0068 (8)
C10	0.0210 (9)	0.0238 (10)	0.0216 (10)	0.0059 (8)	-0.0007 (7)	0.0078 (8)
C11	0.0202 (8)	0.0204 (10)	0.0146 (9)	0.0059 (7)	-0.0017 (7)	0.0049 (8)
C12	0.0187 (9)	0.0248 (10)	0.0218 (10)	0.0060 (8)	0.0038 (7)	0.0092 (8)
C13	0.0233 (9)	0.0292 (11)	0.0229 (10)	0.0120 (8)	-0.0007 (8)	0.0079 (9)
C14	0.0293 (10)	0.0209 (10)	0.0218 (10)	0.0065 (8)	-0.0052 (8)	0.0073 (8)
C15	0.0225 (9)	0.0265 (11)	0.0205 (10)	0.0019 (8)	-0.0006 (8)	0.0116 (9)
C16	0.0200 (9)	0.0264 (10)	0.0147 (9)	0.0063 (8)	0.0007 (7)	0.0057 (8)

Geometric parameters (\AA , $^\circ$)

P1—F2	1.5875 (11)	C5—C6	1.378 (2)
P1—F3	1.5933 (11)	C5—H5	0.9500
P1—F6	1.5942 (12)	C7—C8	1.345 (2)
P1—F1	1.5978 (11)	C7—H7	0.9500
P1—F5	1.6052 (12)	C8—H8	0.9500
P1—F4	1.6066 (11)	C9—H9	0.9500
N1—C9	1.336 (2)	C10—C11	1.505 (2)
N1—C7	1.380 (2)	C10—H10A	0.9900
N1—C6	1.437 (2)	C10—H10B	0.9900
N2—C9	1.323 (2)	C11—C16	1.389 (2)
N2—C8	1.376 (2)	C11—C12	1.396 (2)
N2—C10	1.474 (2)	C12—C13	1.379 (2)
C1—C2	1.385 (2)	C12—H12	0.9500
C1—C6	1.389 (2)	C13—C14	1.391 (2)
C1—H1	0.9500	C13—H13	0.9500
C2—C3	1.383 (2)	C14—C15	1.386 (2)
C2—H2	0.9500	C14—H14	0.9500
C3—C4	1.380 (2)	C15—C16	1.380 (2)
C3—H3	0.9500	C15—H15	0.9500
C4—C5	1.386 (2)	C16—H16	0.9500
C4—H4	0.9500		

F2—P1—F3	90.46 (6)	C5—C6—C1	121.20 (17)
F2—P1—F6	90.77 (7)	C5—C6—N1	120.21 (15)
F3—P1—F6	90.53 (7)	C1—C6—N1	118.58 (15)
F2—P1—F1	179.73 (6)	C8—C7—N1	107.46 (15)
F3—P1—F1	89.79 (6)	C8—C7—H7	126.3
F6—P1—F1	89.34 (6)	N1—C7—H7	126.3
F2—P1—F5	90.23 (7)	C7—C8—N2	107.13 (15)
F3—P1—F5	90.14 (7)	C7—C8—H8	126.4
F6—P1—F5	178.80 (7)	N2—C8—H8	126.4
F1—P1—F5	89.66 (6)	N2—C9—N1	108.96 (14)
F2—P1—F4	89.83 (6)	N2—C9—H9	125.5
F3—P1—F4	179.11 (7)	N1—C9—H9	125.5
F6—P1—F4	90.31 (7)	N2—C10—C11	112.54 (14)
F1—P1—F4	89.92 (6)	N2—C10—H10A	109.1
F5—P1—F4	89.02 (7)	C11—C10—H10A	109.1
C9—N1—C7	107.81 (14)	N2—C10—H10B	109.1
C9—N1—C6	125.48 (14)	C11—C10—H10B	109.1
C7—N1—C6	126.60 (15)	H10A—C10—H10B	107.8
C9—N2—C8	108.63 (14)	C16—C11—C12	118.95 (16)
C9—N2—C10	124.79 (14)	C16—C11—C10	119.61 (15)
C8—N2—C10	126.53 (14)	C12—C11—C10	121.34 (15)
C2—C1—C6	118.79 (17)	C13—C12—C11	120.18 (16)
C2—C1—H1	120.6	C13—C12—H12	119.9
C6—C1—H1	120.6	C11—C12—H12	119.9
C3—C2—C1	120.50 (17)	C12—C13—C14	120.50 (17)
C3—C2—H2	119.8	C12—C13—H13	119.8
C1—C2—H2	119.8	C14—C13—H13	119.8
C4—C3—C2	119.89 (17)	C15—C14—C13	119.44 (17)
C4—C3—H3	120.1	C15—C14—H14	120.3
C2—C3—H3	120.1	C13—C14—H14	120.3
C3—C4—C5	120.38 (17)	C16—C15—C14	120.11 (16)
C3—C4—H4	119.8	C16—C15—H15	119.9
C5—C4—H4	119.8	C14—C15—H15	119.9
C6—C5—C4	119.21 (17)	C15—C16—C11	120.81 (16)
C6—C5—H5	120.4	C15—C16—H16	119.6
C4—C5—H5	120.4	C11—C16—H16	119.6
C6—C1—C2—C3	-1.4 (3)	C8—N2—C9—N1	-0.49 (18)
C1—C2—C3—C4	1.9 (3)	C10—N2—C9—N1	177.44 (14)
C2—C3—C4—C5	-1.0 (3)	C7—N1—C9—N2	0.77 (18)
C3—C4—C5—C6	-0.3 (3)	C6—N1—C9—N2	-175.77 (14)
C4—C5—C6—C1	0.8 (3)	C9—N2—C10—C11	132.27 (16)
C4—C5—C6—N1	-177.50 (15)	C8—N2—C10—C11	-50.2 (2)
C2—C1—C6—C5	0.0 (3)	N2—C10—C11—C16	109.96 (17)
C2—C1—C6—N1	178.40 (15)	N2—C10—C11—C12	-73.7 (2)
C9—N1—C6—C5	-151.02 (16)	C16—C11—C12—C13	0.9 (2)
C7—N1—C6—C5	33.1 (2)	C10—C11—C12—C13	-175.41 (16)
C9—N1—C6—C1	30.6 (2)	C11—C12—C13—C14	-0.4 (3)
C7—N1—C6—C1	-145.30 (16)	C12—C13—C14—C15	-0.4 (3)
C9—N1—C7—C8	-0.76 (18)	C13—C14—C15—C16	0.7 (2)

supplementary materials

C6—N1—C7—C8	175.73 (15)	C14—C15—C16—C11	-0.2 (2)
N1—C7—C8—N2	0.46 (18)	C12—C11—C16—C15	-0.6 (2)
C9—N2—C8—C7	0.01 (18)	C10—C11—C16—C15	175.78 (15)
C10—N2—C8—C7	-177.88 (15)		

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
C7—H7···F1 ⁱ	0.95	2.50	3.099 (2)	121
C8—H8···F6 ⁱ	0.95	2.50	3.392 (2)	156
C9—H9···F5 ⁱⁱ	0.95	2.34	3.247 (2)	159
C10—H10A···F4 ⁱⁱ	0.99	2.49	3.444 (2)	161
C10—H10B···F3 ⁱⁱⁱ	0.99	2.49	3.455 (3)	164

Symmetry codes: (i) $x-1, y-1, z$; (ii) $x-1, y-1, z-1$; (iii) $-x+1, -y+1, -z+1$.

Fig. 1

