

Poly[[aqua(μ_2 -oxalato)(μ_2 -2-oxido-pyridinium-3-carboxylato)holmium(III)] monohydrate]

Hui-Lan Zhu,^a Hui-Ling Lai,^a Lu Han,^a Yi-Fan Luo^{a*} and Rong-Hua Zeng^{a,b}

^aSchool of Chemistry and Environment, South China Normal University, Guangzhou 510006, People's Republic of China, and ^bKey Laboratory of Technology on Electrochemical Energy Storage and Power Generation in Guangdong Universities, South China Normal University, Guangzhou 510006, People's Republic of China
Correspondence e-mail: luoyf2004@yahoo.com.cn


Received 25 August 2009; accepted 8 September 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(C-C) = 0.011$ Å; R factor = 0.034; wR factor = 0.090; data-to-parameter ratio = 10.8.

In the title complex, $\{[Ho(C_2O_4)(C_6H_4NO_3)(H_2O)] \cdot (H_2O)\}_n$, the Ho^{III} ion is coordinated by three O atoms from two 2-oxidopyridinium-3-carboxylate ligands, four O atoms from two oxalate ligands and one water molecule in a distorted bicapped trigonal-prismatic geometry. The 2-oxidopyridinium-3-carboxylate and oxalate ligands link the Ho^{III} ions into a layer in (100). These layers are further connected by intermolecular O—H···O hydrogen bonds involving the coordinated water molecules to assemble a three-dimensional supramolecular network. The uncoordinated water molecule is involved in N—H···O and O—H···O hydrogen bonds within the layer.

Related literature

For general background to molecular self-assembly of supramolecular architectures, see: Deng *et al.* (2008); Zeng *et al.* (2007). For 2-oxidopyridinium-3-carboxylic acid and oxalic acid as building blocks, see: Huang *et al.* (2009).



Experimental

Crystal data

$[Ho(C_2O_4)(C_6H_4NO_3)(H_2O)] \cdot H_2O$	$\gamma = 80.359 (2)^\circ$
$M_r = 427.08$	$V = 562.01 (16) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.5391 (11) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.5305 (16) \text{ \AA}$	$\mu = 7.09 \text{ mm}^{-1}$
$c = 9.7391 (16) \text{ \AA}$	$T = 296 \text{ K}$
$\alpha = 71.810 (2)^\circ$	$0.20 \times 0.18 \times 0.17 \text{ mm}$
$\beta = 78.862 (2)^\circ$	

Data collection

Bruker APEXII CCD diffractometer	2884 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	1983 independent reflections
$T_{min} = 0.332$, $T_{max} = 0.379$	1862 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.090$	$\Delta\rho_{\max} = 2.75 \text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\min} = -2.19 \text{ e \AA}^{-3}$
1983 reflections	
184 parameters	
6 restraints	

Table 1
Selected bond lengths (Å).

Ho1—O1	2.354 (5)	Ho1—O5 ⁱⁱ	2.364 (5)
Ho1—O2 ⁱ	2.348 (5)	Ho1—O6	2.394 (5)
Ho1—O3	2.279 (5)	Ho1—O7 ⁱⁱⁱ	2.392 (5)
Ho1—O4	2.406 (4)	Ho1—O1W	2.348 (5)

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $-x, -y + 1, -z$; (iii) $-x, -y + 1, -z + 1$.

Table 2
Hydrogen-bond geometry (Å, °).

$D—H \cdots A$	$D—H$	$H \cdots A$	$D \cdots A$	$D—H \cdots A$
O1W—H1W···O2 ^{iv}	0.84 (6)	2.02 (6)	2.737 (7)	142 (8)
O1W—H2W···O4 ^v	0.85 (7)	1.96 (4)	2.749 (7)	154 (8)
O2W—H3W···O6	0.84 (7)	2.36 (11)	2.882 (10)	121 (8)
O2W—H4W···O1	0.86 (7)	2.27 (3)	3.082 (8)	157 (9)
N1—H1···O2W ^{vi}	0.86	1.99	2.781 (10)	154

Symmetry codes: (iv) $-x + 1, -y, -z + 1$; (v) $x + 1, y, z$; (vi) $x, y, z - 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Chan Xue Yan Cooperative Special Project of Guangdong Province and the Ministry of Science and Technology of China (project No. 2007A090302046), the Project of Science and Technology of Guangdong Province (project No. 2007A020200002-4) and the Natural Science Foundation of Guangdong Province (No. 9151063101000037) for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2226).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Deng, H., Qiu, Y.-C., Li, Y.-H., Liu, Z.-H., Zeng, R.-H., Zeller, M. & Batten, S. R. (2008). *Chem. Commun.* pp. 2239–2241.
Huang, C.-D., Huang, J.-X., Wu, Y.-Y., Lian, Y.-Y. & Zeng, R.-H. (2009). *Acta Cryst. E*65, m177–m178.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A*64, 112–122.
Zeng, R.-H., Qiu, Y.-C., Cai, Y.-P., Wu, J.-Z. & Deng, H. (2007). *Acta Cryst. E*63, m1666.

supplementary materials

Acta Cryst. (2009). E65, m1217-m1218 [doi:10.1107/S1600536809036344]

Poly[[aqua(μ_2 -oxalato)(μ_2 -2-oxidopyridinium-3-carboxylato)holmium(III)] monohydrate]

H.-L. Zhu, H.-L. Lai, L. Han, Y.-F. Luo and R.-H. Zeng

Comment

Molecular self-assembly of supramolecular architectures has received much attention during recent decades (Deng *et al.*, 2008; Zeng *et al.*, 2007). The structures and properties of such systems depend on the coordination and geometric preferences of both the central metal ions and the bridging building blocks, as well as the influence of weaker non-covalent interactions, such as hydrogen bonds and π – π stacking interactions. As a building block, 2-oxynicotinic acid and oxalic acid are good ligands with multifunctional coordination sites providing a high likelihood for the generation of structures with high dimensions (Huang *et al.*, 2009). Recently, we obtained the title coordination polymer, which was synthesized under hydrothermal conditions.

In the title compound (Fig. 1), the Ho^{III} centre is eight-coordinated by seven O atoms from two 2-oxynicotinate ligands and two oxalate ligands, and by one water molecule in a distorted bicapped trigonal prismatic geometry, with Ho—O distances and O—Ho—O angles ranging from 2.279 (5) to 2.406 (4) Å and 67.30 (17) to 148.26 (17) $^\circ$, respectively (Table 1). The carboxylate groups of the 2-oxynicotinates and oxalates act as bridging ligands, linking the Ho centres into a layer parallel to the (100) plane (Fig. 2). The layers are further connected by intermolecular O—H···O hydrogen bonds involving the coordinated water molecules into a three-dimensional supramolecular network (Table 2). The uncoordinated water molecule is involved in N—H···O and O—H···O hydrogen bonds within the layer (Table 2).

Experimental

A mixture of Ho₂O₃ (0.375 g, 1 mmol), 2-oxynicotinic acid (0.127 g, 1 mmol), oxalic acid (0.09 g, 1 mmol), water (10 ml) in the presence of HNO₃ (0.024 g, 0.385 mmol) was stirred vigorously for 20 min and then sealed in a Teflon-lined stainless-steel autoclave (20 ml capacity). The autoclave was heated and maintained at 446 K for 2 d, and then cooled to room temperature at 5 K h⁻¹, giving colourless block crystals.

Refinement

Water H atoms were tentatively located in difference Fourier maps and refined with distance restraints of O—H = 0.84 (1) and H···H = 1.35 (1) Å, and with a fixed $U_{\text{iso}}(\text{H}) = 0.064 \text{ \AA}^2$. H atoms attached to C and N atoms were placed at calculated positions and treated as riding on their parent atoms, with C—H = 0.93 and N—H = 0.86 Å and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$. The highest residual electron density was found 0.94 Å from Ho1 and the deepest hole 0.91 Å from Ho1.

supplementary materials

Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity. [Symmetry codes: (i) -x, -y, 1 - z; (ii) -x, 1 - y, -z; (iii) -x, 1 - y, 1 - z.]




Fig. 2. View of a layered network of the title compound.

Poly[[aqua(μ_2 -oxalato)(μ_2 -2-oxidopyridinium-3-carboxylato)holmium(III)] monohydrate]

Crystal data

[Ho(C ₂ O ₄)(C ₆ H ₄ NO ₃)(H ₂ O)]·H ₂ O	Z = 2
M _r = 427.08	F ₀₀₀ = 404
Triclinic, P <bar{1}< td=""><td>D_x = 2.524 Mg m⁻³</td></bar{1}<>	D _x = 2.524 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 6.5391 (11) Å	Cell parameters from 2285 reflections
b = 9.5305 (16) Å	θ = 2.2–28.1°
c = 9.7391 (16) Å	μ = 7.09 mm ⁻¹
α = 71.810 (2)°	T = 296 K
β = 78.862 (2)°	Block, colourless
γ = 80.359 (2)°	0.20 × 0.18 × 0.17 mm
V = 562.01 (16) Å ³	

Data collection

Bruker APEXII CCD diffractometer	1983 independent reflections
Radiation source: fine-focus sealed tube	1862 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.018$
T = 296 K	$\theta_{\text{max}} = 25.2^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.2^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -7 \rightarrow 6$
$T_{\text{min}} = 0.332$, $T_{\text{max}} = 0.379$	$k = -11 \rightarrow 11$
2884 measured reflections	$l = -11 \rightarrow 10$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.034$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.090$	$w = 1/[\sigma^2(F_o^2) + (0.0695P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\max} < 0.001$
1983 reflections	$\Delta\rho_{\max} = 2.75 \text{ e \AA}^{-3}$
184 parameters	$\Delta\rho_{\min} = -2.19 \text{ e \AA}^{-3}$
6 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Ho1	0.09083 (4)	0.29059 (3)	0.30045 (3)	0.01784 (15)
O4	-0.1655 (7)	0.3863 (5)	0.1370 (5)	0.0219 (10)
O3	0.1549 (8)	0.1295 (6)	0.1626 (5)	0.0311 (11)
O6	0.1351 (9)	0.3228 (5)	0.5267 (6)	0.0338 (12)
O1	0.2127 (8)	0.0581 (5)	0.4520 (5)	0.0251 (10)
O5	-0.2292 (7)	0.5458 (5)	-0.0767 (5)	0.0258 (10)
N1	0.2729 (11)	-0.0557 (8)	0.0581 (7)	0.0366 (15)
H1	0.2407	0.0054	-0.0227	0.044*
C6	0.2307 (12)	-0.0067 (8)	0.1802 (8)	0.0266 (15)
C8	0.0606 (11)	0.4422 (8)	0.5578 (8)	0.0261 (15)
C5	0.3624 (14)	-0.1948 (10)	0.0574 (10)	0.044 (2)
H5	0.3862	-0.2213	-0.0290	0.052*
C4	0.4170 (14)	-0.2947 (9)	0.1807 (10)	0.048 (3)
H4	0.4851	-0.3882	0.1795	0.057*
C3	0.3695 (13)	-0.2550 (9)	0.3094 (10)	0.0396 (19)
H3	0.3999	-0.3250	0.3960	0.048*
C2	0.2776 (10)	-0.1133 (7)	0.3126 (7)	0.0251 (15)
C1	0.2306 (10)	-0.0747 (7)	0.4529 (7)	0.0236 (14)
C7	-0.1147 (10)	0.4807 (7)	0.0161 (7)	0.0190 (13)
O2W	0.2278 (13)	0.0635 (9)	0.7646 (8)	0.0621 (19)
O2	0.2125 (7)	-0.1813 (5)	0.5703 (5)	0.0262 (10)
O7	0.0730 (9)	0.4748 (6)	0.6691 (5)	0.0358 (13)
O1W	0.4526 (8)	0.2970 (7)	0.2822 (6)	0.0427 (15)
H3W	0.164 (16)	0.149 (7)	0.758 (10)	0.064*
H4W	0.203 (16)	0.041 (10)	0.691 (6)	0.064*
H1W	0.508 (12)	0.263 (11)	0.359 (5)	0.064*
H2W	0.549 (10)	0.332 (11)	0.215 (6)	0.064*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Ho1	0.0166 (2)	0.0197 (2)	0.0158 (2)	-0.00221 (12)	-0.00168 (12)	-0.00363 (13)

supplementary materials

O4	0.018 (2)	0.027 (2)	0.017 (2)	-0.0088 (18)	-0.0019 (18)	0.0028 (19)
O3	0.039 (3)	0.031 (3)	0.020 (2)	0.002 (2)	-0.005 (2)	-0.007 (2)
O6	0.046 (3)	0.023 (3)	0.034 (3)	0.010 (2)	-0.013 (2)	-0.013 (2)
O1	0.027 (3)	0.024 (2)	0.022 (2)	-0.0029 (19)	-0.0024 (19)	-0.0050 (19)
O5	0.016 (2)	0.035 (3)	0.023 (2)	-0.0035 (19)	-0.0071 (19)	-0.001 (2)
N1	0.040 (4)	0.043 (4)	0.028 (3)	-0.009 (3)	0.003 (3)	-0.016 (3)
C6	0.025 (4)	0.027 (4)	0.029 (4)	-0.007 (3)	-0.001 (3)	-0.010 (3)
C8	0.024 (4)	0.026 (4)	0.028 (4)	-0.005 (3)	-0.003 (3)	-0.007 (3)
C5	0.048 (5)	0.046 (5)	0.042 (5)	-0.011 (4)	0.014 (4)	-0.028 (4)
C4	0.055 (6)	0.036 (5)	0.050 (6)	-0.008 (4)	0.013 (5)	-0.021 (5)
C3	0.038 (5)	0.032 (4)	0.047 (5)	0.000 (3)	0.001 (4)	-0.015 (4)
C2	0.018 (3)	0.025 (3)	0.029 (4)	-0.008 (3)	0.007 (3)	-0.006 (3)
C1	0.012 (3)	0.026 (3)	0.028 (4)	0.000 (3)	-0.003 (3)	-0.003 (3)
C7	0.018 (3)	0.020 (3)	0.020 (3)	-0.006 (2)	0.001 (3)	-0.007 (3)
O2W	0.069 (5)	0.078 (5)	0.041 (4)	-0.022 (4)	-0.020 (4)	-0.005 (4)
O2	0.019 (2)	0.027 (3)	0.027 (3)	-0.0043 (19)	-0.0037 (19)	0.002 (2)
O7	0.056 (4)	0.031 (3)	0.024 (3)	0.008 (2)	-0.018 (2)	-0.013 (2)
O1W	0.021 (3)	0.062 (4)	0.029 (3)	-0.012 (3)	-0.008 (2)	0.015 (3)

Geometric parameters (\AA , $^\circ$)

Ho1—O1	2.354 (5)	C6—C2	1.421 (10)
Ho1—O2 ⁱ	2.348 (5)	C8—O7	1.237 (9)
Ho1—O3	2.279 (5)	C8—C8 ⁱⁱⁱ	1.546 (15)
Ho1—O4	2.406 (4)	C5—C4	1.347 (13)
Ho1—O5 ⁱⁱ	2.364 (5)	C5—H5	0.9300
Ho1—O6	2.394 (5)	C4—C3	1.384 (11)
Ho1—O7 ⁱⁱⁱ	2.392 (5)	C4—H4	0.9300
Ho1—O1W	2.348 (5)	C3—C2	1.390 (10)
O4—C7	1.262 (8)	C3—H3	0.9300
O3—C6	1.279 (9)	C2—C1	1.487 (10)
O6—C8	1.263 (9)	C1—O2	1.271 (8)
O1—C1	1.249 (8)	C7—C7 ⁱⁱ	1.554 (12)
O5—C7	1.232 (8)	O2W—H3W	0.84 (7)
O5—Ho1 ⁱⁱ	2.364 (4)	O2W—H4W	0.86 (7)
N1—C5	1.358 (11)	O1W—H1W	0.84 (6)
N1—C6	1.373 (10)	O1W—H2W	0.85 (7)
N1—H1	0.8600		
O3—Ho1—O1W	90.4 (2)	C5—N1—C6	123.9 (7)
O3—Ho1—O2 ⁱ	90.37 (18)	C5—N1—H1	118.0
O1W—Ho1—O2 ⁱ	148.26 (17)	C6—N1—H1	118.0
O3—Ho1—O1	73.45 (17)	O3—C6—N1	116.8 (7)
O1W—Ho1—O1	75.15 (18)	O3—C6—C2	127.2 (7)
O2 ⁱ —Ho1—O1	74.69 (16)	N1—C6—C2	116.1 (7)
O3—Ho1—O5 ⁱⁱ	81.80 (17)	O7—C8—O6	127.4 (7)
O1W—Ho1—O5 ⁱⁱ	67.83 (16)	O7—C8—C8 ⁱⁱⁱ	117.7 (8)

O2 ⁱ —Ho1—O5 ⁱⁱ	143.49 (16)	O6—C8—C8 ⁱⁱⁱ	114.9 (8)
O1—Ho1—O5 ⁱⁱ	134.91 (16)	C4—C5—N1	120.6 (8)
O3—Ho1—O7 ⁱⁱⁱ	148.06 (18)	C4—C5—H5	119.7
O1W—Ho1—O7 ⁱⁱⁱ	105.7 (2)	N1—C5—H5	119.7
O2 ⁱ —Ho1—O7 ⁱⁱⁱ	89.77 (18)	C5—C4—C3	118.5 (8)
O1—Ho1—O7 ⁱⁱⁱ	136.83 (16)	C5—C4—H4	120.7
O5 ⁱⁱ —Ho1—O7 ⁱⁱⁱ	79.25 (18)	C3—C4—H4	120.7
O3—Ho1—O6	144.55 (17)	C4—C3—C2	121.7 (8)
O1W—Ho1—O6	75.0 (2)	C4—C3—H3	119.1
O2 ⁱ —Ho1—O6	86.35 (18)	C2—C3—H3	119.1
O1—Ho1—O6	71.64 (16)	C3—C2—C6	119.1 (7)
O5 ⁱⁱ —Ho1—O6	119.81 (18)	C3—C2—C1	120.1 (7)
O7 ⁱⁱⁱ —Ho1—O6	67.30 (17)	C6—C2—C1	120.8 (6)
O3—Ho1—O4	77.17 (17)	O1—C1—O2	122.8 (6)
O1W—Ho1—O4	135.57 (16)	O1—C1—C2	119.9 (6)
O2 ⁱ —Ho1—O4	75.25 (15)	O2—C1—C2	117.3 (6)
O1—Ho1—O4	137.32 (16)	O5—C7—O4	126.5 (6)
O5 ⁱⁱ —Ho1—O4	68.24 (14)	O5—C7—C7 ⁱⁱ	117.4 (7)
O7 ⁱⁱⁱ —Ho1—O4	72.00 (17)	O4—C7—C7 ⁱⁱ	116.0 (7)
O6—Ho1—O4	135.14 (16)	H3W—O2W—H4W	105 (9)
C7—O4—Ho1	118.2 (4)	C1—O2—Ho1 ⁱ	128.4 (4)
C6—O3—Ho1	136.0 (5)	C8—O7—Ho1 ⁱⁱⁱ	119.8 (5)
C8—O6—Ho1	120.3 (5)	Ho1—O1W—H1W	119 (6)
C1—O1—Ho1	136.8 (4)	Ho1—O1W—H2W	136 (6)
C7—O5—Ho1 ⁱⁱ	119.9 (4)	H1W—O1W—H2W	106 (3)

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $-x, -y+1, -z$; (iii) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O1W—H1W···O2 ^{iv}	0.84 (6)	2.02 (6)	2.737 (7)	142 (8)
O1W—H2W···O4 ^v	0.85 (7)	1.96 (4)	2.749 (7)	154 (8)
O2W—H3W···O6	0.84 (7)	2.36 (11)	2.882 (10)	121 (8)
O2W—H4W···O1	0.86 (7)	2.27 (3)	3.082 (8)	157 (9)
N1—H1···O2W ^{vi}	0.86	1.99	2.781 (10)	154

Symmetry codes: (iv) $-x+1, -y, -z+1$; (v) $x+1, y, z$; (vi) $x, y, z-1$.

supplementary materials

Fig. 1




Fig. 2

