

Poly[[bis[μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihydropyrido[2,3-*d*]-pyrimidine-6-carboxylato]zinc(II)] dihydrate]

Wen Xu,^a Da-Sheng Zhu,^a Xiao-Dan Song^b and Zhe An^{b*}

^aSecond Hospital, Jilin University, Changchun 130041, People's Republic of China, and ^bSchool of Pharmaceutical Science, Harbin Medical University, Harbin, 150086, People's Republic of China

Correspondence e-mail: anzhe6409@sina.com


Received 22 August 2009; accepted 12 September 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; H-atom completeness 89%; disorder in solvent or counterion; R factor = 0.057; wR factor = 0.210; data-to-parameter ratio = 16.2.

The title compound, $\{[\text{Zn}(\text{C}_{14}\text{H}_{16}\text{N}_5\text{O}_3)_2]\cdot 2\text{H}_2\text{O}\}_n$ or $[\text{Zn}(\text{ppa})_2]\cdot 2\text{H}_2\text{O}\}_n$, where ppa = 8-ethyl-5,8-dihydro-5-oxo-2-(1-piperazinyl)-pyrido(2,3-*d*)-pyrimidine-6-carboxylate, was synthesized under hydrothermal conditions. The Zn^{II} atom (site symmetry $\bar{1}$) exhibits a distorted *trans*- ZnN_2O_4 octahedral geometry defined by two monodentate *N*-bonded and two bidentate *O,O*-bonded ppa monoanions. The extended two-dimensional structure arising from this connectivity is a square grid and the disordered uncoordinated water molecules occupy cavities within the grid. An $\text{N}-\text{H}\cdots\text{O}$ hydrogen bond occurs.

Related literature

For manganese complexes of the ppa anion, see: Huang *et al.* (2008). For background to the medicinal uses of pipemicid acid, see: Mizuki *et al.* (1996).



Experimental

Crystal data

$[\text{Zn}(\text{C}_{14}\text{H}_{16}\text{N}_5\text{O}_3)_2]\cdot 2\text{H}_2\text{O}$	$V = 1616.9 (6)\text{ \AA}^3$
$M_r = 704.05$	$Z = 2$
Monoclinic, $P2_1/c$	$\text{Mo K}\alpha$ radiation
$a = 6.1146 (12)\text{ \AA}$	$\mu = 0.82\text{ mm}^{-1}$
$b = 21.424 (4)\text{ \AA}$	$T = 295\text{ K}$
$c = 12.577 (3)\text{ \AA}$	$0.36 \times 0.28 \times 0.18\text{ mm}$
$\beta = 101.10 (3)^\circ$	

Data collection

Rigaku R-AXIS RAPID diffractometer	15697 measured reflections
Absorption correction: multi-scan (<i>CrystalStructure</i> ; Rigaku/MSC, 2002)	3684 independent reflections
	2570 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.045$
	$T_{\min} = 0.756$, $T_{\max} = 0.866$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.210$	$\Delta\rho_{\max} = 0.83\text{ e \AA}^{-3}$
$S = 1.06$	$\Delta\rho_{\min} = -0.83\text{ e \AA}^{-3}$
3684 reflections	
228 parameters	
1 restraint	

Table 1
Selected bond lengths (\AA).

$\text{Zn1}-\text{O1}$	2.031 (3)	$\text{Zn1}-\text{N}5^{\dagger}$	2.275 (3)
$\text{Zn1}-\text{O3}$	2.107 (2)		

Symmetry code: (i) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$H\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N}5-\text{H}5\text{N}\cdots\text{O}2^{\text{ii}}$	0.91 (5)	2.28 (5)	3.168 (5)	166 (4)

Symmetry code: (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

The authors thank the Innovation Science Foundation of Harbin Medical University for financial support (grant No. 060041).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5062).

References

- Huang, J., Hu, W.-P. & An, Z. (2008). *Acta Cryst. E64*, m547.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Mizuki, Y., Fujiwara, I. & Yamaguchi, T. (1996). *J. Antimicrob. Chemother.* **37**, Suppl. A, 41–45.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m1223 [doi:10.1107/S1600536809036939]

Poly[[bis[μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihdropyrido[2,3-*d*]pyrimidine-6-carboxylato]zinc(II)] dihydrate]

W. Xu, D.-S. Zhu, X.-D. Song and Z. An

Comment

Pipemidic acid (Hppa, C₁₄H₁₆N₅O₃, 8-Ethyl-5,8-dihydro-5-oxo-2-(1-piperazinyl)-pyrido(2,3-*d*)-pyrimidine-6-carboxylic acid) is member of a class of quinolones used to treat infections (Mizuki *et al.*, 1996). The manganese complex of the ppa anion has been reported (Huang *et al.*, 2008); the title zinc(II) complex is reported here (Fig. 1).

The zinc(II) atom is coordinated by four oxygen atoms and two N atoms from four ppa ligands (two monodentate-N and two O,O-bidentate) to form a square grid (Fig. 2). The disordered, uncoordinated, water molecules occupy the cavities.

Experimental

A mixture of Zn(CH₃COO)₂.2H₂O (0.055 g, 0.25 mmol), Hppa (0.15 g, 0.5 mmol), sodium hydroxide (0.04 g, 1 mmol) and water (12 ml) was stirred for 30 min in air. The mixture was then transferred to a 23 ml Teflon-lined hydrothermal bomb. The bomb was kept at 433 K for 72 h under autogenous pressure. Upon cooling, colourless prisms of (I) were obtained from the reaction mixture.

Refinement

The carbon-bound H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with U_{iso}(H) = 1.2U_{eq}(C). The H on Nitrogen atoms were located in a difference Fourier map, and were refined with a distance restraint of N—H = 0.86 (1) Å and with U_{iso}(H) = 1.2U_{eq}(N).

The water H atoms could not be placed due to the disorder of the O atoms.

Figures




Fig. 1. The asymmetric unit of (I) extended to show the zinc coordination sphere showing the showing 50% displacement ellipsoids (water molecule O atoms have been omitted for clarity).




Fig. 2. A view of part of a two-dimensional polymeric sheet in (I) showing the square-grid connectivity (H atoms and water molecule O atoms omitted for clarity).

supplementary materials

Poly[[bis[μ_2 -8-ethyl-5-oxo-2-(piperazin-1-yl)-5,8-dihydropyrido[2,3-*d*]pyrimidine-6-carboxylato]zinc(II)] dihydrate]

Crystal data

[Zn(C ₁₄ H ₁₆ N ₅ O ₃) ₂]·2H ₂ O	Z = 2
M _r = 704.05	F ₀₀₀ = 728
Monoclinic, P2 ₁ /c	D _x = 1.442 Mg m ⁻³
Hall symbol: -P 2ybc	Mo K α radiation, λ = 0.71073 Å
a = 6.1146 (12) Å	μ = 0.82 mm ⁻¹
b = 21.424 (4) Å	T = 295 K
c = 12.577 (3) Å	Prism, colorless
β = 101.10 (3) $^\circ$	0.36 × 0.28 × 0.18 mm
V = 1616.9 (6) Å ³	

Data collection

Rigaku R-AXIS RAPID diffractometer	3684 independent reflections
Radiation source: fine-focus sealed tube	2570 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.045$
Detector resolution: 10.000 pixels mm ⁻¹	$\theta_{\text{max}} = 27.5^\circ$
T = 295 K	$\theta_{\text{min}} = 3.3^\circ$
ω scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (CrystalStructure; Rigaku/MSC, 2002)	$k = -27 \rightarrow 25$
$T_{\text{min}} = 0.756$, $T_{\text{max}} = 0.866$	$l = -16 \rightarrow 16$
15697 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.057$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.210$	$w = 1/[\sigma^2(F_o^2) + (0.1254P)^2 + 1.7801P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} < 0.001$
3684 reflections	$\Delta\rho_{\text{max}} = 0.83 \text{ e \AA}^{-3}$
228 parameters	$\Delta\rho_{\text{min}} = -0.83 \text{ e \AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1W	-0.045 (3)	-0.0632 (10)	-0.0746 (11)	0.187 (8)	0.50
O2W	0.340 (5)	0.0205 (10)	-0.0364 (12)	0.251 (14)	0.50
Zn1	0.5000	0.0000	0.5000	0.0265 (2)	
O1	0.6981 (4)	-0.00325 (10)	0.3877 (2)	0.0271 (6)	
O2	0.8573 (7)	0.01818 (18)	0.2500 (3)	0.0616 (11)	
O3	0.3495 (5)	0.07935 (11)	0.4179 (2)	0.0317 (6)	
N1	0.4916 (7)	0.17173 (17)	0.1532 (3)	0.0481 (10)	
N2	0.2252 (6)	0.24690 (15)	0.1677 (3)	0.0386 (8)	
N3	-0.0127 (6)	0.23572 (16)	0.2988 (3)	0.0436 (9)	
N4	-0.0227 (6)	0.32384 (15)	0.1907 (3)	0.0349 (8)	
N5	-0.2450 (5)	0.43908 (14)	0.1084 (2)	0.0273 (7)	
H5N	-0.154 (8)	0.466 (2)	0.152 (4)	0.065 (16)*	
C1	0.7147 (7)	0.02891 (17)	0.3064 (3)	0.0316 (8)	
C2	0.5658 (7)	0.08450 (17)	0.2772 (3)	0.0317 (8)	
C3	0.3947 (6)	0.10453 (16)	0.3346 (3)	0.0271 (7)	
C4	0.2744 (7)	0.15974 (16)	0.2910 (3)	0.0303 (8)	
C5	0.0937 (8)	0.18359 (19)	0.3318 (3)	0.0398 (10)	
H5	0.0454	0.1610	0.3861	0.048*	
C6	0.0671 (7)	0.26762 (18)	0.2197 (3)	0.0327 (8)	
C7	0.3246 (7)	0.19340 (18)	0.2034 (3)	0.0360 (9)	
C8	0.6010 (8)	0.1189 (2)	0.1902 (4)	0.0457 (11)	
H8	0.7096	0.1047	0.1535	0.055*	
C9	0.5540 (11)	0.2051 (3)	0.0585 (5)	0.0665 (16)	
H9B	0.5306	0.2496	0.0653	0.080*	
H9A	0.7107	0.1983	0.0582	0.080*	
C10	0.4247 (16)	0.1834 (5)	-0.0401 (7)	0.116 (3)	
H10C	0.4566	0.1401	-0.0496	0.174*	
H10B	0.4605	0.2071	-0.0993	0.174*	
H10A	0.2693	0.1883	-0.0385	0.174*	
C11	-0.1608 (8)	0.3572 (2)	0.2553 (4)	0.0475 (11)	
H11B	-0.0673	0.3839	0.3077	0.057*	
H11A	-0.2346	0.3275	0.2947	0.057*	
C12	-0.3356 (7)	0.3970 (2)	0.1813 (4)	0.0398 (10)	

supplementary materials

H12B	-0.4429	0.3694	0.1379	0.048*
H12A	-0.4150	0.4218	0.2262	0.048*
C13	-0.1090 (6)	0.40176 (17)	0.0469 (3)	0.0317 (8)
H13A	-0.0405	0.4295	0.0018	0.038*
H13B	-0.2057	0.3734	-0.0006	0.038*
C14	0.0708 (7)	0.36460 (18)	0.1185 (4)	0.0369 (9)
H14B	0.1500	0.3397	0.0737	0.044*
H14A	0.1768	0.3929	0.1610	0.044*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1W	0.209 (18)	0.28 (2)	0.094 (9)	-0.004 (16)	0.091 (11)	0.013 (12)
O2W	0.47 (4)	0.202 (17)	0.077 (9)	0.11 (2)	0.049 (16)	-0.057 (12)
Zn1	0.0317 (4)	0.0194 (3)	0.0294 (3)	0.0006 (2)	0.0079 (2)	0.0028 (2)
O1	0.0276 (13)	0.0218 (12)	0.0333 (14)	0.0019 (9)	0.0089 (10)	0.0029 (10)
O2	0.073 (3)	0.062 (2)	0.060 (2)	0.040 (2)	0.0387 (19)	0.0275 (18)
O3	0.0378 (15)	0.0223 (12)	0.0368 (14)	0.0092 (10)	0.0116 (11)	0.0121 (11)
N1	0.065 (3)	0.0398 (19)	0.046 (2)	0.0223 (18)	0.0265 (19)	0.0195 (17)
N2	0.051 (2)	0.0282 (16)	0.0400 (18)	0.0135 (15)	0.0160 (16)	0.0124 (14)
N3	0.044 (2)	0.0361 (18)	0.055 (2)	0.0166 (16)	0.0219 (17)	0.0231 (17)
N4	0.0390 (19)	0.0283 (16)	0.0415 (18)	0.0096 (14)	0.0177 (15)	0.0119 (14)
N5	0.0265 (16)	0.0219 (14)	0.0324 (15)	0.0064 (12)	0.0028 (12)	0.0034 (13)
C1	0.037 (2)	0.0274 (18)	0.0316 (18)	0.0046 (15)	0.0112 (16)	-0.0002 (16)
C2	0.038 (2)	0.0238 (17)	0.0354 (19)	0.0070 (15)	0.0110 (16)	0.0035 (15)
C3	0.0283 (18)	0.0219 (16)	0.0302 (17)	0.0007 (13)	0.0032 (14)	0.0013 (14)
C4	0.036 (2)	0.0242 (17)	0.0318 (18)	0.0021 (15)	0.0087 (15)	0.0055 (15)
C5	0.048 (3)	0.033 (2)	0.043 (2)	0.0088 (18)	0.0188 (19)	0.0174 (18)
C6	0.033 (2)	0.0273 (18)	0.0380 (19)	0.0046 (15)	0.0078 (16)	0.0077 (16)
C7	0.044 (2)	0.0323 (19)	0.0347 (19)	0.0092 (17)	0.0137 (17)	0.0067 (17)
C8	0.060 (3)	0.037 (2)	0.046 (2)	0.018 (2)	0.026 (2)	0.0107 (19)
C9	0.083 (4)	0.063 (3)	0.062 (3)	0.030 (3)	0.038 (3)	0.024 (3)
C10	0.117 (7)	0.144 (9)	0.090 (5)	0.031 (6)	0.028 (5)	0.024 (6)
C11	0.055 (3)	0.045 (2)	0.047 (2)	0.025 (2)	0.021 (2)	0.020 (2)
C12	0.040 (2)	0.036 (2)	0.047 (2)	0.0157 (17)	0.0179 (18)	0.0161 (19)
C13	0.035 (2)	0.0247 (17)	0.0366 (19)	0.0101 (15)	0.0094 (16)	0.0081 (15)
C14	0.033 (2)	0.0293 (18)	0.051 (2)	0.0043 (16)	0.0147 (17)	0.0119 (18)

Geometric parameters (\AA , °)

Zn1—O1	2.031 (3)	C2—C8	1.370 (6)
Zn1—O1 ⁱ	2.031 (3)	C2—C3	1.446 (5)
Zn1—O3 ⁱ	2.107 (2)	C3—C4	1.444 (5)
Zn1—O3	2.107 (2)	C4—C7	1.399 (5)
Zn1—N5 ⁱⁱ	2.275 (3)	C4—C5	1.401 (6)
Zn1—N5 ⁱⁱⁱ	2.275 (3)	C5—H5	0.9300
O1—C1	1.253 (5)	C8—H8	0.9300
O2—C1	1.247 (5)	C9—C10	1.415 (11)

O3—C3	1.256 (4)	C9—H9B	0.9700
N1—C8	1.351 (5)	C9—H9A	0.9700
N1—C7	1.381 (5)	C10—H10C	0.9600
N1—C9	1.500 (6)	C10—H10B	0.9600
N2—C7	1.334 (5)	C10—H10A	0.9600
N2—C6	1.343 (5)	C11—C12	1.534 (5)
N3—C5	1.319 (5)	C11—H11B	0.9700
N3—C6	1.373 (5)	C11—H11A	0.9700
N4—C6	1.345 (5)	C12—H12B	0.9700
N4—C14	1.454 (5)	C12—H12A	0.9700
N4—C11	1.466 (5)	C13—C14	1.508 (5)
N5—C12	1.468 (5)	C13—H13A	0.9700
N5—C13	1.475 (5)	C13—H13B	0.9700
N5—Zn1 ^{iv}	2.275 (3)	C14—H14B	0.9700
N5—H5N	0.900 (10)	C14—H14A	0.9700
C1—C2	1.501 (5)		
O1—Zn1—O1 ⁱ	180.0	N2—C6—N4	117.3 (3)
O1—Zn1—O3 ⁱ	92.90 (10)	N2—C6—N3	125.3 (4)
O1 ⁱ —Zn1—O3 ⁱ	87.10 (10)	N4—C6—N3	117.3 (4)
O1—Zn1—O3	87.10 (10)	N2—C7—N1	117.6 (3)
O1 ⁱ —Zn1—O3	92.90 (10)	N2—C7—C4	123.6 (4)
O3 ⁱ —Zn1—O3	180.0	N1—C7—C4	118.7 (3)
O1—Zn1—N5 ⁱⁱ	89.74 (11)	N1—C8—C2	125.6 (4)
O1 ⁱ —Zn1—N5 ⁱⁱ	90.26 (11)	N1—C8—H8	117.2
O3 ⁱ —Zn1—N5 ⁱⁱ	90.86 (11)	C2—C8—H8	117.2
O3—Zn1—N5 ⁱⁱ	89.14 (11)	C10—C9—N1	110.8 (7)
O1—Zn1—N5 ⁱⁱⁱ	90.26 (11)	C10—C9—H9B	109.5
O1 ⁱ —Zn1—N5 ⁱⁱⁱ	89.74 (11)	N1—C9—H9B	109.5
O3 ⁱ —Zn1—N5 ⁱⁱⁱ	89.14 (11)	C10—C9—H9A	109.5
O3—Zn1—N5 ⁱⁱⁱ	90.86 (11)	N1—C9—H9A	109.5
N5 ⁱⁱ —Zn1—N5 ⁱⁱⁱ	180.0	H9B—C9—H9A	108.1
C1—O1—Zn1	134.5 (2)	C9—C10—H10C	109.5
C3—O3—Zn1	127.6 (2)	C9—C10—H10B	109.5
C8—N1—C7	119.0 (3)	H10C—C10—H10B	109.5
C8—N1—C9	119.2 (4)	C9—C10—H10A	109.5
C7—N1—C9	121.8 (4)	H10C—C10—H10A	109.5
C7—N2—C6	116.3 (3)	H10B—C10—H10A	109.5
C5—N3—C6	115.3 (4)	N4—C11—C12	110.0 (3)
C6—N4—C14	121.2 (3)	N4—C11—H11B	109.7
C6—N4—C11	122.4 (3)	C12—C11—H11B	109.7
C14—N4—C11	113.0 (3)	N4—C11—H11A	109.7
C12—N5—C13	108.3 (3)	C12—C11—H11A	109.7
C12—N5—Zn1 ^{iv}	115.4 (2)	H11B—C11—H11A	108.2
C13—N5—Zn1 ^{iv}	112.8 (2)	N5—C12—C11	114.7 (3)
C12—N5—H5N	106 (4)	N5—C12—H12B	108.6

supplementary materials

C13—N5—H5N	108 (4)	C11—C12—H12B	108.6
Zn1 ^{iv} —N5—H5N	106 (4)	N5—C12—H12A	108.6
O2—C1—O1	122.6 (4)	C11—C12—H12A	108.6
O2—C1—C2	117.7 (3)	H12B—C12—H12A	107.6
O1—C1—C2	119.7 (3)	N5—C13—C14	113.1 (3)
C8—C2—C3	118.6 (3)	N5—C13—H13A	109.0
C8—C2—C1	116.2 (3)	C14—C13—H13A	109.0
C3—C2—C1	125.2 (3)	N5—C13—H13B	109.0
O3—C3—C4	119.4 (3)	C14—C13—H13B	109.0
O3—C3—C2	125.8 (3)	H13A—C13—H13B	107.8
C4—C3—C2	114.7 (3)	N4—C14—C13	111.2 (3)
C7—C4—C5	114.1 (3)	N4—C14—H14B	109.4
C7—C4—C3	123.2 (4)	C13—C14—H14B	109.4
C5—C4—C3	122.6 (3)	N4—C14—H14A	109.4
N3—C5—C4	124.7 (4)	C13—C14—H14A	109.4
N3—C5—H5	117.6	H14B—C14—H14A	108.0
C4—C5—H5	117.6		
O1 ⁱ —Zn1—O1—C1	-50 (2)	C11—N4—C6—N2	-167.0 (4)
O3 ⁱ —Zn1—O1—C1	179.5 (4)	C14—N4—C6—N3	171.7 (4)
O3—Zn1—O1—C1	-0.5 (4)	C11—N4—C6—N3	14.0 (6)
N5 ⁱⁱ —Zn1—O1—C1	88.7 (4)	C5—N3—C6—N2	7.0 (7)
N5 ⁱⁱⁱ —Zn1—O1—C1	-91.3 (4)	C5—N3—C6—N4	-174.0 (4)
O1—Zn1—O3—C3	0.4 (3)	C6—N2—C7—N1	-178.5 (4)
O1 ⁱ —Zn1—O3—C3	-179.6 (3)	C6—N2—C7—C4	-1.0 (6)
O3 ⁱ —Zn1—O3—C3	65 (100)	C8—N1—C7—N2	177.5 (4)
N5 ⁱⁱ —Zn1—O3—C3	-89.4 (3)	C9—N1—C7—N2	-2.7 (7)
N5 ⁱⁱⁱ —Zn1—O3—C3	90.6 (3)	C8—N1—C7—C4	-0.1 (7)
Zn1—O1—C1—O2	178.2 (3)	C9—N1—C7—C4	179.7 (5)
Zn1—O1—C1—C2	1.0 (6)	C5—C4—C7—N2	6.2 (6)
O2—C1—C2—C8	-0.1 (6)	C3—C4—C7—N2	-174.9 (4)
O1—C1—C2—C8	177.1 (4)	C5—C4—C7—N1	-176.3 (4)
O2—C1—C2—C3	-178.8 (4)	C3—C4—C7—N1	2.5 (6)
O1—C1—C2—C3	-1.6 (6)	C7—N1—C8—C2	-1.9 (8)
Zn1—O3—C3—C4	-179.0 (2)	C9—N1—C8—C2	178.3 (5)
Zn1—O3—C3—C2	-1.1 (5)	C3—C2—C8—N1	1.4 (7)
C8—C2—C3—O3	-177.0 (4)	C1—C2—C8—N1	-177.4 (4)
C1—C2—C3—O3	1.7 (6)	C8—N1—C9—C10	90.7 (7)
C8—C2—C3—C4	1.0 (6)	C7—N1—C9—C10	-89.1 (7)
C1—C2—C3—C4	179.7 (4)	C6—N4—C11—C12	-148.9 (4)
O3—C3—C4—C7	175.2 (4)	C14—N4—C11—C12	51.8 (5)
C2—C3—C4—C7	-2.9 (6)	C13—N5—C12—C11	53.7 (5)
O3—C3—C4—C5	-6.0 (6)	Zn1 ^{iv} —N5—C12—C11	-178.8 (3)
C2—C3—C4—C5	175.9 (4)	N4—C11—C12—N5	-52.7 (5)
C6—N3—C5—C4	-0.8 (7)	C12—N5—C13—C14	-55.0 (4)
C7—C4—C5—N3	-5.2 (7)	Zn1 ^{iv} —N5—C13—C14	176.0 (2)
C3—C4—C5—N3	175.9 (4)	C6—N4—C14—C13	145.9 (4)

supplementary materials

C7—N2—C6—N4	174.8 (4)	C11—N4—C14—C13	−54.5 (5)
C7—N2—C6—N3	−6.2 (7)	N5—C13—C14—N4	56.5 (5)
C14—N4—C6—N2	−9.3 (6)		

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, y-1/2, -z+1/2$; (iii) $x+1, -y+1/2, z+1/2$; (iv) $-x, y+1/2, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N5—H5N \cdots O2 ^v	0.91 (5)	2.28 (5)	3.168 (5)	166 (4)

Symmetry codes: (v) $-x+1, y+1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

