

Bis[4-oxido-2-oxo-2,3-dihydropyrimidin-1-ium-5-carboxylato(1.5-)–κ²O⁴,O⁵]-bis(1,10-phenanthroline-κ²N,N')-dysprosium(III) dihydrate

Xianlin Liu, Huihui Xing, Jing Miao, Maomao Jia and Zilu Chen*

College of Chemistry and Chemical Engineering, Guangxi Normal University, Yucai Road 15, Guilin 541004, People's Republic of China
Correspondence e-mail: chenzilucl@yahoo.co.uk


Received 25 August 2009; accepted 11 September 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.007\text{ \AA}$; R factor = 0.030; wR factor = 0.068; data-to-parameter ratio = 11.8.

In the title compound, $[\text{Dy}(\text{C}_5\text{H}_{2.5}\text{N}_2\text{O}_4)_2(\text{C}_{12}\text{H}_8\text{N}_2)_2]\cdot 2\text{H}_2\text{O}$, the Dy^{III} ion is located on a twofold rotation axis and is coordinated in a square-antiprismatic geometry by two chelating 1,10-phenanthroline molecules as well as two 4-oxido-2-oxo-2,3-dihydropyrimidin-1-ium-5-carboxylato(1.5-) anions. N—H···O and O—H···O hydrogen bonds help to stabilize the crystal structure. The H atom involved in an N—H···N hydrogen bond is disordered around a twofold rotation axis.

Related literature

For isostructural lanthanide complexes with 2,4-dioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid, see: Sun & Jin (2004a,b); Xing *et al.* (2008a); Xiong *et al.* (2008a,b). For other metal complexes of 2,4-dioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid, see: Hueso-Ureña *et al.* (1993, 1996); Baran *et al.* (1996); Luo *et al.* (2002); Maistralis *et al.* (1991, 1992); Xing *et al.* (2008b). For the role played by hydrogen bonds in stabilizing structures, see: Chen *et al.* (2006); Arora *et al.* (2009); Jagan & Sivakumar (2009).



Experimental

Crystal data

$[\text{Dy}(\text{C}_5\text{H}_{2.5}\text{N}_2\text{O}_4)_2(\text{C}_{12}\text{H}_8\text{N}_2)_2]\cdot 2\text{H}_2\text{O}$	$\beta = 100.613(2)^\circ$
$M_r = 868.12$	$V = 3218.3(7)\text{ \AA}^3$
Monoclinic, $C2/c$	$Z = 4$
$a = 17.143(2)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 14.4470(17)\text{ \AA}$	$\mu = 2.40\text{ mm}^{-1}$
$c = 13.2211(16)\text{ \AA}$	$T = 295\text{ K}$
	$0.10 \times 0.03 \times 0.02\text{ mm}$

Data collection

Bruker APEXII CCD area-detector diffractometer	11116 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1998)	2842 independent reflections
$T_{\min} = 0.796$, $T_{\max} = 0.954$	2619 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$	240 parameters
$wR(F^2) = 0.068$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\max} = 0.93\text{ e \AA}^{-3}$
2842 reflections	$\Delta\rho_{\min} = -1.22\text{ e \AA}^{-3}$

Table 1
Selected geometric parameters (\AA , $^\circ$).

Dy—O2	2.256 (3)	Dy—N2	2.554 (3)
Dy—O1	2.313 (2)	Dy—N1	2.576 (3)

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H3···N3 ⁱⁱ	0.86	1.80	2.659 (7)	174
N4—H4···O3 ⁱⁱⁱ	0.86	2.01	2.867 (4)	178
N4—H4···O2 ⁱⁱⁱ	0.86	2.62	3.183 (4)	124
O5—H51···O3 ^{iv}	0.85	2.15	2.993 (5)	175
O5—H52···O4 ^v	0.85	2.15	2.960 (5)	160

Symmetry codes: (ii) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 2$; (iii) $x, -y + 1, z + \frac{1}{2}$; (iv) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$; (v) $x, y, z - 1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors are grateful for financial support from the Guangxi Natural Science Foundation (grant No. 0991008) and the Scientific Research Foundation for Returned Overseas Chinese Scholars, State Education Ministry, China (grant No. [2006]331).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PB2007).

References

- Arora, K. K., Talwelkar, M. S. & Pedireddi, V. R. (2009). *New J. Chem.* **33**, 57–63.
- Baran, E., Mercaser, R. C., Hueso-Ureña, F., Moreno-Carretero, M. N., Quiros-Olozabal, M. & Salas-Peregrin, J. M. (1996). *Polyhedron*, **15**, 1717–1724.

- Bruker (1998). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, Z.-L., Zhang, Y.-Z. & Liang, F.-P. (2006). *Acta Cryst. E* **62**, m1296–m1297.
- Hueso-Ureña, F., Moreno-Carretero, M. N., Romero-Molina, M. A., Salas-Peregrin, J. M., Sanchez-Sanchez, M. P., Alvarez de Cienfuegos-Lopez, G. & Faure, R. (1993). *J. Inorg. Biochem.* **51**, 613–632.
- Hueso-Ureña, F., Moreno-Carretero, M. N., Quirós-Olozábal, M., Salas-Peregrín, J. M., Faure, R. & Cienfuegos-López, G. A. (1996). *Inorg. Chim. Acta*, **241**, 61–69.
- Jagan, R. & Sivakumar, K. (2009). *Acta Cryst. C* **65**, o414–o418.
- Luo, J.-H., Hong, M.-C., Zhao, Y.-J., Cao, R. & Weng, J.-B. (2002). *Chin. J. Struct. Chem.* **21**, 392–395.
- Maistralis, G., Katsaros, N., Mentzas, D. & Terzis, A. (1991). *Acta Cryst. C* **47**, 740–743.
- Maistralis, G., Katsaros, N., Perlepes, S. P. & Kovala-Demertz, D. (1992). *J. Inorg. Biochem.* **45**, 1–12.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sun, C.-Y. & Jin, L.-P. (2004a). *Polyhedron*, **23**, 2085–2093.
- Sun, C.-Y. & Jin, L.-P. (2004b). *Polyhedron*, **23**, 2227–2233.
- Xing, H.-H., Chen, Z.-L. & Ng, S. W. (2008a). *Acta Cryst. E* **64**, m418.
- Xing, H.-H., Chen, Z.-L. & Ng, S. W. (2008b). *Acta Cryst. E* **64**, m419–m420.
- Xiong, W., Xing, H., Su, Y. & Chen, Z. (2008a). *Acta Cryst. E* **64**, m1166–m1167.
- Xiong, W., Xing, H., Su, Y. & Chen, Z. (2008b). *Acta Cryst. E* **64**, m1168–m1169.

supplementary materials

Acta Cryst. (2009). E65, m1224-m1225 [doi:10.1107/S1600536809036848]

Bis[4-oxido-2-oxo-2,3-dihydropyrimidin-1-ium-5-carboxylato(1.5-)- κ^2O^4,O^5]bis(1,10-phenanthroline- κ^2N,N')dysprosium(III) dihydrate

X. Liu, H. Xing, J. Miao, M. Jia and Z. Chen

Comment

2,4-Dioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid has been extensively used in the preparation of robust networks or some porous coordination polymers because of its versatile coordination modes. For further investigation of its coordination behavior to lanthanide(III) ions, we report here a new Dy^{III} complex of 2,4-dioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid, which turned out to be isostructural with the analogues Eu^{III}, Tb^{III}, Yb^{III} (2004a,b), Er^{III} (Xing *et al.*, 2008a) and Y^{III}, Gd^{III} (Xiong *et al.*, 2008a,b) complexes.

In the title complex, (I), the Dy^{III} ion is located on a twofold rotation axis and coordinated in square-antiprismatic geometry completed by four N atoms from two chelating phenanthroline molecules, four O atoms from two chelating 4-oxido-2-oxo-2,3-dihydropyrimidin-1-ium-5-carboxylate(1.5-) anions. The distances of Dy—O are 2.256 (3) and 2.313 (2) Å, and those of Dy—N are 2.554 (3) and 2.576 (3) Å. The average distance of Dy—O is 2.285 Å, which is much shorter than that of Dy—N (2.565 Å). The dihedral angle between two phen molecules is 37.726°. As shown in Table 2 and Fig. 2, each coordination unit [Dy(C₅H₃N₂O₄)(C₅H₂N₂O₄)(C₁₂H₈N₂)₂]⁺ interacts with another four *via* intermolecular N—H···O and N—H···N hydrogen bonds. Each lattice water molecule links two coordination units mentioned above *via* O—H···O hydrogen bonds. As a result, these hydrogen bonds link the title complex units and lattice water molecules into a three-dimensional supramolecular network as shown in the packing diagram (Fig. 2). The H atom involved in an N—H···N hydrogen bond is disordered around a twofold rotation axis with half occupancy.

Experimental

A mixture of 2,4-dioxo-1,2,3,4-tetrahydropyrimidine-5-carboxylic acid (0.0312 g, 0.2 mmol), DyCl₃·6H₂O (0.0754 g, 0.2 mmol), Phen·H₂O (0.0396 g, 0.2 mmol), NaOH (0.00800 g, 0.2 mmol), water (15 ml) was sealed in a 25 ml Teflon-lined stainless-steel reactor and heated to 383 K for 120 h, it was then cooled over 48 h, light yellow crystals were isolated in 80% yield. Elemental analysis for C₃₄H₂₅DyN₈O₁₀, calculated: C 47.04, H 2.90, N 12.91%; found: C 47.49, H 2.96, N 13.24%.

Refinement

H atoms of the water molecule were located in a difference Fourier map and allowed to ride on their parent atoms [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$]. Other H atoms were placed at calculated positions (C—H = 0.93 Å and N—H = 0.86 Å) and were included in the refinement in the riding model approximation, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C}, \text{N})$. The pyrimidine hydrogen atom H3 is shared by two N—H groups and thus has an occupancy factor of 0.5.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom-numbering scheme and 30% displacement ellipsoids.




Fig. 2. A view of the packing diagram of the title compound viewed down the *c* axis. H atoms not involving in hydrogen bonds are omitted for clarity.

Bis[4-oxido-2-oxo-2,3-dihydropyrimidin-1-i um-5-carboxylato(1.5)- κ^2O^5,O^6]bis(1,10-phenanthroline- κ^2N,N')dysprosium(III) dihydrate

Crystal data

[Dy(C ₅ H _{2.5} N ₂ O ₄) ₂ (C ₁₂ H ₈ N ₂) ₂] <cdot>2H₂O</cdot>	$F_{000} = 1724$
$M_r = 868.12$	$D_x = 1.792 \text{ Mg m}^{-3}$
Monoclinic, <i>C</i> 2/ <i>c</i>	Mo <i>K</i> α radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -C 2yc	Cell parameters from 4430 reflections
$a = 17.143 (2) \text{ \AA}$	$\theta = 2.3\text{--}26.9^\circ$
$b = 14.4470 (17) \text{ \AA}$	$\mu = 2.40 \text{ mm}^{-1}$
$c = 13.2211 (16) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 100.613 (2)^\circ$	Prism, yellow
$V = 3218.3 (7) \text{ \AA}^3$	$0.10 \times 0.03 \times 0.02 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	2842 independent reflections
Radiation source: fine-focus sealed tube	2619 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.037$
$T = 295 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 1998)	$h = -20 \rightarrow 20$
$T_{\text{min}} = 0.796$, $T_{\text{max}} = 0.954$	$k = -17 \rightarrow 17$
11116 measured reflections	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H-atom parameters constrained
$wR(F^2) = 0.068$	$w = 1/[\sigma^2(F_o^2) + (0.0366P)^2 + 4.1092P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\max} = 0.001$
2842 reflections	$\Delta\rho_{\max} = 0.93 \text{ e \AA}^{-3}$
240 parameters	$\Delta\rho_{\min} = -1.22 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Dy	0.0000	0.612148 (16)	0.7500	0.02165 (10)	
N1	0.03494 (19)	0.6959 (2)	0.5922 (2)	0.0299 (7)	
N2	0.09455 (19)	0.7496 (2)	0.7881 (3)	0.0318 (8)	
N3	0.2176 (2)	0.3312 (2)	0.9692 (2)	0.0379 (9)	
H3	0.2412	0.2796	0.9869	0.046*	0.50
N4	0.16685 (19)	0.4740 (2)	1.0061 (2)	0.0295 (8)	
H4	0.1617	0.5157	1.0509	0.035*	
O1	0.09579 (16)	0.56623 (18)	0.88770 (19)	0.0302 (6)	
O2	0.07048 (16)	0.50051 (18)	0.68831 (19)	0.0327 (6)	
O3	0.15320 (18)	0.38990 (19)	0.6597 (2)	0.0362 (7)	
O4	0.2306 (2)	0.38078 (19)	1.1355 (2)	0.0437 (8)	
C1	0.1236 (2)	0.4394 (3)	0.7192 (3)	0.0278 (9)	
C2	0.1504 (2)	0.4274 (3)	0.8320 (3)	0.0262 (8)	
C3	0.1915 (3)	0.3504 (3)	0.8700 (3)	0.0363 (10)	
H3A	0.2025	0.3072	0.8224	0.044*	
C4	0.2065 (3)	0.3940 (3)	1.0424 (3)	0.0326 (9)	
C5	0.1350 (2)	0.4930 (2)	0.9059 (3)	0.0240 (8)	
C6	0.1293 (2)	0.7739 (3)	0.8822 (3)	0.0362 (10)	

supplementary materials

H6	0.1307	0.7310	0.9350	0.043*
C7	0.1639 (3)	0.8606 (3)	0.9064 (4)	0.0479 (12)
H7	0.1875	0.8750	0.9736	0.057*
C8	0.1626 (3)	0.9232 (3)	0.8298 (4)	0.0507 (13)
H8	0.1847	0.9815	0.8448	0.061*
C9	0.1283 (3)	0.9012 (3)	0.7289 (4)	0.0416 (11)
C10	0.0959 (2)	0.8115 (3)	0.7112 (3)	0.0303 (9)
C11	0.0644 (2)	0.7832 (3)	0.6071 (3)	0.0315 (9)
C12	0.0658 (3)	0.8453 (3)	0.5253 (4)	0.0451 (11)
C13	0.0962 (3)	0.9367 (4)	0.5480 (4)	0.0622 (15)
H13	0.0954	0.9786	0.4945	0.075*
C14	0.1256 (3)	0.9631 (3)	0.6439 (5)	0.0610 (15)
H14	0.1449	1.0231	0.6559	0.073*
C15	0.0367 (3)	0.8133 (4)	0.4259 (4)	0.0577 (14)
H15	0.0352	0.8526	0.3699	0.069*
C16	0.0108 (3)	0.7253 (4)	0.4111 (3)	0.0514 (13)
H16	-0.0064	0.7029	0.3448	0.062*
C17	0.0099 (2)	0.6686 (3)	0.4955 (3)	0.0384 (10)
H17	-0.0091	0.6085	0.4839	0.046*
O5	0.3087 (3)	0.3123 (3)	0.3401 (3)	0.0926 (15)
H51	0.3201	0.2551	0.3440	0.139*
H52	0.2760	0.3253	0.2854	0.139*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Dy	0.02842 (15)	0.01487 (14)	0.01945 (14)	0.000	-0.00135 (10)	0.000
N1	0.0310 (18)	0.0264 (18)	0.0312 (18)	-0.0013 (14)	0.0023 (14)	-0.0005 (14)
N2	0.0311 (19)	0.0288 (19)	0.0350 (19)	-0.0034 (15)	0.0049 (15)	-0.0040 (15)
N3	0.060 (2)	0.0248 (18)	0.0264 (18)	0.0105 (17)	-0.0002 (16)	0.0024 (15)
N4	0.042 (2)	0.0245 (17)	0.0195 (16)	0.0066 (15)	0.0000 (14)	-0.0021 (13)
O1	0.0425 (16)	0.0218 (14)	0.0224 (13)	0.0096 (13)	-0.0044 (12)	-0.0024 (11)
O2	0.0439 (17)	0.0310 (15)	0.0209 (13)	0.0115 (13)	-0.0003 (12)	0.0015 (12)
O3	0.0519 (18)	0.0361 (16)	0.0211 (14)	0.0150 (14)	0.0079 (13)	-0.0015 (12)
O4	0.060 (2)	0.0342 (17)	0.0305 (16)	0.0059 (15)	-0.0095 (14)	0.0031 (13)
C1	0.036 (2)	0.021 (2)	0.025 (2)	-0.0020 (18)	0.0015 (17)	0.0014 (17)
C2	0.032 (2)	0.024 (2)	0.0217 (19)	0.0052 (17)	0.0024 (16)	0.0012 (16)
C3	0.053 (3)	0.025 (2)	0.029 (2)	0.010 (2)	0.0043 (19)	-0.0041 (18)
C4	0.039 (2)	0.028 (2)	0.028 (2)	-0.0024 (18)	-0.0028 (17)	0.0038 (17)
C5	0.029 (2)	0.021 (2)	0.0211 (19)	-0.0003 (16)	0.0017 (15)	0.0000 (15)
C6	0.038 (2)	0.033 (2)	0.035 (2)	-0.0083 (19)	0.0024 (19)	-0.0072 (19)
C7	0.043 (3)	0.048 (3)	0.052 (3)	-0.014 (2)	0.009 (2)	-0.024 (2)
C8	0.051 (3)	0.034 (3)	0.068 (3)	-0.016 (2)	0.013 (3)	-0.019 (3)
C9	0.042 (3)	0.024 (2)	0.061 (3)	-0.0076 (19)	0.014 (2)	-0.006 (2)
C10	0.025 (2)	0.026 (2)	0.040 (2)	-0.0013 (16)	0.0070 (17)	-0.0005 (18)
C11	0.029 (2)	0.029 (2)	0.037 (2)	0.0015 (17)	0.0071 (18)	0.0051 (18)
C12	0.044 (3)	0.041 (3)	0.052 (3)	-0.003 (2)	0.012 (2)	0.014 (2)
C13	0.077 (4)	0.044 (3)	0.066 (4)	-0.014 (3)	0.013 (3)	0.022 (3)

C14	0.070 (4)	0.028 (3)	0.088 (4)	-0.015 (2)	0.022 (3)	0.005 (3)
C15	0.067 (4)	0.066 (4)	0.038 (3)	-0.010 (3)	0.007 (2)	0.024 (3)
C16	0.056 (3)	0.070 (4)	0.028 (2)	-0.013 (3)	0.007 (2)	0.005 (2)
C17	0.037 (2)	0.042 (3)	0.036 (2)	-0.004 (2)	0.0056 (19)	-0.005 (2)
O5	0.142 (4)	0.047 (2)	0.070 (3)	0.003 (3)	-0.027 (3)	-0.010 (2)

Geometric parameters (\AA , $^{\circ}$)

Dy—O2 ⁱ	2.256 (3)	C2—C5	1.420 (5)
Dy—O2	2.256 (3)	C3—H3A	0.9300
Dy—O1 ⁱ	2.313 (2)	C6—C7	1.398 (6)
Dy—O1	2.313 (2)	C6—H6	0.9300
Dy—N2	2.554 (3)	C7—C8	1.355 (7)
Dy—N2 ⁱ	2.554 (3)	C7—H7	0.9300
Dy—N1 ⁱ	2.576 (3)	C8—C9	1.392 (7)
Dy—N1	2.576 (3)	C8—H8	0.9300
N1—C17	1.331 (5)	C9—C10	1.412 (5)
N1—C11	1.360 (5)	C9—C14	1.431 (7)
N2—C6	1.324 (5)	C10—C11	1.442 (6)
N2—C10	1.358 (5)	C11—C12	1.408 (6)
N3—C3	1.335 (5)	C12—C15	1.397 (7)
N3—C4	1.364 (5)	C12—C13	1.430 (7)
N3—H3	0.8600	C13—C14	1.331 (8)
N4—C5	1.365 (5)	C13—H13	0.9300
N4—C4	1.382 (5)	C14—H14	0.9300
N4—H4	0.8600	C15—C16	1.349 (7)
O1—C5	1.253 (4)	C15—H15	0.9300
O2—C1	1.279 (4)	C16—C17	1.386 (6)
O3—C1	1.239 (5)	C16—H16	0.9300
O4—C4	1.239 (5)	C17—H17	0.9300
C1—C2	1.488 (5)	O5—H51	0.8483
C2—C3	1.363 (6)	O5—H52	0.8510
O2 ⁱ —Dy—O2	88.73 (14)	C5—C2—C1	123.4 (3)
O2 ⁱ —Dy—O1 ⁱ	74.32 (9)	N3—C3—C2	126.1 (4)
O2—Dy—O1 ⁱ	81.97 (10)	N3—C3—H3A	117.0
O2 ⁱ —Dy—O1	81.97 (10)	C2—C3—H3A	117.0
O2—Dy—O1	74.32 (9)	O4—C4—N3	122.5 (4)
O1 ⁱ —Dy—O1	146.67 (13)	O4—C4—N4	121.7 (4)
O2 ⁱ —Dy—N2	147.90 (10)	N3—C4—N4	115.8 (3)
O2—Dy—N2	105.34 (10)	O1—C5—N4	117.4 (3)
O1 ⁱ —Dy—N2	135.32 (10)	O1—C5—C2	126.3 (3)
O1—Dy—N2	74.62 (10)	N4—C5—C2	116.3 (3)
O2 ⁱ —Dy—N2 ⁱ	105.34 (10)	N2—C6—C7	123.4 (4)
O2—Dy—N2 ⁱ	147.90 (10)	N2—C6—H6	118.3
O1 ⁱ —Dy—N2 ⁱ	74.62 (10)	C7—C6—H6	118.3
O1—Dy—N2 ⁱ	135.32 (10)	C8—C7—C6	118.7 (4)

supplementary materials

N2—Dy—N2 ⁱ	77.95 (15)	C8—C7—H7	120.6
O2 ⁱ —Dy—N1 ⁱ	79.80 (10)	C6—C7—H7	120.6
O2—Dy—N1 ⁱ	148.05 (10)	C7—C8—C9	120.6 (4)
O1 ⁱ —Dy—N1 ⁱ	122.34 (10)	C7—C8—H8	119.7
O1—Dy—N1 ⁱ	74.60 (9)	C9—C8—H8	119.7
N2—Dy—N1 ⁱ	73.07 (10)	C8—C9—C10	116.9 (4)
N2 ⁱ —Dy—N1 ⁱ	63.96 (10)	C8—C9—C14	123.8 (4)
O2 ⁱ —Dy—N1	148.05 (10)	C10—C9—C14	119.3 (4)
O2—Dy—N1	79.80 (10)	N2—C10—C9	122.7 (4)
O1 ⁱ —Dy—N1	74.60 (9)	N2—C10—C11	118.3 (3)
O1—Dy—N1	122.34 (10)	C9—C10—C11	119.0 (4)
N2—Dy—N1	63.96 (10)	N1—C11—C12	122.6 (4)
N2 ⁱ —Dy—N1	73.07 (10)	N1—C11—C10	117.7 (3)
N1 ⁱ —Dy—N1	123.98 (14)	C12—C11—C10	119.8 (4)
C17—N1—C11	117.3 (4)	C15—C12—C11	117.2 (4)
C17—N1—Dy	123.9 (3)	C15—C12—C13	123.8 (4)
C11—N1—Dy	117.0 (2)	C11—C12—C13	118.9 (4)
C6—N2—C10	117.6 (3)	C14—C13—C12	121.6 (5)
C6—N2—Dy	123.4 (3)	C14—C13—H13	119.2
C10—N2—Dy	117.5 (2)	C12—C13—H13	119.2
C3—N3—C4	119.6 (3)	C13—C14—C9	121.3 (5)
C3—N3—H3	120.2	C13—C14—H14	119.3
C4—N3—H3	120.2	C9—C14—H14	119.3
C5—N4—C4	126.0 (3)	C16—C15—C12	120.0 (4)
C5—N4—H4	117.0	C16—C15—H15	120.0
C4—N4—H4	117.0	C12—C15—H15	120.0
C5—O1—Dy	132.2 (2)	C15—C16—C17	119.4 (4)
C1—O2—Dy	140.8 (2)	C15—C16—H16	120.3
O3—C1—O2	123.1 (3)	C17—C16—H16	120.3
O3—C1—C2	118.8 (3)	N1—C17—C16	123.3 (4)
O2—C1—C2	118.1 (3)	N1—C17—H17	118.3
C3—C2—C5	116.1 (3)	C16—C17—H17	118.3
C3—C2—C1	120.5 (3)	H51—O5—H52	112.0
O2 ⁱ —Dy—N1—C17	−6.9 (4)	C1—C2—C3—N3	−178.9 (4)
O2—Dy—N1—C17	63.8 (3)	C3—N3—C4—O4	−178.8 (4)
O1 ⁱ —Dy—N1—C17	−20.6 (3)	C3—N3—C4—N4	1.0 (6)
O1—Dy—N1—C17	127.8 (3)	C5—N4—C4—O4	−176.8 (4)
N2—Dy—N1—C17	176.6 (3)	C5—N4—C4—N3	3.4 (6)
N2 ⁱ —Dy—N1—C17	−98.8 (3)	Dy—O1—C5—N4	−158.4 (3)
N1 ⁱ —Dy—N1—C17	−139.5 (3)	Dy—O1—C5—C2	22.1 (6)
O2 ⁱ —Dy—N1—C11	157.4 (2)	C4—N4—C5—O1	174.7 (4)
O2—Dy—N1—C11	−132.0 (3)	C4—N4—C5—C2	−5.7 (6)
O1 ⁱ —Dy—N1—C11	143.6 (3)	C3—C2—C5—O1	−177.0 (4)
O1—Dy—N1—C11	−68.0 (3)	C1—C2—C5—O1	2.6 (6)
N2—Dy—N1—C11	−19.2 (3)	C3—C2—C5—N4	3.4 (5)

N2 ⁱ —Dy—N1—C11	65.4 (3)	C1—C2—C5—N4	-177.0 (3)
N1 ⁱ —Dy—N1—C11	24.7 (2)	C10—N2—C6—C7	2.7 (6)
O2 ⁱ —Dy—N2—C6	8.4 (4)	Dy—N2—C6—C7	-162.9 (3)
O2—Dy—N2—C6	-104.8 (3)	N2—C6—C7—C8	-0.4 (7)
O1 ⁱ —Dy—N2—C6	161.1 (3)	C6—C7—C8—C9	-0.9 (7)
O1—Dy—N2—C6	-36.2 (3)	C7—C8—C9—C10	-0.2 (7)
N2 ⁱ —Dy—N2—C6	108.1 (3)	C7—C8—C9—C14	-179.1 (5)
N1 ⁱ —Dy—N2—C6	42.0 (3)	C6—N2—C10—C9	-3.9 (6)
N1—Dy—N2—C6	-175.0 (3)	Dy—N2—C10—C9	162.6 (3)
O2 ⁱ —Dy—N2—C10	-157.3 (3)	C6—N2—C10—C11	174.8 (4)
O2—Dy—N2—C10	89.6 (3)	Dy—N2—C10—C11	-18.7 (5)
O1 ⁱ —Dy—N2—C10	-4.5 (3)	C8—C9—C10—N2	2.7 (6)
O1—Dy—N2—C10	158.1 (3)	C14—C9—C10—N2	-178.4 (4)
N2 ⁱ —Dy—N2—C10	-57.6 (2)	C8—C9—C10—C11	-176.0 (4)
N1 ⁱ —Dy—N2—C10	-123.7 (3)	C14—C9—C10—C11	2.9 (6)
N1—Dy—N2—C10	19.3 (3)	C17—N1—C11—C12	2.9 (6)
O2 ⁱ —Dy—O1—C5	68.0 (3)	Dy—N1—C11—C12	-162.4 (3)
O2—Dy—O1—C5	-22.9 (3)	C17—N1—C11—C10	-176.5 (4)
O1 ⁱ —Dy—O1—C5	23.4 (3)	Dy—N1—C11—C10	18.3 (4)
N2—Dy—O1—C5	-134.2 (4)	N2—C10—C11—N1	0.1 (5)
N2 ⁱ —Dy—O1—C5	171.7 (3)	C9—C10—C11—N1	178.8 (4)
N1 ⁱ —Dy—O1—C5	149.6 (4)	N2—C10—C11—C12	-179.3 (4)
N1—Dy—O1—C5	-89.6 (3)	C9—C10—C11—C12	-0.6 (6)
O2 ⁱ —Dy—O2—C1	-72.1 (4)	N1—C11—C12—C15	-1.3 (7)
O1 ⁱ —Dy—O2—C1	-146.4 (4)	C10—C11—C12—C15	178.1 (4)
O1—Dy—O2—C1	9.9 (4)	N1—C11—C12—C13	178.6 (4)
N2—Dy—O2—C1	78.7 (4)	C10—C11—C12—C13	-2.0 (6)
N2 ⁱ —Dy—O2—C1	170.4 (4)	C15—C12—C13—C14	-177.8 (6)
N1 ⁱ —Dy—O2—C1	-3.8 (5)	C11—C12—C13—C14	2.3 (8)
N1—Dy—O2—C1	137.9 (4)	C12—C13—C14—C9	0.1 (9)
Dy—O2—C1—O3	-176.1 (3)	C8—C9—C14—C13	176.1 (5)
Dy—O2—C1—C2	4.2 (6)	C10—C9—C14—C13	-2.8 (8)
O3—C1—C2—C3	-15.5 (6)	C11—C12—C15—C16	-1.7 (8)
O2—C1—C2—C3	164.1 (4)	C13—C12—C15—C16	178.4 (5)
O3—C1—C2—C5	164.9 (4)	C12—C15—C16—C17	2.9 (8)
O2—C1—C2—C5	-15.4 (6)	C11—N1—C17—C16	-1.7 (6)
C4—N3—C3—C2	-3.0 (7)	Dy—N1—C17—C16	162.5 (3)
C5—C2—C3—N3	0.7 (7)	C15—C16—C17—N1	-1.2 (8)

Symmetry codes: (i) $-x, y, -z+3/2$.

Hydrogen-bond geometry (\AA , $^\circ$)


$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N3—H3 \cdots N3 ⁱⁱ	0.86	1.80	2.659 (7)	174
N4—H4 \cdots O3 ⁱⁱⁱ	0.86	2.01	2.867 (4)	178

supplementary materials

N4—H4···O2 ⁱⁱⁱ	0.86	2.62	3.183 (4)	124
O5—H51···O3 ^{iv}	0.85	2.15	2.993 (5)	175
O5—H52···O4 ^v	0.85	2.15	2.960 (5)	160

Symmetry codes: (ii) $-x+1/2, -y+1/2, -z+2$; (iii) $x, -y+1, z+1/2$; (iv) $-x+1/2, -y+1/2, -z+1$; (v) $x, y, z-1$.

Fig. 1



supplementary materials

Fig. 2

