

1-Allyl-3-chloro-6-nitro-1*H*-indazole

Nabil El Brahmi,^a Benchidmi Mohamed,^a El Mokhtar Essassi,^a Hafid Zouihri^b and Seik Weng Ng^{c*}

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Université Mohammed V-Agdal, BP 1014 Avenue Ibn Batout, Rabat, Morocco, ^bCNRST Division UATRS, Angle Allal Fassi/FAR, BP 8027 Hay Riad, 10000 Rabat, Morocco, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 25 August 2009; accepted 26 August 2009

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$; R factor = 0.056; wR factor = 0.185; data-to-parameter ratio = 13.1.

The indazole system in each of the two independent molecules of the title compound, $\text{C}_{10}\text{H}_8\text{ClN}_3\text{O}_2$, is planar (r.m.s. deviations = 0.005 and 0.005 Å). The nitro group is coplanar with the fused-ring system [dihedral angles = 1.3 (3) and 4.8 (3) Å].

Related literature

For a review of indazoles, see: Elguero (1996); Elguero *et al.* (1995).



Experimental

Crystal data

$\text{C}_{10}\text{H}_8\text{ClN}_3\text{O}_2$
 $M_r = 237.64$
Monoclinic, $P2_1/n$
 $a = 7.6804 (1)\text{ \AA}$
 $b = 9.9559 (2)\text{ \AA}$
 $c = 28.4344 (4)\text{ \AA}$
 $\beta = 95.144 (1)^\circ$

$V = 2165.49 (6)\text{ \AA}^3$
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.34\text{ mm}^{-1}$
 $T = 295\text{ K}$
 $0.4 \times 0.3 \times 0.2\text{ mm}$

Data collection

Bruker APEXII diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.884$, $T_{\max} = 0.934$

19833 measured reflections
3777 independent reflections
2665 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.032$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.185$
 $S = 1.07$
3777 reflections

289 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.78\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.29\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank Université Mohammed V-Agdal and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2531).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2005). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Elguero, J. (1996). *Comprehensive Heterocyclic Chemistry II*, edited by I. Shinkai, Vol. 3, p. 1. Oxford: Elsevier Science.
Elguero, J., Fruchier, A., Tjiou, E. M. & Trofimenko, S. (1995). *Chem. Heterocycl. Compd. (Engl. Transl.)*, **31**, 1006–1026.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o2320 [doi:10.1107/S1600536809034138]

1-Allyl-3-chloro-6-nitro-1*H*-indazole

N. El Brahmi, B. Mohamed, E. M. Essassi, H. Zouihri and S. W. Ng

Experimental

3-Chloro-6-nitroindazole (5 mmol) and allyl bromide (10 mmol) were reacted in THF (40 ml) in the presence of potassium carbonate (10 mmol) and tetra-*n*-butylammonium bromide (0.5 mmol). The mixture was stirred for 24 h, filtered, and the THF removed under vacuum. The product was separated by chromatography on silica gel with a hexane:ethyl acetate (9:1) solvent system. The compound was obtained as yellow crystals in 50% yield; m.p. 351 K.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.97 Å) and were included in the refinement in the riding model approximation with $U(\text{H})$ set to $1.2U(\text{C})$.

Although data were measured to a high 2θ limit, those reflections beyond 50° were not used as their inclusion significantly raised the R index.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of the two independent molecules of $\text{C}_{10}\text{H}_8\text{ClN}_3\text{O}_2$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

1-Allyl-3-chloro-6-nitro-1*H*-indazole

Crystal data

$\text{C}_{10}\text{H}_8\text{ClN}_3\text{O}_2$	$F_{000} = 976$
$M_r = 237.64$	$D_x = 1.458 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	$\text{Mo } K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 6549 reflections
$a = 7.6804 (1) \text{ \AA}$	$\theta = 2.2\text{--}29.5^\circ$
$b = 9.9559 (2) \text{ \AA}$	$\mu = 0.34 \text{ mm}^{-1}$
$c = 28.4344 (4) \text{ \AA}$	$T = 295 \text{ K}$
$\beta = 95.144 (1)^\circ$	Prism, yellow
$V = 2165.49 (6) \text{ \AA}^3$	$0.4 \times 0.3 \times 0.2 \text{ mm}$
$Z = 8$	

supplementary materials

Data collection

Bruker APEX2 diffractometer	3777 independent reflections
Radiation source: fine-focus sealed tube	2665 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.032$
$T = 295 \text{ K}$	$\theta_{\text{max}} = 25.0^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.884, T_{\text{max}} = 0.934$	$k = -11 \rightarrow 11$
19833 measured reflections	$l = -33 \rightarrow 33$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.056$	H-atom parameters constrained
$wR(F^2) = 0.185$	$w = 1/[\sigma^2(F_o^2) + (0.1031P)^2 + 0.7179P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} = 0.001$
3777 reflections	$\Delta\rho_{\text{max}} = 0.78 \text{ e \AA}^{-3}$
289 parameters	$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.30814 (12)	0.06578 (10)	0.03474 (3)	0.0718 (3)
Cl2	0.94159 (18)	0.35838 (12)	0.02733 (4)	0.0958 (4)
O1	0.6481 (4)	0.0762 (3)	0.28950 (9)	0.0866 (9)
O2	0.7296 (4)	0.2788 (3)	0.27899 (9)	0.0867 (9)
O3	1.2186 (4)	0.2329 (4)	0.28256 (10)	0.1089 (11)
O4	1.0809 (4)	0.0460 (3)	0.27538 (10)	0.0928 (9)
N1	0.5287 (3)	0.3527 (3)	0.10386 (9)	0.0563 (7)
N2	0.4505 (4)	0.2960 (3)	0.06380 (9)	0.0606 (8)
N3	0.6601 (4)	0.1750 (3)	0.26458 (9)	0.0582 (7)
N4	0.8630 (3)	0.0501 (3)	0.09910 (10)	0.0529 (7)
N5	0.8504 (4)	0.1220 (3)	0.05846 (10)	0.0622 (8)
N6	1.1300 (4)	0.1509 (4)	0.25934 (11)	0.0659 (8)
C1	0.4072 (4)	0.1742 (4)	0.07565 (11)	0.0523 (8)
C2	0.4547 (3)	0.1450 (3)	0.12352 (11)	0.0451 (7)
C3	0.4413 (4)	0.0352 (3)	0.15351 (11)	0.0509 (8)
H3	0.3882	-0.0441	0.1424	0.061*
C4	0.5079 (4)	0.0470 (3)	0.19951 (11)	0.0510 (8)

H4	0.5003	-0.0244	0.2203	0.061*
C5	0.5876 (4)	0.1675 (3)	0.21511 (10)	0.0458 (7)
C6	0.6039 (4)	0.2781 (3)	0.18747 (10)	0.0447 (7)
H6	0.6570	0.3568	0.1991	0.054*
C7	0.5353 (4)	0.2648 (3)	0.14059 (11)	0.0447 (7)
C8	0.6026 (5)	0.4877 (4)	0.10227 (14)	0.0678 (10)
H8A	0.6737	0.5039	0.1317	0.081*
H8B	0.6793	0.4909	0.0770	0.081*
C9	0.4777 (6)	0.5939 (5)	0.0952 (2)	0.0994 (15)
H9	0.4034	0.6041	0.1191	0.119*
C10	0.4537 (7)	0.6752 (5)	0.0618 (2)	0.1091 (18)
H10A	0.5232	0.6710	0.0366	0.131*
H10B	0.3663	0.7398	0.0620	0.131*
C11	0.9279 (4)	0.2371 (4)	0.06933 (11)	0.0588 (9)
C12	0.9941 (4)	0.2448 (3)	0.11660 (11)	0.0488 (8)
C13	1.0839 (4)	0.3397 (3)	0.14605 (13)	0.0563 (8)
H13	1.1130	0.4232	0.1343	0.068*
C14	1.1278 (4)	0.3072 (4)	0.19233 (13)	0.0563 (9)
H14	1.1885	0.3681	0.2124	0.068*
C15	1.0808 (4)	0.1816 (3)	0.20921 (11)	0.0491 (8)
C16	0.9938 (3)	0.0852 (3)	0.18217 (11)	0.0453 (7)
H16	0.9651	0.0023	0.1944	0.054*
C17	0.9510 (3)	0.1194 (3)	0.13522 (11)	0.0435 (7)
C18	0.8035 (4)	-0.0888 (3)	0.09845 (13)	0.0615 (9)
H18A	0.6969	-0.0963	0.0776	0.074*
H18B	0.7766	-0.1142	0.1299	0.074*
C19	0.9352 (5)	-0.1820 (4)	0.08251 (14)	0.0701 (10)
H19	0.9748	-0.1658	0.0531	0.084*
C20	0.9988 (6)	-0.2813 (5)	0.1052 (2)	0.1019 (16)
H20A	0.9629	-0.3012	0.1348	0.122*
H20B	1.0817	-0.3350	0.0924	0.122*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0785 (6)	0.0739 (7)	0.0602 (6)	-0.0018 (5)	-0.0089 (4)	-0.0182 (4)
Cl2	0.1491 (11)	0.0756 (9)	0.0649 (6)	0.0153 (7)	0.0215 (6)	0.0187 (5)
O1	0.128 (2)	0.072 (2)	0.0577 (15)	0.0029 (16)	-0.0050 (15)	0.0184 (15)
O2	0.116 (2)	0.081 (2)	0.0593 (16)	-0.0218 (17)	-0.0108 (15)	-0.0064 (14)
O3	0.126 (2)	0.121 (3)	0.0733 (19)	-0.021 (2)	-0.0282 (18)	-0.0127 (19)
O4	0.133 (3)	0.074 (2)	0.0688 (18)	0.0066 (18)	-0.0030 (16)	0.0176 (16)
N1	0.0649 (16)	0.0493 (18)	0.0536 (16)	-0.0039 (13)	-0.0018 (12)	0.0018 (13)
N2	0.0677 (16)	0.063 (2)	0.0494 (16)	0.0022 (14)	-0.0031 (13)	-0.0010 (14)
N3	0.0651 (16)	0.060 (2)	0.0495 (16)	0.0074 (14)	0.0058 (13)	-0.0029 (15)
N4	0.0516 (14)	0.0464 (18)	0.0606 (17)	0.0000 (12)	0.0055 (12)	-0.0086 (13)
N5	0.0684 (17)	0.062 (2)	0.0560 (17)	0.0127 (15)	0.0025 (13)	-0.0061 (15)
N6	0.0645 (17)	0.071 (2)	0.0614 (19)	0.0134 (16)	-0.0007 (15)	-0.0084 (18)
C1	0.0503 (16)	0.053 (2)	0.0530 (19)	0.0056 (14)	0.0033 (14)	-0.0088 (16)

supplementary materials

C2	0.0388 (14)	0.043 (2)	0.0543 (18)	0.0061 (12)	0.0063 (13)	-0.0087 (14)
C3	0.0506 (16)	0.041 (2)	0.062 (2)	0.0009 (13)	0.0101 (14)	-0.0064 (16)
C4	0.0567 (17)	0.042 (2)	0.0556 (19)	0.0068 (14)	0.0119 (15)	0.0031 (15)
C5	0.0443 (15)	0.047 (2)	0.0469 (17)	0.0094 (13)	0.0072 (13)	-0.0022 (14)
C6	0.0449 (15)	0.0386 (19)	0.0504 (17)	0.0050 (12)	0.0027 (13)	-0.0055 (14)
C7	0.0414 (14)	0.0427 (19)	0.0503 (17)	0.0079 (13)	0.0066 (12)	-0.0006 (15)
C8	0.078 (2)	0.054 (2)	0.069 (2)	-0.0089 (19)	-0.0050 (18)	0.0130 (18)
C9	0.085 (3)	0.078 (3)	0.134 (4)	-0.011 (2)	0.000 (3)	0.021 (3)
C10	0.108 (3)	0.080 (4)	0.132 (4)	-0.019 (3)	-0.023 (3)	0.046 (3)
C11	0.072 (2)	0.050 (2)	0.055 (2)	0.0154 (17)	0.0094 (17)	0.0019 (16)
C12	0.0496 (16)	0.040 (2)	0.0581 (18)	0.0094 (13)	0.0139 (14)	-0.0024 (15)
C13	0.0630 (19)	0.036 (2)	0.072 (2)	-0.0020 (14)	0.0176 (17)	-0.0019 (16)
C14	0.0510 (16)	0.047 (2)	0.072 (2)	-0.0019 (14)	0.0107 (15)	-0.0194 (17)
C15	0.0446 (15)	0.045 (2)	0.0577 (19)	0.0105 (13)	0.0057 (14)	-0.0034 (15)
C16	0.0460 (15)	0.0322 (18)	0.0589 (18)	0.0040 (12)	0.0105 (13)	0.0000 (14)
C17	0.0389 (14)	0.0354 (18)	0.0567 (18)	0.0060 (12)	0.0075 (13)	-0.0060 (14)
C18	0.0579 (18)	0.050 (2)	0.077 (2)	-0.0065 (15)	0.0049 (16)	-0.0166 (18)
C19	0.085 (2)	0.056 (3)	0.072 (2)	0.001 (2)	0.0204 (19)	-0.002 (2)
C20	0.096 (3)	0.082 (4)	0.131 (4)	0.013 (3)	0.030 (3)	0.027 (3)

Geometric parameters (\AA , $^\circ$)

C11—C1	1.714 (3)	C6—H6	0.9300
C12—C11	1.708 (4)	C8—C9	1.430 (6)
O1—N3	1.221 (4)	C8—H8A	0.9700
O2—N3	1.218 (4)	C8—H8B	0.9700
O3—N6	1.219 (4)	C9—C10	1.250 (6)
O4—N6	1.213 (4)	C9—H9	0.9300
N1—C7	1.360 (4)	C10—H10A	0.9300
N1—N2	1.362 (4)	C10—H10B	0.9300
N1—C8	1.461 (4)	C11—C12	1.396 (4)
N2—C1	1.309 (5)	C12—C13	1.402 (5)
N3—C5	1.468 (4)	C12—C17	1.408 (4)
N4—N5	1.356 (4)	C13—C14	1.368 (5)
N4—C17	1.365 (4)	C13—H13	0.9300
N4—C18	1.456 (4)	C14—C15	1.398 (5)
N5—C11	1.316 (5)	C14—H14	0.9300
N6—C15	1.474 (4)	C15—C16	1.367 (4)
C1—C2	1.408 (4)	C16—C17	1.388 (4)
C2—C3	1.396 (4)	C16—H16	0.9300
C2—C7	1.410 (4)	C18—C19	1.474 (5)
C3—C4	1.366 (4)	C18—H18A	0.9700
C3—H3	0.9300	C18—H18B	0.9700
C4—C5	1.401 (4)	C19—C20	1.255 (6)
C4—H4	0.9300	C19—H19	0.9300
C5—C6	1.365 (4)	C20—H20A	0.9300
C6—C7	1.395 (4)	C20—H20B	0.9300
C7—N1—N2	111.0 (3)	H8A—C8—H8B	107.5
C7—N1—C8	128.9 (3)	C10—C9—C8	129.6 (6)

N2—N1—C8	119.9 (3)	C10—C9—H9	115.2
C1—N2—N1	105.8 (3)	C8—C9—H9	115.2
O2—N3—O1	122.8 (3)	C9—C10—H10A	120.0
O2—N3—C5	118.9 (3)	C9—C10—H10B	120.0
O1—N3—C5	118.3 (3)	H10A—C10—H10B	120.0
N5—N4—C17	111.6 (3)	N5—C11—C12	113.1 (3)
N5—N4—C18	119.5 (3)	N5—C11—Cl2	120.4 (3)
C17—N4—C18	128.6 (3)	C12—C11—Cl2	126.5 (3)
C11—N5—N4	105.3 (3)	C11—C12—C13	136.8 (3)
O4—N6—O3	123.4 (4)	C11—C12—C17	103.6 (3)
O4—N6—C15	118.8 (3)	C13—C12—C17	119.5 (3)
O3—N6—C15	117.8 (4)	C14—C13—C12	118.7 (3)
N2—C1—C2	113.0 (3)	C14—C13—H13	120.6
N2—C1—Cl1	121.1 (3)	C12—C13—H13	120.6
C2—C1—Cl1	125.9 (3)	C13—C14—C15	119.6 (3)
C3—C2—C1	136.5 (3)	C13—C14—H14	120.2
C3—C2—C7	120.4 (3)	C15—C14—H14	120.2
C1—C2—C7	103.1 (3)	C16—C15—C14	124.3 (3)
C4—C3—C2	118.4 (3)	C16—C15—N6	117.9 (3)
C4—C3—H3	120.8	C14—C15—N6	117.8 (3)
C2—C3—H3	120.8	C15—C16—C17	115.5 (3)
C3—C4—C5	119.5 (3)	C15—C16—H16	122.2
C3—C4—H4	120.3	C17—C16—H16	122.2
C5—C4—H4	120.3	N4—C17—C16	131.2 (3)
C6—C5—C4	124.7 (3)	N4—C17—C12	106.4 (3)
C6—C5—N3	117.7 (3)	C16—C17—C12	122.4 (3)
C4—C5—N3	117.7 (3)	N4—C18—C19	112.2 (3)
C5—C6—C7	115.3 (3)	N4—C18—H18A	109.2
C5—C6—H6	122.4	C19—C18—H18A	109.2
C7—C6—H6	122.4	N4—C18—H18B	109.2
N1—C7—C6	131.1 (3)	C19—C18—H18B	109.2
N1—C7—C2	107.2 (3)	H18A—C18—H18B	107.9
C6—C7—C2	121.8 (3)	C20—C19—C18	125.7 (4)
C9—C8—N1	115.3 (3)	C20—C19—H19	117.1
C9—C8—H8A	108.5	C18—C19—H19	117.1
N1—C8—H8A	108.5	C19—C20—H20A	120.0
C9—C8—H8B	108.5	C19—C20—H20B	120.0
N1—C8—H8B	108.5	H20A—C20—H20B	120.0
C7—N1—N2—C1	0.7 (3)	N2—N1—C8—C9	70.4 (5)
C8—N1—N2—C1	176.4 (3)	N1—C8—C9—C10	-116.9 (5)
C17—N4—N5—C11	-1.0 (3)	N4—N5—C11—C12	0.3 (4)
C18—N4—N5—C11	-174.4 (3)	N4—N5—C11—Cl2	179.8 (2)
N1—N2—C1—C2	-0.2 (4)	N5—C11—C12—C13	-179.6 (3)
N1—N2—C1—Cl1	-178.6 (2)	Cl2—C11—C12—C13	0.9 (6)
N2—C1—C2—C3	-179.7 (3)	N5—C11—C12—C17	0.4 (3)
Cl1—C1—C2—C3	-1.4 (5)	Cl2—C11—C12—C17	-179.0 (2)
N2—C1—C2—C7	-0.3 (3)	C11—C12—C13—C14	-179.7 (3)
Cl1—C1—C2—C7	178.1 (2)	C17—C12—C13—C14	0.2 (4)
C1—C2—C3—C4	179.3 (3)	C12—C13—C14—C15	-0.7 (4)

supplementary materials

C7—C2—C3—C4	0.0 (4)	C13—C14—C15—C16	0.9 (5)
C2—C3—C4—C5	-0.3 (4)	C13—C14—C15—N6	-179.7 (3)
C3—C4—C5—C6	0.5 (4)	O4—N6—C15—C16	-5.2 (4)
C3—C4—C5—N3	-179.0 (3)	O3—N6—C15—C16	174.8 (3)
O2—N3—C5—C6	0.7 (4)	O4—N6—C15—C14	175.4 (3)
O1—N3—C5—C6	-179.1 (3)	O3—N6—C15—C14	-4.6 (4)
O2—N3—C5—C4	-179.8 (3)	C14—C15—C16—C17	-0.5 (4)
O1—N3—C5—C4	0.4 (4)	N6—C15—C16—C17	-179.9 (2)
C4—C5—C6—C7	-0.5 (4)	N5—N4—C17—C16	-180.0 (3)
N3—C5—C6—C7	179.0 (2)	C18—N4—C17—C16	-7.3 (5)
N2—N1—C7—C6	179.3 (3)	N5—N4—C17—C12	1.3 (3)
C8—N1—C7—C6	4.0 (5)	C18—N4—C17—C12	174.0 (3)
N2—N1—C7—C2	-0.8 (3)	C15—C16—C17—N4	-178.6 (3)
C8—N1—C7—C2	-176.1 (3)	C15—C16—C17—C12	0.0 (4)
C5—C6—C7—N1	-180.0 (3)	C11—C12—C17—N4	-1.0 (3)
C5—C6—C7—C2	0.2 (4)	C13—C12—C17—N4	179.0 (3)
C3—C2—C7—N1	-179.8 (3)	C11—C12—C17—C16	-179.9 (3)
C1—C2—C7—N1	0.7 (3)	C13—C12—C17—C16	0.2 (4)
C3—C2—C7—C6	0.1 (4)	N5—N4—C18—C19	81.6 (4)
C1—C2—C7—C6	-179.5 (3)	C17—N4—C18—C19	-90.5 (4)
C7—N1—C8—C9	-114.7 (4)	N4—C18—C19—C20	123.4 (5)

Fig. 1

