

2,2'-(1,3,5,7-Tetraoxo-1,2,3,5,6,7-hexahydropyrrolo[3,4-f]isoindole-2,6-diyl)-diacetic acid *N,N*-dimethylformamide disolvate

Chunhua Ge,* Xiangqian Li, Xiangdong Zhang, Yang Zhao and Rui Zhang

College of Chemistry, Liaoning University, Shenyang 110036, People's Republic of China

Correspondence e-mail: chhge@lnu.edu.cn


Received 25 August 2009; accepted 1 September 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; disorder in solvent or counterion; R factor = 0.039; wR factor = 0.111; data-to-parameter ratio = 13.5.

The asymmetric unit of the title compound, $\text{C}_{14}\text{H}_8\text{N}_2\text{O}_8 \cdot 2\text{C}_3\text{H}_7\text{NO}$ or $L \cdot 2\text{DMF}$ (DMF = *N,N*-dimethylformamide), contains one half of the centrosymmetric molecule L and one solvent molecule, which is disordered between two orientations in a 0.555 (4):0.445 (4) ratio. Intermolecular $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds link one L and two DMF molecules into a centrosymmetric hydrogen-bonded cluster. The crystal packing is further stabilized by weak intermolecular $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds.

Related literature

For recent developments in the chemistry of naphthalene diimides, see Bhosale *et al.* (2008). For pyromellitic diimides, see: Gabriel & Iverson (2002); Ghosh & Ramakrishnan (2005); Kimizuka *et al.* (1995). For details of the synthesis, see Barooah *et al.* (2006).



Experimental

Crystal data

$\text{C}_{14}\text{H}_8\text{N}_2\text{O}_8 \cdot 2\text{C}_3\text{H}_7\text{NO}$

$M_r = 478.42$

Monoclinic, $P2_1/c$
 $a = 7.7470 (15)\text{ \AA}$
 $b = 9.3100 (19)\text{ \AA}$
 $c = 16.334 (5)\text{ \AA}$
 $\beta = 104.02 (3)^\circ$
 $V = 1143.0 (5)\text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.11\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.30 \times 0.25 \times 0.25\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.958$, $T_{\max} = 0.973$

6227 measured reflections
2236 independent reflections
1910 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.111$
 $S = 1.06$
2236 reflections

166 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.15\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
O3-H3A \cdots O5A	0.82	1.72	2.526 (2)	166
O4-H4B \cdots O5B	0.82	1.70	2.496 (2)	162
C2-H2A \cdots O3 ⁱ	0.97	2.41	3.230 (2)	142
C6-H6 \cdots O1 ⁱⁱ	0.93	2.50	3.414 (2)	166
C9-H9A \cdots O2 ⁱⁱⁱ	0.96	2.57	3.432 (3)	149

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x - 1, y, z$; (iii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Project for Innovation Teams of Liaoning Province, China (No. 2007 T052), the Project for Provincial Key Laboratories of Liaoning Province, China (No. 2008S104) and the Doctoral Startup Project, Liaoning University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2607).

References

- Barooah, N., Sarma, R. J. & Baruah, J. B. (2006). *CrystEngComm*, **8**, 608–615.
Bhosale, S. V., Jani, C. H. & Langford, S. J. (2008). *Chem. Soc. Rev.* **37**, 331–342.
Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Gabriel, G. J. & Iverson, B. L. (2002). *J. Am. Chem. Soc.* **124**, 15174–15175.
Ghosh, S. & Ramakrishnan, S. (2005). *Angew. Chem. Int. Ed.* **44**, 5441–5447.
Kimizuka, N., Kawasaki, T., Hirata, K. & Kunitake, T. (1995). *J. Am. Chem. Soc.* **117**, 6360–6361.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o2400 [doi:10.1107/S1600536809035107]

2,2'-(1,3,5,7-Tetraoxo-1,2,3,5,6,7-hexahydropyrrolo[3,4-*f*]isoindole-2,6-diyl)diacetic acid *N,N*-dimethylformamide disolvate

C. Ge, X. Li, X. Zhang, Y. Zhao and R. Zhang

Comment

Interest in the derivatives of diimide such as pyromellitic diimides, naphthalene diimides, and perylene diimides has arisen because of their potential applications for supramolecular and new functional materials (Bhosale *et al.*, 2008). There have been a number of studies investigating the host–guest chemistry about pyromellitic diimide (Gabriel & Iverson, 2002; Ghosh & Ramakrishnan, 2005; Kimizuka *et al.*, 1995). Supramolecular host of *L* and inclusion compounds with aromatic guests have been described (Barooah *et al.*, 2006). In this paper, we report the crystal structure of the title compound, obtained by the recrystallization in DMF-MeOH.

In the molecule *L* (=2,2'-(1,3,5,7-tetraoxo-5,7-dihydropyrrolo[3,4-*f*]isoindole-2,6-diyl)diacetic acid), two acetic acid groups are placed on upper and lower sides of the rigid conjugate plane (Fig. 1). Intermolecular O—H···O hydrogen bonds (Table 1) link one *L* and two solvent molecules into centrosymmetric hydrogen-bonded cluster. The crystal packing is further stabilized by weak intermolecular C—H···O hydrogen bonds (Table 1).


Experimental

2,2'-(1,3,5,7-Tetraoxo-5,7-dihydropyrrolo[3,4-*f*]isoindole-2,6-diyl)diacetic acid was synthesized according to the literature (Barooah *et al.*, 2006). *N,N*'-dimethylformamide (DMF) 5 ml was added into a solution of compound mentioned above 0.1 mmol in 20 ml MeOH. The resultant colourless solution was filtered. Crystals suitable for X-ray analysis were formed after three days at room temperature.

Refinement

All H atoms were placed in calculated positions and included in a riding-model approximation, with C—H = 0.93 - 0.97 Å, O—H = 0.82 Å and $U_{\text{iso}}(\text{H})$ = 1.2-1.5 U_{eq} of the parent atom. The solvent molecule is disordered between two orientations with the occupancies refined to 0.555 (4) and 0.445 (4), respectively. The hydroxy H atom is also disordered between two positions - H3A and H4B - with the same occupancies, respectively.

Figures



supplementary materials

2,2'-(1,3,5,7-Tetraoxo-1,2,3,5,6,7-hexahydropyrrolo[3,4-f]isoindole- 2,6-diyl)diacetic acid *N,N*-dimethylformamide disolvate

Crystal data

C ₁₄ H ₈ N ₂ O ₈ ·2C ₃ H ₇ NO	$F_{000} = 500$
$M_r = 478.42$	$D_x = 1.390 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 541 reflections
$a = 7.7470 (15) \text{ \AA}$	$\theta = 2.5\text{--}22.7^\circ$
$b = 9.3100 (19) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$c = 16.334 (5) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 104.02 (3)^\circ$	Block, colourless
$V = 1143.0 (5) \text{ \AA}^3$	$0.30 \times 0.25 \times 0.25 \text{ mm}$
$Z = 2$	

Data collection

Bruker SMART CCD area-detector diffractometer	2236 independent reflections
Radiation source: fine-focus sealed tube	1910 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.030$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 26.0^\circ$
ϕ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.958$, $T_{\text{max}} = 0.973$	$k = -11 \rightarrow 10$
6227 measured reflections	$l = -20 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.111$	$w = 1/[\sigma^2(F_o^2) + (0.0528P)^2 + 0.2565P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.06$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2236 reflections	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$
166 parameters	$\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O3	0.46550 (18)	0.49571 (11)	0.66856 (7)	0.0615 (3)	
H3A	0.4139	0.5238	0.6214	0.061*	0.555 (4)
O4	0.48981 (19)	0.29416 (13)	0.59801 (7)	0.0689 (4)	
H4B	0.4357	0.3423	0.5579	0.069*	0.445 (4)
O1	0.86755 (14)	0.43154 (15)	0.88375 (8)	0.0638 (4)	
O2	0.27767 (14)	0.31116 (13)	0.79491 (7)	0.0573 (3)	
N1	0.58016 (15)	0.35733 (14)	0.82137 (7)	0.0451 (3)	
C1	0.51315 (19)	0.36686 (16)	0.66427 (9)	0.0457 (3)	
C2	0.6117 (2)	0.29280 (17)	0.74458 (10)	0.0504 (4)	
H2A	0.5759	0.1928	0.7422	0.060*	
H2B	0.7383	0.2956	0.7476	0.060*	
C3	0.71166 (19)	0.42013 (16)	0.88520 (9)	0.0447 (3)	
C4	0.41265 (18)	0.35918 (15)	0.84052 (9)	0.0429 (3)	
C5	0.43776 (17)	0.43160 (14)	0.92473 (8)	0.0388 (3)	
C6	0.31300 (17)	0.46298 (15)	0.97165 (9)	0.0410 (3)	
H6	0.1933	0.4390	0.9531	0.049*	
C7	0.61865 (17)	0.46738 (15)	0.95191 (8)	0.0390 (3)	
N2	0.09136 (18)	0.56664 (15)	0.40440 (9)	0.0545 (4)	
C10	0.2173 (2)	0.5240 (2)	0.46976 (11)	0.0592 (4)	
H10A	0.2662	0.4330	0.4691	0.071*	0.555 (4)
H10B	0.2583	0.5849	0.5155	0.071*	0.445 (4)
C9	0.0213 (3)	0.4691 (2)	0.33408 (13)	0.0726 (5)	
H9A	0.0839	0.3793	0.3439	0.109*	
H9B	0.0370	0.5110	0.2827	0.109*	
H9C	-0.1031	0.4529	0.3295	0.109*	
C8	0.0168 (3)	0.7115 (2)	0.40126 (13)	0.0707 (5)	
H8A	0.0662	0.7600	0.4536	0.106*	
H8B	-0.1101	0.7056	0.3925	0.106*	
H8C	0.0454	0.7640	0.3557	0.106*	
O5A	0.2757 (3)	0.6081 (2)	0.53640 (13)	0.0644 (8)	0.555 (4)
O5B	0.2835 (4)	0.3945 (3)	0.46915 (17)	0.0705 (11)	0.445 (4)

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O3	0.0914 (9)	0.0416 (6)	0.0451 (6)	0.0103 (6)	0.0044 (6)	-0.0013 (5)
O4	0.0965 (10)	0.0517 (7)	0.0497 (7)	0.0094 (6)	0.0004 (6)	-0.0124 (5)
O1	0.0363 (6)	0.0933 (9)	0.0612 (7)	-0.0030 (6)	0.0105 (5)	-0.0050 (6)
O2	0.0464 (6)	0.0670 (7)	0.0513 (6)	-0.0117 (5)	-0.0021 (5)	-0.0040 (5)
N1	0.0416 (6)	0.0520 (7)	0.0393 (6)	-0.0001 (5)	0.0049 (5)	0.0040 (5)
C1	0.0469 (8)	0.0430 (8)	0.0452 (8)	-0.0043 (6)	0.0074 (6)	0.0013 (6)
C2	0.0553 (9)	0.0465 (8)	0.0478 (8)	0.0061 (7)	0.0096 (7)	0.0031 (6)
C3	0.0373 (7)	0.0499 (8)	0.0432 (7)	0.0010 (6)	0.0025 (6)	0.0091 (6)
C4	0.0407 (7)	0.0417 (7)	0.0413 (7)	-0.0020 (6)	0.0006 (6)	0.0092 (6)
C5	0.0345 (7)	0.0380 (7)	0.0391 (7)	-0.0020 (5)	-0.0007 (5)	0.0102 (5)
C6	0.0291 (6)	0.0451 (7)	0.0439 (7)	-0.0036 (5)	-0.0010 (5)	0.0088 (6)
C7	0.0321 (7)	0.0408 (7)	0.0409 (7)	0.0004 (5)	0.0027 (5)	0.0108 (6)
N2	0.0486 (7)	0.0539 (8)	0.0562 (8)	0.0005 (6)	0.0033 (6)	0.0000 (6)
C10	0.0557 (10)	0.0610 (10)	0.0573 (10)	0.0013 (8)	0.0069 (8)	0.0063 (8)
C9	0.0696 (12)	0.0693 (12)	0.0691 (12)	0.0050 (10)	-0.0019 (9)	-0.0119 (9)
C8	0.0690 (12)	0.0566 (10)	0.0766 (12)	0.0060 (9)	-0.0016 (9)	-0.0009 (9)
O5A	0.0737 (15)	0.0581 (13)	0.0527 (13)	0.0055 (11)	-0.0016 (10)	0.0063 (10)
O5B	0.081 (2)	0.069 (2)	0.0497 (16)	0.0120 (15)	-0.0070 (13)	0.0009 (13)

Geometric parameters (\AA , $^\circ$)

O3—C1	1.2621 (18)	C6—C7 ⁱ	1.393 (2)
O3—H3A	0.8200	C6—H6	0.9300
O4—C1	1.2519 (18)	C7—C6 ⁱ	1.393 (2)
O4—H4B	0.8200	N2—C10	1.321 (2)
O1—C3	1.2182 (18)	N2—C9	1.462 (2)
O2—C4	1.2128 (17)	N2—C8	1.463 (2)
N1—C3	1.3971 (19)	C10—O5B	1.311 (3)
N1—C4	1.4067 (19)	C10—O5A	1.328 (3)
N1—C2	1.464 (2)	C10—H10A	0.9300
C1—C2	1.515 (2)	C10—H10B	0.9300
C2—H2A	0.9700	C9—H9A	0.9600
C2—H2B	0.9700	C9—H9B	0.9600
C3—C7	1.510 (2)	C9—H9C	0.9600
C4—C5	1.502 (2)	C8—H8A	0.9600
C5—C6	1.402 (2)	C8—H8B	0.9600
C5—C7	1.4037 (19)	C8—H8C	0.9600
C1—O3—H3A	109.5	C6 ⁱ —C7—C5	121.85 (13)
C1—O4—H4B	109.5	C6 ⁱ —C7—C3	129.67 (12)
C3—N1—C4	111.97 (12)	C5—C7—C3	108.48 (12)
C3—N1—C2	124.62 (13)	C10—N2—C9	120.63 (15)
C4—N1—C2	123.38 (13)	C8—N2—C10	120.88 (15)
O4—C1—O3	125.38 (14)	C9—N2—C8	118.48 (14)
O4—C1—C2	116.07 (13)	O5B—C10—N2	118.93 (19)

O3—C1—C2	118.53 (13)	O5B—C10—O5A	119.5 (2)
N1—C2—C1	113.67 (12)	N2—C10—O5A	121.58 (18)
N1—C2—H2A	108.8	N2—C10—H10A	119.2
C1—C2—H2A	108.8	O5A—C10—H10A	119.2
N1—C2—H2B	108.8	O5B—C10—H10B	120.5
C1—C2—H2B	108.8	N2—C10—H10B	120.5
H2A—C2—H2B	107.7	H10A—C10—H10B	120.2
O1—C3—N1	124.69 (14)	N2—C9—H9A	109.5
O1—C3—C7	129.57 (14)	N2—C9—H9B	109.5
N1—C3—C7	105.74 (12)	H9A—C9—H9B	109.5
O2—C4—N1	124.30 (14)	N2—C9—H9C	109.5
O2—C4—C5	128.99 (14)	H9A—C9—H9C	109.5
N1—C4—C5	106.70 (11)	H9B—C9—H9C	109.5
C6—C5—C7	123.04 (13)	N2—C8—H8A	109.5
C6—C5—C4	129.85 (12)	N2—C8—H8B	109.5
C7—C5—C4	107.11 (13)	H8A—C8—H8B	109.5
C7 ⁱ —C6—C5	115.11 (12)	N2—C8—H8C	109.5
C7 ⁱ —C6—H6	122.4	H8A—C8—H8C	109.5
C5—C6—H6	122.4	H8B—C8—H8C	109.5
C3—N1—C2—C1	-117.78 (16)	N1—C4—C5—C7	-0.77 (14)
C4—N1—C2—C1	64.21 (19)	C7—C5—C6—C7 ⁱ	-0.1 (2)
O4—C1—C2—N1	-158.82 (14)	C4—C5—C6—C7 ⁱ	-179.65 (13)
O3—C1—C2—N1	22.9 (2)	C6—C5—C7—C6 ⁱ	0.1 (2)
C4—N1—C3—O1	179.90 (15)	C4—C5—C7—C6 ⁱ	179.75 (12)
C2—N1—C3—O1	1.7 (2)	C6—C5—C7—C3	-178.94 (12)
C4—N1—C3—C7	-0.18 (16)	C4—C5—C7—C3	0.67 (14)
C2—N1—C3—C7	-178.38 (12)	O1—C3—C7—C6 ⁱ	0.6 (3)
C3—N1—C4—O2	179.89 (14)	N1—C3—C7—C6 ⁱ	-179.32 (13)
C2—N1—C4—O2	-1.9 (2)	O1—C3—C7—C5	179.59 (15)
C3—N1—C4—C5	0.58 (15)	N1—C3—C7—C5	-0.33 (15)
C2—N1—C4—C5	178.81 (12)	C9—N2—C10—O5B	2.6 (3)
O2—C4—C5—C6	-0.5 (2)	C8—N2—C8—O5B	-178.4 (2)
N1—C4—C5—C6	178.80 (13)	C9—N2—C10—O5A	-176.1 (2)
O2—C4—C5—C7	179.96 (15)	C8—N2—C8—O5A	2.9 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O3—H3A \cdots O5A	0.82	1.72	2.526 (2)	166
O4—H4B \cdots O5B	0.82	1.70	2.496 (2)	162
C2—H2A \cdots O3 ⁱⁱ	0.97	2.41	3.230 (2)	142
C6—H6 \cdots O1 ⁱⁱⁱ	0.93	2.50	3.414 (2)	166
C9—H9A \cdots O2 ^{iv}	0.96	2.57	3.432 (3)	149

Symmetry codes: (ii) $-x+1, y-1/2, -z+3/2$; (iii) $x-1, y, z$; (iv) $x, -y+1/2, z-1/2$.

supplementary materials

Fig. 1

