

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(S)-1-Carboxy-2-(4-nitrophenyl)-ethanaminium bromide

Bo Wang

Ordered Matter Science Research Center, College of Chemistry and Chemical Engineering, Southeast University, Nanjing 210096, People's Republic of China
Correspondence e-mail: fudavid88@yahoo.com.cn


Received 19 August 2009; accepted 1 September 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.032; wR factor = 0.069; data-to-parameter ratio = 18.0.

In the crystal structure of the title compound, $\text{C}_9\text{H}_{11}\text{N}_2\text{O}_4^+\text{Br}^-$, the ethanaminium cations and Br^- anions are linked together by $\text{N}-\text{H}\cdots\text{Br}$ and $\text{O}-\text{H}\cdots\text{Br}$ hydrogen bonding. In the cation, the nitro group is twisted with respect to the benzene ring, making a dihedral angle of 21.43 (5)°.

Related literature

For amino acid derivatives as ligands for the construction of metal-organic frameworks, see: Fu *et al.* (2007).



Experimental

Crystal data

$\text{C}_9\text{H}_{11}\text{N}_2\text{O}_4^+\text{Br}^-$
 $M_r = 291.11$
Monoclinic, $P2_1$
 $a = 5.5378$ (11) Å

$b = 7.4158$ (15) Å
 $c = 14.246$ (3) Å
 $\beta = 91.15$ (3)°
 $V = 584.9$ (2) Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 3.52$ mm⁻¹

$T = 298$ K
 $0.40 \times 0.05 \times 0.05$ mm

Data collection

Rigaku Mercury2 diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku, 2005)
 $T_{\min} = 0.76$, $T_{\max} = 0.84$

5994 measured reflections
2633 independent reflections
2427 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.069$
 $S = 1.04$
2633 reflections
146 parameters
1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.26$ e Å⁻³
 $\Delta\rho_{\min} = -0.31$ e Å⁻³
Absolute structure: Flack (1983),
1202 Friedel pairs
Flack parameter: -0.025 (11)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{Br1}^i$	0.89	2.54	3.355 (3)	153
$\text{N1}-\text{H1B}\cdots\text{Br1}$	0.89	2.46	3.340 (2)	168
$\text{N1}-\text{H1C}\cdots\text{Br1}^{ii}$	0.89	2.59	3.440 (3)	161
$\text{O1}-\text{H1}\cdots\text{Br1}^{iii}$	0.85	2.38	3.174 (3)	155

Symmetry codes: (i) $-x + 1, y - \frac{1}{2}, -z$; (ii) $x - 1, y, z$; (iii) $x - 1, y - 1, z$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXTL/PC* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL/PC*; molecular graphics: *SHELXTL/PC*; software used to prepare material for publication: *SHELXTL/PC*.

This work was supported by a start-up grant from Southeast University for Professor Ren-Gen Xiong.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2595).

References

- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Fu, D.-W., Song, Y.-M., Wang, G.-X., Ye, Q., Xiong, R.-G., Akutagawa, T., Nakamura, T., Chan, P. W. H. & Huang, S. P. D. (2007). *J. Am. Chem. Soc.* **129**, 5346–5347.
Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o2420 [doi:10.1107/S1600536809035211]

(S)-1-Carboxy-2-(4-nitrophenyl)ethanaminium bromide

B. Wang

Comment

Amino acid derivatives are a class of excellent ligands for the construction of novel metal-organic frameworks (Fu *et al.*, 2007). We report here the crystal structure of the title compound.

The title compound is built up from a Br⁻ anion and a protonated amino group cation (Fig. 1). The nitro group is twisted from the benzene ring plane by a dihedral angle of 21.43 (5)°, and the 2-aminopropanoate substituent group is a zig-zag chain.

The crystal packing is stabilized by cation-anion N—H...Br and O—H...Br H-bonds building an infinite two-dimensional network developing parallel to the (1 1 0) plane (Table 1).

Experimental

A mixture of 2-amino-3-phenylpropanoic acid (4.71 g, 30 mmol), concentrated nitric acid (4.0 ml, 14 M) and concentrated sulfuric acid (1.5 ml, 18 M) was stirred at 383 K for 3 h under nitrogen atmosphere. The resulting solution was poured into ice water (100 ml), then filtered and washed with distilled water. The crude product was recrystallized with distilled water by adding dilute HBr (4 ml, 4 M) to yield colorless needle-like single crystals.

Refinement

H atoms were positioned geometrically and treated as riding with C—H = 0.93 (aromatic), 0.97 (methylene), 0.98 Å (methine) and N—H = 0.89 Å, O—H = 0.85 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O,N})$.

Figures




Fig. 1. The structure of the title compound with the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

(S)-1-Carboxy-2-(4-nitrophenyl)ethanaminium bromide

Crystal data

C₉H₁₁N₂O₄⁺·Br⁻

$M_r = 291.11$

$F_{000} = 292$

$D_x = 1.653 \text{ Mg m}^{-3}$

supplementary materials

Monoclinic, $P2_1$
Hall symbol: P 2yb
 $a = 5.5378$ (11) Å
 $b = 7.4158$ (15) Å
 $c = 14.246$ (3) Å
 $\beta = 91.15$ (3)°
 $V = 584.9$ (2) Å³
 $Z = 2$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2427 reflections
 $\theta = 3.1$ – 27.4 °
 $\mu = 3.52$ mm⁻¹
 $T = 298$ K
Needle, colourless
 $0.40 \times 0.05 \times 0.05$ mm

Data collection

Rigaku Mercury2
diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
Detector resolution: 13.6612 pixels mm⁻¹
 $T = 298$ K
CCD profile fitting scans
Absorption correction: multi-scan
(CrystalClear; Rigaku, 2005)
 $T_{\min} = 0.76$, $T_{\max} = 0.84$
5994 measured reflections

2633 independent reflections
2427 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$
 $\theta_{\text{max}} = 27.4$ °
 $\theta_{\text{min}} = 3.1$ °
 $h = -7 \rightarrow 7$
 $k = -9 \rightarrow 9$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.069$
 $S = 1.04$
2633 reflections
146 parameters
1 restraint
Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.02P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³
Extinction correction: none
Absolute structure: Flack (1983), 1202 Friedel pairs
Flack parameter: -0.025 (11)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculat-

ing R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O2	-0.1236 (3)	0.1957 (5)	0.09374 (15)	0.0526 (5)
C7	0.3705 (6)	0.3156 (4)	0.2428 (2)	0.0396 (7)
H7A	0.3926	0.2030	0.2766	0.048*
H7B	0.5266	0.3746	0.2405	0.048*
C9	0.0598 (5)	0.1544 (4)	0.1348 (2)	0.0365 (8)
C8	0.2848 (6)	0.2725 (4)	0.1416 (2)	0.0346 (7)
H8	0.4158	0.2079	0.1109	0.041*
C6	0.2007 (5)	0.4342 (4)	0.29749 (19)	0.0361 (7)
C2	0.0984 (6)	0.7325 (5)	0.3539 (2)	0.0488 (10)
H2	0.1295	0.8556	0.3573	0.059*
C4	-0.1471 (7)	0.4776 (6)	0.3932 (2)	0.0488 (10)
H4	-0.2795	0.4309	0.4242	0.059*
C3	-0.0974 (6)	0.6591 (5)	0.3970 (2)	0.0474 (10)
C5	0.0006 (6)	0.3660 (4)	0.3434 (2)	0.0418 (8)
H5	-0.0333	0.2433	0.3402	0.050*
C1	0.2497 (6)	0.6173 (5)	0.3048 (2)	0.0427 (8)
H1D	0.3860	0.6640	0.2765	0.051*
O4	-0.3908 (5)	0.7160 (9)	0.5059 (2)	0.1119 (15)
N2	-0.2621 (7)	0.7828 (7)	0.4470 (3)	0.0783 (12)
O3	-0.2547 (8)	0.9417 (7)	0.4254 (4)	0.1225 (16)
N1	0.2332 (4)	0.4392 (4)	0.08589 (18)	0.0375 (6)
H1A	0.2284	0.4121	0.0250	0.056*
H1B	0.3490	0.5200	0.0971	0.056*
H1C	0.0915	0.4846	0.1024	0.056*
O1	0.0979 (5)	-0.0009 (3)	0.1777 (2)	0.0697 (9)
H1	-0.0290	-0.0651	0.1733	0.104*
Br1	0.72355 (5)	0.69798 (6)	0.112290 (18)	0.04362 (11)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O2	0.0343 (11)	0.0513 (12)	0.0719 (14)	-0.0139 (17)	-0.0077 (10)	0.010 (2)
C7	0.0327 (17)	0.0424 (18)	0.0434 (19)	-0.0014 (14)	-0.0097 (13)	-0.0022 (14)
C9	0.0373 (16)	0.032 (2)	0.0404 (16)	-0.0075 (12)	-0.0008 (13)	-0.0020 (12)
C8	0.0312 (16)	0.0363 (16)	0.0361 (16)	-0.0054 (12)	-0.0016 (12)	-0.0049 (13)
C6	0.0351 (16)	0.0460 (18)	0.0267 (15)	-0.0042 (14)	-0.0064 (12)	0.0013 (13)
C2	0.060 (2)	0.047 (3)	0.0399 (17)	-0.0022 (17)	-0.0021 (16)	-0.0023 (15)
C4	0.0346 (18)	0.076 (3)	0.036 (2)	-0.0130 (19)	0.0044 (14)	-0.0054 (17)
C3	0.0413 (19)	0.065 (3)	0.0356 (17)	0.0010 (17)	-0.0043 (13)	-0.0170 (16)
C5	0.0438 (19)	0.0447 (19)	0.0368 (18)	-0.0144 (14)	-0.0050 (15)	0.0010 (14)
C1	0.045 (2)	0.052 (2)	0.0310 (17)	-0.0108 (15)	0.0004 (14)	-0.0024 (14)
O4	0.074 (2)	0.156 (4)	0.108 (2)	-0.038 (3)	0.0436 (19)	-0.074 (3)

supplementary materials

N2	0.064 (3)	0.099 (3)	0.072 (3)	-0.001 (2)	0.002 (2)	-0.046 (2)
O3	0.123 (4)	0.095 (3)	0.150 (4)	0.034 (3)	0.027 (3)	-0.037 (3)
N1	0.0355 (14)	0.0441 (16)	0.0330 (13)	-0.0151 (12)	0.0001 (10)	-0.0006 (12)
O1	0.072 (2)	0.0427 (15)	0.092 (2)	-0.0215 (13)	-0.0365 (17)	0.0197 (14)
Br1	0.04125 (17)	0.04304 (17)	0.04656 (17)	-0.01590 (17)	0.00035 (11)	0.00473 (18)

Geometric parameters (Å, °)

O2—C9	1.201 (3)	C2—H2	0.9300
C7—C6	1.515 (4)	C4—C5	1.371 (5)
C7—C8	1.543 (4)	C4—C3	1.375 (6)
C7—H7A	0.9700	C4—H4	0.9300
C7—H7B	0.9700	C3—N2	1.486 (5)
C9—O1	1.318 (4)	C5—H5	0.9300
C9—C8	1.525 (4)	C1—H1D	0.9300
C8—N1	1.493 (4)	O4—N2	1.217 (5)
C8—H8	0.9800	N2—O3	1.219 (6)
C6—C1	1.388 (5)	N1—H1A	0.8900
C6—C5	1.393 (4)	N1—H1B	0.8900
C2—C3	1.369 (5)	N1—H1C	0.8900
C2—C1	1.395 (5)	O1—H1	0.8500
C6—C7—C8	114.7 (3)	C5—C4—H4	120.3
C6—C7—H7A	108.6	C3—C4—H4	120.3
C8—C7—H7A	108.6	C2—C3—C4	122.1 (3)
C6—C7—H7B	108.6	C2—C3—N2	117.9 (4)
C8—C7—H7B	108.6	C4—C3—N2	119.9 (4)
H7A—C7—H7B	107.6	C4—C5—C6	120.8 (3)
O2—C9—O1	125.0 (3)	C4—C5—H5	119.6
O2—C9—C8	124.5 (3)	C6—C5—H5	119.6
O1—C9—C8	110.5 (3)	C6—C1—C2	121.2 (3)
N1—C8—C9	107.1 (2)	C6—C1—H1D	119.4
N1—C8—C7	112.2 (2)	C2—C1—H1D	119.4
C9—C8—C7	114.4 (3)	O4—N2—O3	126.2 (5)
N1—C8—H8	107.6	O4—N2—C3	117.0 (5)
C9—C8—H8	107.6	O3—N2—C3	116.8 (4)
C7—C8—H8	107.6	C8—N1—H1A	109.5
C1—C6—C5	118.4 (3)	C8—N1—H1B	109.5
C1—C6—C7	119.0 (3)	H1A—N1—H1B	109.5
C5—C6—C7	122.6 (3)	C8—N1—H1C	109.5
C3—C2—C1	118.0 (3)	H1A—N1—H1C	109.5
C3—C2—H2	121.0	H1B—N1—H1C	109.5
C1—C2—H2	121.0	C9—O1—H1	109.3
C5—C4—C3	119.3 (3)		
O2—C9—C8—N1	-1.9 (4)	C5—C4—C3—N2	177.2 (3)
O1—C9—C8—N1	176.0 (3)	C3—C4—C5—C6	0.6 (5)
O2—C9—C8—C7	123.1 (3)	C1—C6—C5—C4	1.2 (5)
O1—C9—C8—C7	-58.9 (4)	C7—C6—C5—C4	179.1 (3)
C6—C7—C8—N1	55.2 (4)	C5—C6—C1—C2	-2.3 (5)
C6—C7—C8—C9	-67.1 (3)	C7—C6—C1—C2	179.6 (3)

C8—C7—C6—C1	-99.1 (3)	C3—C2—C1—C6	1.7 (5)
C8—C7—C6—C5	82.9 (4)	C2—C3—N2—O4	-159.6 (4)
C1—C2—C3—C4	0.2 (5)	C4—C3—N2—O4	21.8 (5)
C1—C2—C3—N2	-178.4 (3)	C2—C3—N2—O3	20.1 (6)
C5—C4—C3—C2	-1.3 (5)	C4—C3—N2—O3	-158.4 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1A...Br1 ⁱ	0.89	2.54	3.355 (3)	153
N1—H1B...Br1	0.89	2.46	3.340 (2)	168
N1—H1C...Br1 ⁱⁱ	0.89	2.59	3.440 (3)	161
O1—H1...Br1 ⁱⁱⁱ	0.85	2.38	3.174 (3)	155

Symmetry codes: (i) $-x+1, y-1/2, -z$; (ii) $x-1, y, z$; (iii) $x-1, y-1, z$.

Fig. 1

