

Triphenyl[(*E*)-4-[4-(phenyldiazenyl)-phenyl]-4*H*-1,2,4-triazol-1-yl]boron

Daisuke Urakami,^a Katsuya Inoue^{a,b} and Shinya Hayami^{c*}

^aDepartment of Chemistry, Graduate School of Science, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan, ^bInstitute for Advanced Materials Research, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima 739-8526, Japan, and ^cDepartment of Chemistry, Graduate School of Science and Technology, Kumamoto University, 2-39-1 Kurokami, Kumamoto 860-8555, Japan
Correspondence e-mail: hayami@sci.kumamoto-u.ac.jp


Received 25 July 2009; accepted 7 September 2009

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.050; wR factor = 0.138; data-to-parameter ratio = 16.5.

In the title compound, $\text{C}_{32}\text{H}_{26}\text{BN}_5$ or $[(\text{C}_{14}\text{H}_{11}\text{N}_5)\text{B}(\text{C}_6\text{H}_5)_3]$, the B atom is approximately tetrahedrally coordinated. The diazo unit is in a *trans* conformation, which is generally more stable than a *cis* one for aromatic azo compounds. The crystal structure features very weak $\text{C}-\text{H}\cdots\pi$ interactions. The dihedral angles between the central benzene ring and the terminal rings in the heterocycle are 62.64, 73.54 and 61.60°.

Related literature

For the use of azobenzenes for holographic information storage, see: Rasmussen *et al.* (1999).



Experimental

Crystal data

$\text{C}_{32}\text{H}_{26}\text{BN}_5$
 $M_r = 491.39$
Monoclinic, $P2_1/n$

$a = 17.4049$ (6) Å
 $b = 7.1284$ (2) Å
 $c = 21.0645$ (7) Å

$\beta = 97.0810$ (15)°
 $V = 2593.52$ (14) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.08$ mm⁻¹
 $T = 113$ K
 $0.40 \times 0.20 \times 0.10$ mm

Data collection

Rigaku R-Axis RAPID diffractometer
Absorption correction: multi-scan (Higashi, 2001)
 $T_{\min} = 0.971$, $T_{\max} = 0.993$

19971 measured reflections
5668 independent reflections
4068 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.138$
 $S = 1.03$
5668 reflections

343 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.26$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C4}-\text{H3}\cdots\text{Cg1}^i$	0.95	2.91	3.822 (2)	162

Symmetry code: (i) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$. Cg1 is the centroid of the C27-C31 ring.

Data collection: *PROCESS-AUTO* (Rigaku, 1998); cell refinement: *PROCESS-AUTO*; data reduction: *CrystalClear* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Yadokari-XG* (Wakita, 2000); software used to prepare material for publication: *SHELXL97*.

This work was supported by the project 'Development of Molecular Devices in Ferroelectric Metallomesogens' in 2006 of the New Energy and Industrial Technology Development Organization (NEDO) of Japan and by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of the Japanese Government (No. 20350028).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BX2228).

References

- Higashi, T. (2001). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
Rasmussen, P. H., Ramanujam, P. S., Hvilsted, S. & Berg, R. H. (1999). *J. Am. Chem. Soc.* **121**, 4738–4743.
Rigaku (1998). *PROCESS-AUTO*. Rigaku Corporation, Tokyo, Japan.
Rigaku/MSK (2002). *CrystalClear*. Rigaku/MSK, The Woodlands, Texas, USA, and Rigaku Corporation, Tokyo, Japan.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Wakita, K. (2000). *Yadokari-XG*. Department of Chemistry, Graduate School of Science, University of Tokyo, Japan.

supplementary materials

Acta Cryst. (2009). E65, o2425 [doi:10.1107/S1600536809036071]

Triphenyl{(E)-4-[4-(phenyldiazenyl)phenyl]-4H-1,2,4-triazol-1-yl}boron

D. Urakami, K. Inoue and S. Hayami

Comment

Photochromic materials are of interest as media for recording, storage, and reproduction of information in nonlinear optics and holography. Aromatic azo compounds occupy an important place among such materials, therefore many photoisomerizable azobenzene groups were synthesized to apply for photoswitchable systems or optical data storage device (Rasmussen *et al.*, 1999). Here we report the synthesis and crystal structure of the title complex. The addition compound of a triazole derivative with diazobenzene and a triphenylboron contains a B atom approximately tetrahedrally coordinated with a B—N distance of 1.636 (2) Å and B—C bond lengths averaging 1.625 (2) Å. The diazo moiety forms a *trans* conformation, and the bond distance N4—N5 is 1.2563 (19) Å. The crystal structure is stabilized by weak C—H \cdots π (C27/C32) interactions between phenyl hydrogens from (*E*)-4-(4-(phenyldiazenyl)phenyl)-4H-1,2,4-triazole fragment to phenyl rings from triphenylboron fragment. Fig 2, Table 1.

Experimental

The compound (I) was synthesized by refluxed for an hour with (*E*)-4-(4-(phenyldiazenyl)phenyl)-4H-1,2,4-triazole (3 mmol, 0.748 g), sodium tetraphenylborate (4 mmol, 1.369 g) and iron chloride tetrahydrate (2 mmol, 0.390 g) in ethanol (50 ml). The resulting yellow solution was filtered and the single crystals of (I) suitable X-ray diffraction were obtained by slowly evaporation of ethanol.

Refinement

All H atoms were positioned geometrically (C—H = 0.95 Å) and were refined as riding, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. ORTEP drawing for the compound (I) showing 50% probability displacement ellipsoids.




Fig. 2. Packing structure for the compound (I).

Triphenyl[(*E*)-4-[4-(phenyldiazenyl)phenyl]-4*H*-1,2,4-triazol-1-yl]boron

Crystal data

$C_{32}H_{26}BN_5$	$F_{000} = 1032$
$M_r = 491.39$	$D_x = 1.258 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 13301 reflections
$a = 17.4049 (6) \text{ \AA}$	$\theta = 2.7\text{--}27.4^\circ$
$b = 7.1284 (2) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 21.0645 (7) \text{ \AA}$	$T = 113 \text{ K}$
$\beta = 97.0810 (15)^\circ$	Block, orange
$V = 2593.52 (14) \text{ \AA}^3$	$0.40 \times 0.20 \times 0.10 \text{ mm}$
$Z = 4$	

Data collection

Rigaku R-Axis RAPID diffractometer	5668 independent reflections
Radiation source: fine-focus sealed tube	4068 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.050$
$T = 113 \text{ K}$	$\theta_{\text{max}} = 27.4^\circ$
ω scans	$\theta_{\text{min}} = 1.4^\circ$
Absorption correction: multi-scan (Higashi, 2001)	$h = -21 \rightarrow 22$
$T_{\text{min}} = 0.971$, $T_{\text{max}} = 0.993$	$k = -9 \rightarrow 8$
19971 measured reflections	$l = -27 \rightarrow 27$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.050$	H-atom parameters constrained
$wR(F^2) = 0.138$	$w = 1/[\sigma^2(F_o^2) + (0.0835P)^2]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
5668 reflections	$(\Delta/\sigma)_{\text{max}} = 0.002$
343 parameters	$\Delta\rho_{\text{max}} = 0.26 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.27 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
B1	0.21836 (11)	0.3083 (3)	0.04914 (9)	0.0190 (4)
C1	0.29403 (9)	0.0119 (2)	0.10360 (8)	0.0197 (4)
H1	0.2800	-0.0758	0.0702	0.024*
C2	0.34594 (11)	0.1427 (2)	0.19029 (8)	0.0285 (4)
H2	0.3765	0.1618	0.2304	0.034*
C3	0.38777 (9)	-0.1882 (2)	0.17537 (8)	0.0201 (4)
C4	0.41059 (10)	-0.2207 (2)	0.23988 (8)	0.0256 (4)
H3	0.3942	-0.1399	0.2714	0.031*
C5	0.45769 (11)	-0.3730 (2)	0.25743 (8)	0.0273 (4)
H4	0.4750	-0.3949	0.3014	0.033*
C6	0.48000 (9)	-0.4945 (2)	0.21125 (8)	0.0214 (4)
C7	0.45478 (10)	-0.4616 (2)	0.14683 (8)	0.0219 (4)
H5	0.4687	-0.5460	0.1153	0.026*
C8	0.40960 (9)	-0.3068 (2)	0.12870 (8)	0.0217 (4)
H6	0.3937	-0.2820	0.0847	0.026*
C9	0.58832 (10)	-0.9199 (2)	0.21814 (8)	0.0225 (4)
C10	0.62167 (10)	-0.9365 (3)	0.28142 (8)	0.0247 (4)
H7	0.6102	-0.8473	0.3124	0.030*
C11	0.67159 (10)	-1.0832 (3)	0.29885 (9)	0.0296 (4)
H8	0.6944	-1.0950	0.3420	0.036*
C12	0.68874 (11)	-1.2142 (3)	0.25356 (9)	0.0330 (5)
H9	0.7238	-1.3137	0.2657	0.040*
C13	0.65460 (12)	-1.1989 (3)	0.19090 (10)	0.0353 (5)
H10	0.6659	-1.2888	0.1600	0.042*
C14	0.60408 (11)	-1.0529 (3)	0.17306 (9)	0.0313 (4)
H11	0.5802	-1.0435	0.1301	0.038*
C15	0.17458 (9)	0.1567 (2)	-0.00001 (8)	0.0194 (4)
C16	0.12558 (9)	0.0210 (2)	0.02111 (8)	0.0222 (4)
H12	0.1184	0.0189	0.0651	0.027*
C17	0.08711 (10)	-0.1107 (2)	-0.01958 (9)	0.0267 (4)
H13	0.0547	-0.2015	-0.0033	0.032*
C18	0.09617 (11)	-0.1094 (2)	-0.08390 (9)	0.0292 (4)
H14	0.0702	-0.1994	-0.1121	0.035*
C19	0.14345 (10)	0.0242 (3)	-0.10691 (8)	0.0286 (4)
H15	0.1495	0.0268	-0.1511	0.034*
C20	0.18194 (10)	0.1542 (2)	-0.06551 (8)	0.0230 (4)
H16	0.2143	0.2445	-0.0821	0.028*
C21	0.28127 (9)	0.4280 (2)	0.01512 (7)	0.0190 (4)

supplementary materials

C22	0.35283 (10)	0.3533 (2)	0.00407 (8)	0.0233 (4)
H17	0.3672	0.2329	0.0208	0.028*
C23	0.40343 (10)	0.4484 (3)	-0.03036 (8)	0.0266 (4)
H18	0.4515	0.3931	-0.0367	0.032*
C24	0.38387 (10)	0.6248 (3)	-0.05563 (8)	0.0282 (4)
H19	0.4185	0.6911	-0.0790	0.034*
C25	0.31309 (10)	0.7026 (3)	-0.04624 (8)	0.0258 (4)
H20	0.2988	0.8223	-0.0636	0.031*
C26	0.26302 (10)	0.6055 (2)	-0.01144 (8)	0.0219 (4)
H21	0.2149	0.6612	-0.0054	0.026*
C27	0.15982 (10)	0.4430 (2)	0.08339 (8)	0.0195 (4)
C28	0.18757 (10)	0.5898 (2)	0.12474 (8)	0.0222 (4)
H22	0.2419	0.6043	0.1354	0.027*
C29	0.13847 (11)	0.7138 (2)	0.15035 (8)	0.0256 (4)
H23	0.1594	0.8127	0.1774	0.031*
C30	0.05859 (11)	0.6942 (3)	0.13665 (8)	0.0277 (4)
H24	0.0247	0.7785	0.1543	0.033*
C31	0.02944 (10)	0.5507 (3)	0.09706 (9)	0.0275 (4)
H25	-0.0250	0.5348	0.0876	0.033*
C32	0.07955 (10)	0.4284 (2)	0.07081 (8)	0.0241 (4)
H26	0.0581	0.3313	0.0432	0.029*
N1	0.26879 (8)	0.1851 (2)	0.10506 (6)	0.0194 (3)
N2	0.30188 (9)	0.2706 (2)	0.16050 (7)	0.0285 (4)
N3	0.34303 (8)	-0.02185 (19)	0.15713 (6)	0.0204 (3)
N4	0.52992 (8)	-0.6447 (2)	0.23468 (7)	0.0238 (3)
N5	0.53680 (8)	-0.7720 (2)	0.19453 (7)	0.0245 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
B1	0.0204 (9)	0.0185 (9)	0.0176 (9)	0.0035 (8)	0.0006 (7)	0.0034 (8)
C1	0.0226 (8)	0.0201 (8)	0.0159 (8)	0.0007 (8)	0.0001 (6)	0.0013 (7)
C2	0.0378 (10)	0.0253 (10)	0.0205 (9)	0.0030 (9)	-0.0045 (8)	-0.0015 (7)
C3	0.0211 (8)	0.0176 (8)	0.0210 (8)	0.0019 (7)	0.0000 (7)	0.0034 (7)
C4	0.0351 (10)	0.0244 (9)	0.0166 (8)	0.0058 (8)	0.0005 (7)	-0.0011 (7)
C5	0.0369 (10)	0.0268 (10)	0.0165 (9)	0.0035 (8)	-0.0036 (7)	0.0015 (7)
C6	0.0216 (8)	0.0229 (9)	0.0190 (8)	0.0015 (8)	-0.0002 (7)	0.0034 (7)
C7	0.0240 (8)	0.0227 (9)	0.0189 (8)	0.0028 (8)	0.0030 (7)	-0.0001 (7)
C8	0.0227 (8)	0.0266 (9)	0.0151 (8)	0.0009 (8)	-0.0005 (6)	0.0020 (7)
C9	0.0232 (9)	0.0229 (9)	0.0222 (9)	0.0031 (8)	0.0057 (7)	0.0046 (7)
C10	0.0266 (9)	0.0245 (9)	0.0235 (9)	0.0028 (8)	0.0050 (7)	0.0029 (7)
C11	0.0296 (10)	0.0304 (10)	0.0285 (10)	0.0027 (9)	0.0023 (8)	0.0071 (8)
C12	0.0319 (10)	0.0275 (10)	0.0409 (12)	0.0094 (9)	0.0102 (9)	0.0093 (9)
C13	0.0449 (12)	0.0287 (10)	0.0347 (11)	0.0094 (10)	0.0140 (9)	0.0002 (9)
C14	0.0403 (11)	0.0302 (10)	0.0234 (9)	0.0054 (9)	0.0041 (8)	0.0009 (8)
C15	0.0192 (8)	0.0190 (8)	0.0196 (8)	0.0061 (7)	0.0001 (6)	0.0014 (7)
C16	0.0214 (9)	0.0234 (9)	0.0215 (9)	0.0041 (8)	0.0016 (7)	0.0018 (7)
C17	0.0224 (9)	0.0216 (9)	0.0343 (10)	0.0007 (8)	-0.0031 (7)	0.0026 (8)

C18	0.0296 (10)	0.0219 (9)	0.0325 (10)	0.0030 (8)	-0.0102 (8)	-0.0068 (8)
C19	0.0350 (10)	0.0300 (10)	0.0190 (9)	0.0061 (9)	-0.0031 (7)	-0.0029 (7)
C20	0.0263 (9)	0.0220 (9)	0.0203 (9)	0.0025 (8)	0.0009 (7)	0.0001 (7)
C21	0.0223 (8)	0.0209 (8)	0.0129 (8)	-0.0009 (7)	-0.0008 (6)	-0.0022 (6)
C22	0.0241 (9)	0.0234 (9)	0.0225 (9)	0.0027 (8)	0.0029 (7)	0.0009 (7)
C23	0.0237 (9)	0.0305 (10)	0.0266 (9)	0.0036 (8)	0.0071 (7)	0.0002 (8)
C24	0.0284 (9)	0.0334 (10)	0.0238 (9)	-0.0030 (9)	0.0081 (7)	0.0029 (8)
C25	0.0303 (9)	0.0259 (9)	0.0205 (9)	0.0009 (8)	0.0008 (7)	0.0047 (7)
C26	0.0224 (8)	0.0251 (9)	0.0181 (8)	0.0030 (8)	0.0017 (7)	-0.0004 (7)
C27	0.0250 (9)	0.0181 (8)	0.0158 (8)	0.0023 (8)	0.0041 (6)	0.0037 (7)
C28	0.0248 (9)	0.0217 (9)	0.0205 (9)	0.0030 (8)	0.0044 (7)	0.0025 (7)
C29	0.0379 (10)	0.0204 (9)	0.0194 (9)	-0.0003 (8)	0.0076 (7)	-0.0004 (7)
C30	0.0345 (10)	0.0270 (9)	0.0238 (9)	0.0088 (9)	0.0125 (8)	0.0012 (8)
C31	0.0239 (9)	0.0298 (10)	0.0295 (10)	0.0021 (8)	0.0059 (7)	0.0024 (8)
C32	0.0260 (9)	0.0240 (9)	0.0225 (9)	0.0014 (8)	0.0032 (7)	-0.0008 (7)
N1	0.0217 (7)	0.0208 (7)	0.0153 (7)	-0.0017 (6)	0.0006 (5)	-0.0010 (6)
N2	0.0382 (9)	0.0267 (8)	0.0182 (7)	0.0030 (7)	-0.0063 (6)	-0.0037 (6)
N3	0.0231 (7)	0.0211 (7)	0.0165 (7)	0.0009 (6)	0.0003 (6)	0.0000 (6)
N4	0.0265 (8)	0.0247 (8)	0.0201 (7)	0.0031 (7)	0.0015 (6)	0.0025 (6)
N5	0.0269 (8)	0.0261 (8)	0.0203 (7)	0.0038 (7)	0.0023 (6)	0.0023 (6)

Geometric parameters (Å, °)

B1—C15	1.621 (2)	C15—C20	1.401 (2)
B1—C21	1.623 (2)	C16—C17	1.387 (2)
B1—C27	1.631 (2)	C16—H12	0.9500
B1—N1	1.636 (2)	C17—C18	1.383 (3)
C1—N1	1.312 (2)	C17—H13	0.9500
C1—N3	1.349 (2)	C18—C19	1.385 (3)
C1—H1	0.9500	C18—H14	0.9500
C2—N2	1.302 (2)	C19—C20	1.387 (2)
C2—N3	1.363 (2)	C19—H15	0.9500
C2—H2	0.9500	C20—H16	0.9500
C3—C4	1.388 (2)	C21—C22	1.400 (2)
C3—C8	1.385 (2)	C21—C26	1.404 (2)
C3—N3	1.444 (2)	C22—H17	0.9500
C4—C5	1.383 (2)	C22—C23	1.385 (2)
C4—H3	0.9500	C23—C24	1.391 (2)
C5—C6	1.393 (2)	C23—H18	0.9500
C5—H4	0.9500	C24—C25	1.387 (2)
C6—C7	1.394 (2)	C24—H19	0.9500
C6—N4	1.428 (2)	C25—C26	1.390 (2)
C7—H5	0.9500	C25—H20	0.9500
C7—C8	1.381 (2)	C26—H21	0.9500
C8—H6	0.9500	C27—C28	1.408 (2)
C9—C10	1.392 (2)	C27—C32	1.393 (2)
C9—C14	1.393 (2)	C28—C29	1.384 (2)
C9—N5	1.433 (2)	C28—H22	0.9500
C10—C11	1.380 (2)	C29—C30	1.392 (3)

supplementary materials

C10—H7	0.9500	C29—H23	0.9500
C11—C12	1.393 (3)	C30—C31	1.377 (3)
C11—H8	0.9500	C30—H24	0.9500
C12—C13	1.383 (3)	C31—C32	1.394 (2)
C12—H9	0.9500	C31—H25	0.9500
C13—C14	1.384 (3)	C32—H26	0.9500
C13—H10	0.9500	N1—N2	1.3787 (18)
C14—H11	0.9500	N4—N5	1.2563 (19)
C15—C16	1.398 (2)		
B1—C15—C16	121.02 (14)	C15—C20—C19	122.11 (17)
B1—C15—C20	122.99 (15)	C15—C20—H16	118.9
B1—C21—C22	122.41 (15)	C16—C15—C20	115.98 (15)
B1—C21—C26	121.19 (14)	C16—C17—C18	119.75 (17)
B1—C27—C32	122.56 (15)	C16—C17—H13	120.1
B1—C27—C28	121.70 (15)	C17—C16—H12	118.7
B1—N1—C1	129.79 (13)	C17—C18—C19	119.48 (16)
B1—N1—N2	120.24 (13)	C17—C18—H14	120.3
C1—N1—N2	109.25 (13)	C18—C17—H13	120.1
C1—N3—C2	104.76 (14)	C18—C19—C20	120.08 (17)
C1—N3—C3	129.28 (14)	C18—C19—H15	120.0
C2—N2—N1	104.99 (14)	C19—C18—H14	120.3
C2—N3—C3	125.84 (14)	C19—C20—H16	118.9
C3—C4—C5	118.81 (16)	C20—C19—H15	120.0
C3—C4—H3	120.6	C21—B1—C27	112.21 (14)
C3—C8—C7	119.23 (15)	C21—B1—N1	105.60 (12)
C3—C8—H6	120.4	C21—C22—C23	122.48 (16)
C4—C3—C8	121.50 (16)	C21—C22—H17	118.8
C4—C3—N3	118.53 (15)	C21—C26—C25	122.20 (16)
C4—C5—C6	120.56 (16)	C21—C26—H21	118.9
C4—C5—H4	119.7	C22—C21—C26	116.02 (15)
C5—C4—H3	120.6	C22—C23—C24	120.08 (16)
C5—C6—C7	119.60 (16)	C22—C23—H18	120.0
C5—C6—N4	115.79 (14)	C23—C22—H17	118.8
C6—C5—H4	119.7	C23—C24—C25	119.10 (16)
C6—C7—C8	120.25 (16)	C23—C24—H19	120.4
C6—C7—H5	119.9	C24—C23—H18	120.0
C6—N4—N5	114.42 (14)	C24—C25—C26	120.12 (16)
C7—C6—N4	124.60 (15)	C24—C25—H20	119.9
C7—C8—H6	120.4	C25—C24—H19	120.4
C8—C3—N3	119.91 (14)	C25—C26—H21	118.9
C8—C7—H5	119.9	C26—C25—H20	119.9
C9—C10—C11	119.59 (17)	C27—B1—N1	107.85 (13)
C9—C10—H7	120.2	C27—C28—C29	122.30 (16)
C9—C14—C13	119.91 (17)	C27—C28—H22	118.9
C9—C14—H11	120.0	C27—C32—C31	122.65 (17)
C9—N5—N4	113.92 (14)	C27—C32—H26	118.7
C10—C9—C14	120.07 (16)	C28—C27—C32	115.62 (15)
C10—C9—N5	124.35 (16)	C28—C29—C30	120.25 (17)
C10—C11—C12	120.43 (18)	C28—C29—H23	119.9

C10—C11—H8	119.8	C29—C28—H22	118.9
C11—C10—H7	120.2	C29—C30—C31	119.00 (16)
C11—C12—C13	119.84 (18)	C29—C30—H24	120.5
C11—C12—H9	120.1	C30—C29—H23	119.9
C12—C11—H8	119.8	C30—C31—C32	120.17 (17)
C12—C13—C14	120.13 (18)	C30—C31—H25	119.9
C12—C13—H10	119.9	C31—C30—H24	120.5
C13—C12—H9	120.1	C31—C32—H26	118.7
C13—C14—H11	120.0	C32—C31—H25	119.9
C14—C9—N5	115.58 (15)	N1—C1—N3	109.09 (14)
C14—C13—H10	119.9	N1—C1—H1	125.5
C15—B1—C21	110.99 (13)	N2—C2—N3	111.91 (15)
C15—B1—C27	113.89 (14)	N2—C2—H2	124.0
C15—B1—N1	105.67 (13)	N3—C1—H1	125.5
C15—C16—C17	122.59 (16)	N3—C2—H2	124.0
C15—C16—H12	118.7		
B1—C15—C16—C17	179.90 (15)	C20—C15—B1—C21	-9.8 (2)
B1—C15—C20—C19	-179.44 (15)	C20—C15—B1—C27	118.01 (17)
B1—C21—C22—C23	173.63 (15)	C20—C15—B1—N1	-123.80 (16)
B1—C21—C26—C25	-173.56 (15)	C20—C15—C16—C17	0.9 (2)
B1—C27—C32—C31	-176.03 (16)	C20—C19—C18—C17	0.6 (3)
B1—N1—C1—N3	170.60 (14)	C22—C21—B1—C15	-75.75 (19)
B1—N1—N2—C2	-171.42 (15)	C22—C21—B1—C27	155.54 (15)
C1—N1—B1—C15	22.6 (2)	C22—C21—B1—N1	38.3 (2)
C1—N1—B1—C21	-95.12 (19)	C22—C21—C26—C25	-0.5 (2)
C1—N1—B1—C27	144.72 (16)	C22—C23—C24—C25	-0.5 (3)
C1—N1—N2—C2	-0.18 (19)	C24—C23—C22—C21	-0.2 (3)
C1—N3—C2—N2	0.4 (2)	C24—C25—C26—C21	-0.2 (3)
C1—N3—C3—C4	-156.01 (17)	C26—C21—B1—C15	96.88 (17)
C1—N3—C3—C8	26.6 (3)	C26—C21—B1—C27	-31.8 (2)
C2—N3—C1—N1	-0.54 (18)	C26—C21—B1—N1	-149.07 (14)
C2—N3—C3—C4	28.5 (3)	C26—C21—C22—C23	0.6 (2)
C2—N3—C3—C8	-148.83 (17)	C26—C25—C24—C23	0.6 (3)
C3—C4—C5—C6	-1.8 (3)	C27—C28—C29—C30	1.3 (3)
C3—C8—C7—C6	-2.0 (3)	C28—C27—B1—C15	-176.33 (14)
C3—N3—C1—N1	-176.73 (15)	C28—C27—B1—C21	-49.2 (2)
C3—N3—C2—N2	176.80 (15)	C28—C27—B1—N1	66.73 (19)
C5—C6—C7—C8	1.7 (3)	C28—C27—C32—C31	0.1 (2)
C6—N4—N5—C9	178.72 (14)	C29—C28—C27—B1	175.07 (15)
C7—C6—C5—C4	0.2 (3)	C29—C28—C27—C32	-1.1 (2)
C7—C8—C3—C4	0.4 (3)	C30—C31—C32—C27	0.7 (3)
C7—C8—C3—N3	177.64 (15)	C31—C30—C29—C28	-0.4 (3)
C8—C3—C4—C5	1.5 (3)	C32—C27—B1—C15	-0.4 (2)
C10—C9—C14—C13	-1.7 (3)	C32—C27—B1—C21	126.77 (17)
C10—C9—N5—N4	7.0 (2)	C32—C27—B1—N1	-117.34 (16)
C10—C11—C12—C13	-1.0 (3)	C32—C31—C30—C29	-0.6 (3)
C12—C11—C10—C9	0.1 (3)	N1—N2—C2—N3	-0.2 (2)
C12—C13—C14—C9	0.8 (3)	N2—N1—B1—C15	-168.22 (13)
C14—C9—C10—C11	1.3 (3)	N2—N1—B1—C21	74.09 (17)

supplementary materials

C14—C9—N5—N4	-173.24 (16)	N2—N1—B1—C27	-46.06 (19)
C14—C13—C12—C11	0.6 (3)	N2—N1—C1—N3	0.46 (18)
C15—C20—C19—C18	-0.3 (3)	N3—C3—C4—C5	-175.80 (16)
C16—C15—B1—C21	171.30 (14)	N4—C6—C5—C4	178.77 (16)
C16—C15—B1—C27	-60.91 (19)	N4—C6—C7—C8	-176.73 (15)
C16—C15—B1—N1	57.29 (18)	N5—C9—C10—C11	-178.98 (16)
C16—C15—C20—C19	-0.5 (2)	N5—C9—C14—C13	178.52 (17)
C16—C17—C18—C19	-0.2 (3)	N5—N4—C6—C5	166.55 (16)
C18—C17—C16—C15	-0.6 (2)	N5—N4—C6—C7	-15.0 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H3 \cdots Cg1 ⁱ	0.95	2.91	3.822 (2)	162

Symmetry codes: (i) $-x+1/2, y-1/2, -z+1/2$.

Fig. 1




Fig. 2

