

## 3,4-O-Isopropylidene-2,7-di-O-p-tolylsulfonyl-*α*-L-xylo-3-heptulo-3,6-furanosononitrile

Liang Ma,<sup>a</sup> Yun-Feng Li,<sup>a\*</sup> Xiang-Bao Meng<sup>a</sup> and Zhong-Jun Li<sup>b</sup>

<sup>a</sup>Department of Chemical Biology, School of Pharmaceutical Sciences, Peking University, Beijing 100191, People's Republic of China, and <sup>b</sup>State Key Laboratory of Natural and Biomimetic Drugs, Peking University, Beijing 100191, People's Republic of China  
Correspondence e-mail: zjli@bjmu.edu.cn


Received 7 April 2009; accepted 17 August 2009

Key indicators: single-crystal X-ray study;  $T = 113\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.049;  $wR$  factor = 0.127; data-to-parameter ratio = 14.1.

In the title compound,  $\text{C}_{24}\text{H}_{27}\text{NO}_{10}\text{S}_2$ , derived from L-sorbofuranose, the fused five-membered rings display envelope conformations. The two tosylate branches are in equatorial positions with respect to the furanose ring, while the hydroxy group is in the axial position. In the crystal structure, the hydroxy group is involved in intermolecular  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds, linking molecules in chains along [100].

### Related literature

For details of the synthesis, see: Bianchi *et al.* (2001); Georges & Fraser-Reid (1984); Sharma *et al.* (2003); Szarek *et al.* (1997).



### Experimental

#### Crystal data

$\text{C}_{24}\text{H}_{27}\text{NO}_{10}\text{S}_2$   
 $M_r = 553.59$   
Monoclinic,  $P2_1$   
 $a = 5.6342 (11)\text{ \AA}$   
 $b = 28.771 (6)\text{ \AA}$   
 $c = 8.3226 (17)\text{ \AA}$   
 $\beta = 103.00 (3)^\circ$

$V = 1314.5 (5)\text{ \AA}^3$   
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.26\text{ mm}^{-1}$   
 $T = 113\text{ K}$   
 $0.16 \times 0.12 \times 0.08\text{ mm}$

#### Data collection

Rigaku Saturn CCD area-detector diffractometer  
Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)  
 $T_{\min} = 0.960$ ,  $T_{\max} = 0.980$

8980 measured reflections  
4781 independent reflections  
4228 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.039$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.127$   
 $S = 1.05$   
4781 reflections  
339 parameters  
1 restraint

H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.31\text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.51\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
1570 Friedel pairs  
Flack parameter: 0.07 (7)

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O5—H5 $\cdots$ O4 <sup>i</sup>	0.82	2.03	2.844 (3)	169

Symmetry code: (i)  $x + 1, y, z$ .

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We acknowledge financial support from the National Natural Science Foundation of China.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2226).

### References

- Bianchi, P., Roda, G., Riva, S., Danieli, B., Zabelinskaja-Mackova, A. & Griengl, H. (2001). *Tetrahedron*, **57**, 2213–2220.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Georges, M. & Fraser-Reid, B. (1984). *Carbohydr. Res.* **127**, 162–164.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sharma, G. V. M., Begum, A., Reddy, K. R., Sankar, A. R. & Kunwar, A. C. (2003). *Tetrahedron Asymmetry*, **14**, 3899–3905.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Szarek, M. A., Wu, X. F. & Szarek, W. A. (1997). *Carbohydr. Res.* **299**, 165–170.

## **supplementary materials**

*Acta Cryst.* (2009). E65, o2437 [doi:10.1107/S1600536809032565]

### 3,4-*O*-Isopropylidene-2,7-di-*O*-*p*-tolylsulfonyl- $\alpha$ -L-xylo-3-heptulo-3,6-furanosononitrile

L. Ma, Y.-F. Li, X.-B. Meng and Z.-J. Li

#### Comment

The skeleton of the title compound is a derivative of L-sorbofuranose and consists of two tosyl, a cyano and a bridged ring. Both five-membered rings display an envelope conformation. The hydroxy group lies in axial bond of the furanose ring. The two tosylate branches lie in equatorial bonds of the five-membered furanose ring. The synthesis has been adapted from published procedures (Bianchi *et al.*, 2001; Georges & Fraser-Reid, 1984; Sharma *et al.*, 2003; Szarek *et al.*, 1997).

#### Experimental

The title compound was synthesized from the original compound 2,3:4,6-di-*O*-isopropylidene- $\alpha$ -L-sorbofuranose and the procedures are as follows: 2,3:4,6-di-*O*-isopropylidene- $\alpha$ -L-xylo-hexos-2-ulofuranose (obtained from the oxidization of 2,3:4,6-di-*O*-isopropylidene- $\alpha$ -L-sorbofuranose) was dissolved in HOAc (80%), followed by addition of NaCN, and after stirred at room temperature for 24 h., 3,4:5,7-di-*O*-isopropylidene- $\alpha$ -L-xylo-3-heptulo-3,6-furanosononitrile was formed. The reaction mixture was then directly heated at 323 K for 12 h. to remove 4,6-isopropylidene, yielding 3,4-*O*-isopropylidene- $\alpha$ -L-xylo-3-heptulo-3,6-furanosononitrile as a white solid after concentration *in vacuo* and subsequent column chromatography (petroleum-ethyl acetate 1:1). Finally 3,4-*O*-isopropylidene- $\alpha$ -L-xylo-3-heptulo-3,6-furanosononitrile was dissolved in pyridine followed by addition of tosyl chloride, and the reaction mixture was stirred at room temperature for 12 h., to obtain the title compound (55% yield over four steps), which was crystallized from petroleum-ethyl acetate (4:1). After a week at room temperature colourless block-like crystals were obtained.

#### Refinement

The completeness of diffraction data was limited to 0.945, because the crystal was lost before data collection was completed. The hydroxy H atom H5 was found in a difference map and refined with O—H bond length constrained to 0.82 Å, using a riding approximation, with  $U_{\text{iso}}(\text{H}5) = 1.5U_{\text{eq}}(\text{O}5)$ . C-bound H atoms were positioned geometrically (C—H = 0.93–0.98 Å) and refined as riding, with  $U_{\text{iso}}(\text{H})=1.2\text{--}1.5U_{\text{eq}}(\text{C})$ . The expected absolute configuration was confirmed by the refinement of a Flack parameter based on 1570 measured Friedel pairs.

#### Figures




Fig. 1. A view of the title molecule, showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented by circles of arbitrary size.

# supplementary materials

---

## 3,4-O-Isopropylidene-2,7-di-O-p-tolylsulfonyl- $\alpha$ -L-xylo-3-heptulo-3,6-furanosononitrile

### Crystal data

C <sub>24</sub> H <sub>27</sub> NO <sub>10</sub> S <sub>2</sub>	$F_{000} = 580$
$M_r = 553.59$	$D_x = 1.399 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: P 2yb	Cell parameters from 3787 reflections
$a = 5.6342 (11) \text{ \AA}$	$\theta = 1.5\text{--}27.5^\circ$
$b = 28.771 (6) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$c = 8.3226 (17) \text{ \AA}$	$T = 113 \text{ K}$
$\beta = 103.00 (3)^\circ$	Block, colourless
$V = 1314.5 (5) \text{ \AA}^3$	$0.16 \times 0.12 \times 0.08 \text{ mm}$
$Z = 2$	

### Data collection

Rigaku Saturn CCD area-detector diffractometer	4781 independent reflections
Radiation source: rotating anode	4228 reflections with $I > 2\sigma(I)$
Monochromator: confocal	$R_{\text{int}} = 0.039$
Detector resolution: 7.31 pixels $\text{mm}^{-1}$	$\theta_{\text{max}} = 27.9^\circ$
$T = 113 \text{ K}$	$\theta_{\text{min}} = 2.5^\circ$
$\omega$ and $\varphi$ scans	$h = -7 \rightarrow 7$
Absorption correction: multi-scan (CrystalClear; Rigaku, 2005)	$k = -37 \rightarrow 36$
$T_{\text{min}} = 0.960$ , $T_{\text{max}} = 0.980$	$l = -8 \rightarrow 10$
8980 measured reflections	

### Refinement

Refinement on $F^2$	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.049$	$w = 1/[\sigma^2(F_o^2) + (0.0656P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.127$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.05$	$\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$
4781 reflections	$\Delta\rho_{\text{min}} = -0.51 \text{ e \AA}^{-3}$
339 parameters	Extinction correction: none
1 restraint	Absolute structure: Flack (1983), 1570 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.07 (7)
Secondary atom site location: difference Fourier map	

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	-0.32882 (13)	0.19945 (2)	0.07345 (11)	0.01793 (18)
S2	-0.17570 (13)	0.48019 (2)	0.32977 (11)	0.01893 (18)
N1	0.2652 (6)	0.26412 (11)	0.3805 (5)	0.0315 (7)
O1	-0.4314 (4)	0.48295 (9)	0.3207 (3)	0.0262 (6)
O2	-0.0775 (4)	0.49967 (8)	0.2009 (3)	0.0252 (6)
O3	-0.1224 (4)	0.42629 (8)	0.3418 (3)	0.0210 (5)
O4	-0.0948 (4)	0.34836 (7)	0.1318 (3)	0.0196 (5)
O5	0.4380 (4)	0.35399 (8)	0.2068 (3)	0.0244 (5)
H5	0.5668	0.3487	0.1801	0.037*
O6	0.0421 (4)	0.35979 (9)	-0.2024 (3)	0.0267 (6)
O7	-0.2485 (4)	0.31572 (8)	-0.1269 (3)	0.0206 (5)
O8	-0.2598 (4)	0.25334 (7)	0.1007 (3)	0.0196 (5)
O9	-0.2258 (4)	0.18218 (8)	-0.0567 (3)	0.0232 (5)
O10	-0.5865 (4)	0.19956 (9)	0.0566 (3)	0.0232 (5)
C1	-0.0181 (6)	0.50230 (11)	0.5188 (5)	0.0199 (7)
C2	-0.1140 (7)	0.49476 (12)	0.6580 (5)	0.0283 (9)
H2	-0.2577	0.4781	0.6489	0.034*
C3	0.0070 (7)	0.51242 (13)	0.8098 (5)	0.0313 (9)
H3	-0.0570	0.5076	0.9021	0.038*
C4	0.2236 (7)	0.53735 (12)	0.8252 (6)	0.0300 (9)
C5	0.3154 (6)	0.54421 (13)	0.6857 (6)	0.0309 (9)
H5A	0.4591	0.5609	0.6944	0.037*
C6	0.1973 (6)	0.52669 (12)	0.5336 (5)	0.0245 (8)
H6	0.2627	0.5313	0.4417	0.029*
C7	0.3506 (8)	0.55689 (15)	0.9890 (6)	0.0409 (11)
H7A	0.5230	0.5576	0.9960	0.061*
H7B	0.3171	0.5377	1.0757	0.061*
H7C	0.2932	0.5879	1.0000	0.061*
C8	0.1045 (5)	0.40951 (11)	0.3035 (5)	0.0219 (8)
H8A	0.1761	0.3855	0.3814	0.026*
H8B	0.2201	0.4349	0.3119	0.026*
C9	0.0493 (5)	0.39035 (10)	0.1331 (5)	0.0188 (7)
H9	-0.0421	0.4132	0.0560	0.023*
C10	0.2720 (5)	0.37409 (11)	0.0732 (5)	0.0201 (7)
H10	0.3464	0.3996	0.0237	0.024*
C11	0.1613 (5)	0.33768 (11)	-0.0543 (4)	0.0203 (7)
H11	0.2793	0.3142	-0.0711	0.024*
C12	-0.1785 (6)	0.33503 (13)	-0.2688 (5)	0.0234 (7)
C13	-0.0521 (5)	0.31694 (10)	0.0094 (5)	0.0176 (7)
C14	-0.3723 (7)	0.36825 (15)	-0.3504 (5)	0.0363 (10)
H14A	-0.3906	0.3922	-0.2737	0.054*
H14B	-0.5236	0.3519	-0.3856	0.054*
H14C	-0.3271	0.3820	-0.4443	0.054*
C15	-0.1348 (8)	0.29602 (17)	-0.3803 (6)	0.0454 (12)
H15A	-0.0884	0.3088	-0.4753	0.068*

## supplementary materials

---

H15B	-0.2813	0.2782	-0.4149	0.068*
H15C	-0.0068	0.2763	-0.3216	0.068*
C16	-0.0202 (5)	0.26825 (11)	0.0850 (4)	0.0177 (7)
H16	0.0402	0.2471	0.0109	0.021*
C17	0.1446 (6)	0.26673 (11)	0.2511 (5)	0.0227 (7)
C18	-0.1844 (5)	0.17379 (10)	0.2592 (4)	0.0159 (6)
C19	0.0301 (6)	0.14921 (11)	0.2676 (5)	0.0222 (7)
H19	0.0925	0.1446	0.1745	0.027*
C20	0.1489 (6)	0.13163 (11)	0.4215 (5)	0.0252 (8)
H20	0.2918	0.1147	0.4297	0.030*
C21	0.0598 (6)	0.13870 (11)	0.5629 (5)	0.0240 (8)
C22	-0.1557 (6)	0.16375 (12)	0.5487 (5)	0.0258 (8)
H22	-0.2179	0.1687	0.6418	0.031*
C23	-0.2790 (6)	0.18142 (11)	0.3976 (5)	0.0212 (7)
H23	-0.4226	0.1981	0.3891	0.025*
C24	0.1933 (8)	0.12001 (14)	0.7275 (5)	0.0345 (10)
H24A	0.3418	0.1054	0.7159	0.052*
H24B	0.0928	0.0976	0.7662	0.052*
H24C	0.2304	0.1451	0.8052	0.052*

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0172 (3)	0.0193 (3)	0.0174 (5)	-0.0009 (3)	0.0043 (3)	-0.0008 (3)
S2	0.0184 (4)	0.0219 (4)	0.0171 (5)	-0.0002 (3)	0.0051 (3)	-0.0031 (3)
N1	0.0297 (15)	0.0327 (16)	0.030 (2)	0.0054 (13)	0.0026 (15)	0.0022 (15)
O1	0.0184 (10)	0.0325 (13)	0.0280 (16)	0.0019 (10)	0.0060 (11)	-0.0039 (12)
O2	0.0333 (13)	0.0257 (11)	0.0195 (15)	0.0003 (10)	0.0122 (12)	0.0025 (11)
O3	0.0190 (10)	0.0209 (11)	0.0250 (15)	-0.0015 (9)	0.0092 (10)	-0.0048 (10)
O4	0.0154 (10)	0.0175 (10)	0.0292 (15)	-0.0021 (8)	0.0120 (10)	-0.0044 (10)
O5	0.0133 (10)	0.0336 (13)	0.0263 (15)	0.0013 (9)	0.0044 (10)	-0.0012 (11)
O6	0.0279 (12)	0.0332 (12)	0.0208 (15)	-0.0061 (10)	0.0095 (12)	0.0024 (11)
O7	0.0162 (10)	0.0277 (12)	0.0160 (14)	-0.0020 (9)	-0.0004 (10)	0.0049 (10)
O8	0.0144 (9)	0.0201 (10)	0.0253 (15)	0.0019 (9)	0.0063 (10)	0.0009 (10)
O9	0.0295 (12)	0.0239 (11)	0.0186 (14)	-0.0007 (10)	0.0107 (11)	-0.0050 (10)
O10	0.0142 (9)	0.0299 (11)	0.0238 (15)	-0.0014 (9)	0.0009 (10)	0.0019 (12)
C1	0.0204 (14)	0.0194 (14)	0.021 (2)	0.0003 (12)	0.0069 (15)	-0.0031 (14)
C2	0.0321 (18)	0.0293 (17)	0.025 (2)	-0.0092 (14)	0.0108 (18)	0.0008 (16)
C3	0.041 (2)	0.0345 (19)	0.020 (2)	-0.0061 (16)	0.0082 (18)	0.0003 (16)
C4	0.036 (2)	0.0228 (16)	0.031 (3)	0.0041 (14)	0.0057 (18)	-0.0008 (16)
C5	0.0225 (17)	0.0346 (19)	0.035 (3)	-0.0055 (15)	0.0051 (17)	-0.0051 (18)
C6	0.0235 (16)	0.0269 (16)	0.026 (2)	-0.0009 (13)	0.0120 (16)	-0.0033 (15)
C7	0.050 (2)	0.036 (2)	0.032 (3)	0.0001 (18)	-0.001 (2)	-0.005 (2)
C8	0.0146 (13)	0.0243 (16)	0.028 (2)	0.0017 (12)	0.0065 (14)	-0.0043 (15)
C9	0.0149 (13)	0.0161 (13)	0.026 (2)	-0.0016 (11)	0.0054 (14)	-0.0006 (13)
C10	0.0150 (13)	0.0256 (15)	0.020 (2)	-0.0014 (12)	0.0055 (14)	-0.0026 (14)
C11	0.0156 (14)	0.0270 (15)	0.020 (2)	-0.0012 (12)	0.0070 (14)	-0.0046 (14)
C12	0.0213 (15)	0.0362 (18)	0.0130 (19)	-0.0070 (14)	0.0048 (14)	-0.0006 (15)

C13	0.0138 (13)	0.0168 (14)	0.022 (2)	0.0024 (11)	0.0030 (13)	-0.0015 (14)
C14	0.0342 (19)	0.043 (2)	0.029 (3)	-0.0029 (17)	0.0004 (19)	0.0153 (19)
C15	0.039 (2)	0.062 (3)	0.040 (3)	-0.011 (2)	0.018 (2)	-0.026 (2)
C16	0.0134 (13)	0.0208 (14)	0.0194 (19)	-0.0005 (11)	0.0047 (13)	-0.0009 (13)
C17	0.0215 (15)	0.0241 (15)	0.024 (2)	0.0028 (13)	0.0083 (16)	0.0011 (14)
C18	0.0181 (14)	0.0180 (14)	0.0123 (18)	-0.0009 (11)	0.0049 (14)	-0.0020 (12)
C19	0.0217 (15)	0.0247 (15)	0.021 (2)	0.0018 (13)	0.0055 (15)	0.0009 (14)
C20	0.0202 (15)	0.0239 (16)	0.030 (2)	0.0010 (13)	0.0032 (16)	0.0024 (15)
C21	0.0273 (17)	0.0215 (15)	0.018 (2)	-0.0019 (13)	-0.0061 (16)	0.0020 (14)
C22	0.0336 (19)	0.0255 (16)	0.020 (2)	0.0025 (14)	0.0090 (17)	0.0019 (15)
C23	0.0256 (16)	0.0247 (15)	0.0153 (19)	0.0011 (13)	0.0087 (15)	0.0014 (14)
C24	0.041 (2)	0.0337 (19)	0.021 (2)	-0.0005 (16)	-0.0092 (19)	0.0051 (18)

*Geometric parameters (Å, °)*

S1—O10	1.427 (2)	C8—H8A	0.9700
S1—O9	1.428 (2)	C8—H8B	0.9700
S1—O8	1.602 (2)	C9—C10	1.524 (4)
S1—C18	1.741 (4)	C9—H9	0.9800
S2—O2	1.427 (3)	C10—C11	1.521 (4)
S2—O1	1.428 (2)	C10—H10	0.9800
S2—O3	1.578 (2)	C11—C13	1.539 (4)
S2—C1	1.743 (4)	C11—H11	0.9800
N1—C17	1.139 (5)	C12—C14	1.494 (5)
O3—C8	1.467 (3)	C12—C15	1.512 (5)
O4—C13	1.423 (4)	C13—C16	1.530 (4)
O4—C9	1.454 (3)	C14—H14A	0.9600
O5—C10	1.406 (4)	C14—H14B	0.9600
O5—H5	0.8200	C14—H14C	0.9600
O6—C11	1.415 (4)	C15—H15A	0.9600
O6—C12	1.431 (4)	C15—H15B	0.9600
O7—C13	1.396 (4)	C15—H15C	0.9600
O7—C12	1.438 (4)	C16—C17	1.483 (5)
O8—C16	1.450 (3)	C16—H16	0.9800
C1—C6	1.383 (4)	C18—C19	1.388 (4)
C1—C2	1.401 (5)	C18—C23	1.391 (5)
C2—C3	1.390 (6)	C19—C20	1.399 (5)
C2—H2	0.9300	C19—H19	0.9300
C3—C4	1.396 (5)	C20—C21	1.394 (5)
C3—H3	0.9300	C20—H20	0.9300
C4—C5	1.388 (6)	C21—C22	1.394 (5)
C4—C7	1.499 (6)	C21—C24	1.506 (5)
C5—C6	1.386 (6)	C22—C23	1.389 (5)
C5—H5A	0.9300	C22—H22	0.9300
C6—H6	0.9300	C23—H23	0.9300
C7—H7A	0.9600	C24—H24A	0.9600
C7—H7B	0.9600	C24—H24B	0.9600
C7—H7C	0.9600	C24—H24C	0.9600
C8—C9	1.488 (5)		

## supplementary materials

---

O10—S1—O9	120.29 (15)	O6—C11—C13	102.7 (2)
O10—S1—O8	102.76 (13)	C10—C11—C13	104.9 (3)
O9—S1—O8	108.47 (13)	O6—C11—H11	112.9
O10—S1—C18	110.50 (14)	C10—C11—H11	112.9
O9—S1—C18	109.72 (15)	C13—C11—H11	112.9
O8—S1—C18	103.57 (14)	O6—C12—O7	104.5 (3)
O2—S2—O1	119.71 (16)	O6—C12—C14	109.8 (3)
O2—S2—O3	109.48 (13)	O7—C12—C14	108.2 (3)
O1—S2—O3	103.49 (13)	O6—C12—C15	111.1 (3)
O2—S2—C1	109.25 (16)	O7—C12—C15	109.3 (3)
O1—S2—C1	109.28 (16)	C14—C12—C15	113.5 (4)
O3—S2—C1	104.48 (15)	O7—C13—O4	111.5 (2)
C8—O3—S2	118.17 (19)	O7—C13—C16	108.0 (2)
C13—O4—C9	110.4 (2)	O4—C13—C16	108.1 (3)
C10—O5—H5	109.5	O7—C13—C11	105.6 (3)
C11—O6—C12	108.3 (3)	O4—C13—C11	105.7 (2)
C13—O7—C12	110.1 (2)	C16—C13—C11	117.9 (2)
C16—O8—S1	118.35 (17)	C12—C14—H14A	109.5
C6—C1—C2	119.9 (4)	C12—C14—H14B	109.5
C6—C1—S2	121.6 (3)	H14A—C14—H14B	109.5
C2—C1—S2	118.5 (3)	C12—C14—H14C	109.5
C3—C2—C1	119.6 (3)	H14A—C14—H14C	109.5
C3—C2—H2	120.2	H14B—C14—H14C	109.5
C1—C2—H2	120.2	C12—C15—H15A	109.5
C2—C3—C4	120.8 (4)	C12—C15—H15B	109.5
C2—C3—H3	119.6	H15A—C15—H15B	109.5
C4—C3—H3	119.6	C12—C15—H15C	109.5
C5—C4—C3	118.6 (4)	H15A—C15—H15C	109.5
C5—C4—C7	121.1 (4)	H15B—C15—H15C	109.5
C3—C4—C7	120.4 (4)	O8—C16—C17	107.9 (3)
C6—C5—C4	121.3 (3)	O8—C16—C13	106.4 (2)
C6—C5—H5A	119.3	C17—C16—C13	113.7 (3)
C4—C5—H5A	119.3	O8—C16—H16	109.6
C1—C6—C5	119.8 (3)	C17—C16—H16	109.6
C1—C6—H6	120.1	C13—C16—H16	109.6
C5—C6—H6	120.1	N1—C17—C16	177.1 (4)
C4—C7—H7A	109.5	C19—C18—C23	122.0 (3)
C4—C7—H7B	109.5	C19—C18—S1	119.2 (3)
H7A—C7—H7B	109.5	C23—C18—S1	118.6 (2)
C4—C7—H7C	109.5	C18—C19—C20	117.5 (3)
H7A—C7—H7C	109.5	C18—C19—H19	121.2
H7B—C7—H7C	109.5	C20—C19—H19	121.2
O3—C8—C9	108.9 (3)	C21—C20—C19	122.1 (3)
O3—C8—H8A	109.9	C21—C20—H20	119.0
C9—C8—H8A	109.9	C19—C20—H20	119.0
O3—C8—H8B	109.9	C22—C21—C20	118.4 (3)
C9—C8—H8B	109.9	C22—C21—C24	120.6 (4)
H8A—C8—H8B	108.3	C20—C21—C24	121.0 (3)
O4—C9—C8	108.1 (3)	C23—C22—C21	120.9 (4)

O4—C9—C10	104.0 (2)	C23—C22—H22	119.5
C8—C9—C10	114.5 (3)	C21—C22—H22	119.5
O4—C9—H9	110.0	C22—C23—C18	119.0 (3)
C8—C9—H9	110.0	C22—C23—H23	120.5
C10—C9—H9	110.0	C18—C23—H23	120.5
O5—C10—C11	111.3 (3)	C21—C24—H24A	109.5
O5—C10—C9	108.5 (3)	C21—C24—H24B	109.5
C11—C10—C9	101.6 (2)	H24A—C24—H24B	109.5
O5—C10—H10	111.6	C21—C24—H24C	109.5
C11—C10—H10	111.6	H24A—C24—H24C	109.5
C9—C10—H10	111.6	H24B—C24—H24C	109.5
O6—C11—C10	109.7 (3)		
O2—S2—O3—C8	33.2 (3)	C13—O7—C12—O6	-16.2 (3)
O1—S2—O3—C8	161.9 (3)	C13—O7—C12—C14	-133.2 (3)
C1—S2—O3—C8	-83.7 (3)	C13—O7—C12—C15	102.7 (3)
O10—S1—O8—C16	-166.0 (2)	C12—O7—C13—O4	112.8 (3)
O9—S1—O8—C16	-37.6 (3)	C12—O7—C13—C16	-128.6 (3)
C18—S1—O8—C16	78.9 (3)	C12—O7—C13—C11	-1.6 (3)
O2—S2—C1—C6	-12.2 (3)	C9—O4—C13—O7	-105.8 (3)
O1—S2—C1—C6	-144.9 (3)	C9—O4—C13—C16	135.5 (3)
O3—S2—C1—C6	104.8 (3)	C9—O4—C13—C11	8.4 (3)
O2—S2—C1—C2	168.0 (3)	O6—C11—C13—O7	18.8 (3)
O1—S2—C1—C2	35.3 (3)	C10—C11—C13—O7	133.5 (3)
O3—S2—C1—C2	-75.0 (3)	O6—C11—C13—O4	-99.5 (3)
C6—C1—C2—C3	0.8 (5)	C10—C11—C13—O4	15.2 (3)
S2—C1—C2—C3	-179.4 (3)	O6—C11—C13—C16	139.6 (3)
C1—C2—C3—C4	-0.3 (6)	C10—C11—C13—C16	-105.7 (3)
C2—C3—C4—C5	0.1 (6)	S1—O8—C16—C17	-92.6 (3)
C2—C3—C4—C7	179.0 (4)	S1—O8—C16—C13	145.0 (2)
C3—C4—C5—C6	-0.3 (6)	O7—C13—C16—O8	-48.4 (3)
C7—C4—C5—C6	-179.2 (4)	O4—C13—C16—O8	72.5 (3)
C2—C1—C6—C5	-1.0 (5)	C11—C13—C16—O8	-167.8 (3)
S2—C1—C6—C5	179.2 (3)	O7—C13—C16—C17	-167.0 (3)
C4—C5—C6—C1	0.8 (6)	O4—C13—C16—C17	-46.2 (3)
S2—O3—C8—C9	-100.7 (3)	C11—C13—C16—C17	73.5 (4)
C13—O4—C9—C8	-150.7 (2)	O10—S1—C18—C19	147.6 (3)
C13—O4—C9—C10	-28.7 (3)	O9—S1—C18—C19	12.7 (3)
O3—C8—C9—O4	-67.9 (3)	O8—S1—C18—C19	-103.0 (3)
O3—C8—C9—C10	176.7 (3)	O10—S1—C18—C23	-36.9 (3)
O4—C9—C10—O5	-81.1 (3)	O9—S1—C18—C23	-171.8 (2)
C8—C9—C10—O5	36.6 (4)	O8—S1—C18—C23	72.5 (3)
O4—C9—C10—C11	36.3 (3)	C23—C18—C19—C20	0.7 (5)
C8—C9—C10—C11	154.0 (3)	S1—C18—C19—C20	176.0 (2)
C12—O6—C11—C10	-140.6 (3)	C18—C19—C20—C21	-0.9 (5)
C12—O6—C11—C13	-29.4 (3)	C19—C20—C21—C22	0.6 (5)
O5—C10—C11—O6	-166.2 (2)	C19—C20—C21—C24	-179.1 (3)
C9—C10—C11—O6	78.4 (3)	C20—C21—C22—C23	-0.2 (5)
O5—C10—C11—C13	84.1 (3)	C24—C21—C22—C23	179.5 (3)
C9—C10—C11—C13	-31.3 (3)	C21—C22—C23—C18	0.0 (5)

## supplementary materials

---

C11—O6—C12—O7	29.1 (3)	C19—C18—C23—C22	-0.3 (5)
C11—O6—C12—C14	145.0 (3)	S1—C18—C23—C22	-175.7 (3)
C11—O6—C12—C15	-88.6 (4)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$\text{H}\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O5—H5 $\cdots$ O4 <sup>i</sup>	0.82	2.03	2.844 (3)	169

Symmetry codes: (i)  $x+1, y, z$ .

**Fig. 1**

