

4-Bromomethyl-7-methyl-6,8-dinitrocoumarin

Ramakrishna Gowda,^{a*} Ganesh N. Alawandi,^b Manohar V. Kulkarni^b and K. V. Arjuna Gowda^c

^aDepartment of Physics, Goverment College for Women, Kolar 563 101, Karnataka, India, ^bDepartment of Chemistry, Karnatak University, Dharwad 580 003, Karnataka, India, and ^cDepartment of Physics, Goverment First Grade College, K.R. Pura, Bangalore 560 036, Karnataka, India

Correspondence e-mail: arjunagowda@indiainfo.com


Received 12 July 2009; accepted 5 September 2009

Key indicators: single-crystal X-ray study; $T = 294\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.010\text{ \AA}$; R factor = 0.060; wR factor = 0.171; data-to-parameter ratio = 12.1.

The crystal structure of the title compound, $\text{C}_{11}\text{H}_7\text{BrN}_2\text{O}_6$, establishes the substitution positions of the nitro groups from the nitration reaction of 7-methyl-4-bromomethyl coumarin. The mean planes of the nitro groups form dihedral angles of 43.9 (8) and 52.7 (10) $^\circ$ with the essentially planar [maximum deviation 0.031 (6) \AA] benzopyran ring system.


Related literature

For background information on the nitration of coumarin compounds, see: Kulkarni *et al.* (1983); Clayton *et al.* (1910). For a related structure, see: Vasudevan *et al.* (1990). For *ab initio* calculations on 6-methyl-4-bromomethylcoumarins, see: Sortur *et al.* (2006).



Experimental

Crystal data



Orthorhombic, $Pbca$
 $a = 8.122 (2)\text{ \AA}$
 $b = 11.091 (4)\text{ \AA}$
 $c = 27.723 (6)\text{ \AA}$
 $V = 2497.3 (12)\text{ \AA}^3$

$Z = 8$
Mo $K\alpha$ radiation
 $\mu = 3.32\text{ mm}^{-1}$
 $T = 294\text{ K}$
 $0.2 \times 0.2 \times 0.1\text{ mm}$

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.520$, $T_{\max} = 0.72$
2196 measured reflections

2196 independent reflections
1148 reflections with $I > 2\sigma(I)$
2 standard reflections
frequency: 60 min
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$
 $wR(F^2) = 0.171$
 $S = 1.05$
2196 reflections

182 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.63\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.82\text{ e \AA}^{-3}$

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

KVAG gratefully thanks the DST for financial support through the SERC Fast Track Young Scientists Scheme and RG thanks MVJ College of Engineering Bangalore (Research Center). The authors also thanks Professor T. N. Guru Row, Chairman, SSCU IISc, Bangalore, for the X-ray data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2864).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
Clayton, A. (1910). *J. Chem. Soc.* **97**, 1397–1408.
Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
Kulkarni, M. V., Pujar, B. G. & Patil, V. D. (1983). *Arch. Pharm.* **316**, 15–21.
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Sortur, V., Yenagi, J., Tonannavar, J., Jadhav, V. B. & Kulkarni, M. V. (2006). *Spectrochim. Acta A*, **64**, 301–307.
Vasudevan, K. T., Puttaraja, & Kulkarni, M. V. (1990). *Acta Cryst. C* **46**, 2129–2131.

supplementary materials

Acta Cryst. (2009). E65, o2446 [doi:10.1107/S160053680903596X]

4-Bromomethyl-7-methyl-6,8-dinitrocoumarin

R. Gowda, G. N. Alawandi, M. V. Kulkarni and K. V. A. Gowda

Comment

The molecular structure of the title compound is shown in Fig. 1. The bromomethyl group is twisted out of the plane of the benzopyran ring as described by the torsion angle of $104.6(7)^\circ$ for C2-C3-C10-Br. This is in agreement with the ab initio calculations on 6-methyl-4-bromomethylcoumarins (Sortur *et al.*, 2006). Positions C-6 and C-8 (refers to positions from systematic naming scheme) become activated due to the electron donating methyl group at C-7 and hence nitration occurs at C-6 and C-8 consistent with the title compound which is also in agreement with earlier reports (Clayton, 1910; Kulkarni *et al.*, 1983).

Experimental

5.06 g of 7-methyl-4-bromomethyl coumarin (0.02 mol) was dissolved in conc. sulfuric acid (10 ml) and treated with a nitrating mixture 15 ml (10 ml H_2SO_4 + 5 ml HNO_3) at ice bath temperatures (273–278 K). The reaction mixture was then allowed to stand at room temperature for two hours and the reaction mixture was poured over crushed ice. The separated solid was washed with excess of water, dried and recrystallized from glacial acetic acid. Crystals suitable for diffraction studies were grown by slow evaporation of an ethanol solution of the title compound.

Refinement

Hydrogen atoms were positioned geometrically with C—H = 0.93–0.97 Å and included in the refinement in a riding-model approximation with $U_{iso}(H) = 1.2 U_{eq}(C)$ or $1.5 U_{eq}(C)$ for methyl C atoms.

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atomic numbering.

4-Bromomethyl-7-methyl-6,8-dinitrocoumarin

Crystal data

| | |
|---------------------|---------------------------------|
| $C_{11}H_7BrN_2O_6$ | $F_{000} = 1360$ |
| $M_r = 343.09$ | $D_x = 1.825 \text{ Mg m}^{-3}$ |

supplementary materials

| | |
|--------------------------------|---|
| Orthorhombic, <i>Pbca</i> | Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$ |
| Hall symbol: -P 2ac 2ab | Cell parameters from 25 reflections |
| $a = 8.122(2) \text{ \AA}$ | $\theta = 10\text{--}15^\circ$ |
| $b = 11.091(4) \text{ \AA}$ | $\mu = 3.32 \text{ mm}^{-1}$ |
| $c = 27.723(6) \text{ \AA}$ | $T = 294 \text{ K}$ |
| $V = 2497.3(12) \text{ \AA}^3$ | Plate, colourless |
| $Z = 8$ | $0.2 \times 0.2 \times 0.1 \text{ mm}$ |

Data collection

| | |
|--|------------------------------------|
| Enraf–Nonius CAD-4 diffractometer | $R_{\text{int}} = 0.0000$ |
| Radiation source: fine-focus sealed tube | $\theta_{\text{max}} = 25.0^\circ$ |
| Monochromator: graphite | $\theta_{\text{min}} = 2.9^\circ$ |
| $T = 294 \text{ K}$ | $h = 0 \rightarrow 9$ |
| ω - 2θ scans | $k = 0 \rightarrow 13$ |
| Absorption correction: ψ scan (North <i>et al.</i> , 1968) | $l = 0 \rightarrow 32$ |
| $T_{\text{min}} = 0.520$, $T_{\text{max}} = 0.72$ | 2 standard reflections |
| 2196 measured reflections | every 60 min |
| 2196 independent reflections | intensity decay: none |
| 1148 reflections with $I > 2\sigma(I)$ | |

Refinement

| | |
|---------------------------------|--|
| Refinement on F^2 | 182 parameters |
| Least-squares matrix: full | H-atom parameters constrained |
| $R[F^2 > 2\sigma(F^2)] = 0.060$ | $w = 1/[\sigma^2(F_o^2) + (0.0687P)^2 + 11.8667P]$ |
| | where $P = (F_o^2 + 2F_c^2)/3$ |
| $wR(F^2) = 0.171$ | $(\Delta/\sigma)_{\text{max}} = 0.001$ |
| $S = 1.05$ | $\Delta\rho_{\text{max}} = 0.63 \text{ e \AA}^{-3}$ |
| 2196 reflections | $\Delta\rho_{\text{min}} = -0.82 \text{ e \AA}^{-3}$ |

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|--------------|-------------|--------------|----------------------------------|
| C1 | 0.7703 (11) | 0.2886 (7) | 0.6819 (3) | 0.047 (2) |
| C2 | 0.9438 (10) | 0.2873 (7) | 0.6922 (3) | 0.045 (2) |
| H3 | 0.9856 | 0.3447 | 0.7134 | 0.054* |
| C3 | 1.0484 (9) | 0.2073 (6) | 0.6727 (3) | 0.0380 (18) |
| C4 | 1.0779 (9) | 0.0341 (6) | 0.6146 (3) | 0.0364 (18) |
| H5 | 1.1898 | 0.0281 | 0.6213 | 0.044* |
| C5 | 1.0085 (9) | -0.0431 (6) | 0.5812 (2) | 0.0348 (17) |
| C6 | 0.8400 (9) | -0.0445 (6) | 0.5709 (3) | 0.0395 (19) |
| C7 | 0.7498 (9) | 0.0406 (6) | 0.5954 (2) | 0.0358 (16) |
| C8 | 0.9860 (8) | 0.1194 (6) | 0.6381 (3) | 0.0326 (17) |
| C9 | 0.8194 (9) | 0.1222 (6) | 0.6280 (3) | 0.0339 (17) |
| C10 | 1.2259 (10) | 0.2058 (7) | 0.6867 (3) | 0.049 (2) |
| H11A | 1.2542 | 0.2804 | 0.7030 | 0.059* |
| H11B | 1.2942 | 0.1987 | 0.6581 | 0.059* |
| C11 | 0.7574 (12) | -0.1318 (7) | 0.5373 (3) | 0.057 (2) |
| H12A | 0.8385 | -0.1842 | 0.5234 | 0.085* |
| H12B | 0.7020 | -0.0882 | 0.5122 | 0.085* |
| H12C | 0.6787 | -0.1790 | 0.5550 | 0.085* |
| N1 | 1.1229 (9) | -0.1278 (6) | 0.5564 (3) | 0.0483 (17) |
| N2 | 0.5699 (8) | 0.0459 (6) | 0.5897 (3) | 0.0477 (17) |
| O1 | 0.7146 (6) | 0.2044 (4) | 0.64815 (18) | 0.0406 (13) |
| O2 | 0.6689 (8) | 0.3553 (6) | 0.6980 (2) | 0.0649 (18) |
| O3 | 1.2252 (9) | -0.1757 (6) | 0.5791 (3) | 0.085 (2) |
| O4 | 1.1032 (9) | -0.1384 (6) | 0.5134 (3) | 0.079 (2) |
| O5 | 0.5137 (9) | 0.0940 (7) | 0.5564 (3) | 0.099 (3) |
| O6 | 0.4882 (8) | 0.0020 (9) | 0.6212 (3) | 0.113 (3) |
| Br | 1.26518 (11) | 0.06847 (7) | 0.72976 (3) | 0.0597 (4) |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-----------|-----------|-----------|------------|------------|------------|
| C1 | 0.061 (6) | 0.043 (4) | 0.037 (4) | 0.004 (5) | -0.002 (5) | -0.002 (4) |
| C2 | 0.060 (6) | 0.036 (4) | 0.040 (4) | -0.001 (4) | -0.002 (4) | -0.006 (4) |
| C3 | 0.040 (4) | 0.039 (4) | 0.034 (4) | -0.008 (4) | -0.002 (3) | 0.004 (3) |
| C4 | 0.034 (4) | 0.038 (4) | 0.038 (4) | 0.000 (3) | -0.004 (3) | -0.001 (3) |
| C5 | 0.039 (4) | 0.036 (4) | 0.029 (4) | 0.007 (3) | 0.003 (3) | -0.002 (3) |
| C6 | 0.045 (5) | 0.040 (4) | 0.034 (4) | -0.003 (4) | -0.009 (4) | 0.001 (3) |
| C7 | 0.030 (4) | 0.039 (4) | 0.039 (4) | -0.002 (4) | -0.002 (4) | 0.006 (3) |
| C8 | 0.028 (4) | 0.034 (4) | 0.036 (4) | -0.004 (3) | 0.000 (3) | -0.003 (3) |
| C9 | 0.033 (4) | 0.037 (4) | 0.032 (4) | 0.007 (3) | 0.004 (3) | 0.005 (3) |
| C10 | 0.056 (6) | 0.048 (4) | 0.044 (4) | -0.013 (4) | -0.004 (4) | -0.001 (4) |
| C11 | 0.072 (6) | 0.040 (4) | 0.058 (5) | -0.001 (5) | -0.024 (5) | -0.011 (4) |
| N1 | 0.049 (5) | 0.049 (4) | 0.045 (5) | 0.001 (4) | 0.001 (4) | -0.001 (4) |
| N2 | 0.037 (4) | 0.046 (4) | 0.058 (5) | 0.001 (3) | -0.012 (4) | 0.001 (4) |

supplementary materials

| | | | | | | |
|----|------------|------------|------------|------------|-------------|-------------|
| O1 | 0.031 (3) | 0.049 (3) | 0.042 (3) | 0.010 (2) | -0.003 (2) | -0.002 (3) |
| O2 | 0.071 (4) | 0.063 (4) | 0.061 (4) | 0.026 (4) | 0.003 (3) | -0.013 (3) |
| O3 | 0.074 (5) | 0.092 (5) | 0.088 (5) | 0.039 (4) | -0.010 (4) | -0.023 (4) |
| O4 | 0.102 (6) | 0.076 (4) | 0.059 (5) | 0.023 (4) | 0.012 (4) | -0.014 (4) |
| O5 | 0.060 (5) | 0.130 (7) | 0.108 (6) | 0.004 (4) | -0.033 (5) | 0.048 (5) |
| O6 | 0.038 (4) | 0.160 (8) | 0.141 (8) | -0.011 (5) | -0.002 (5) | 0.073 (7) |
| Br | 0.0621 (6) | 0.0560 (5) | 0.0611 (6) | 0.0056 (5) | -0.0181 (5) | -0.0042 (4) |

Geometric parameters (\AA , $^\circ$)

| | | | |
|-------------|------------|---------------|------------|
| C1—O2 | 1.194 (9) | C7—C9 | 1.397 (10) |
| C1—O1 | 1.398 (9) | C7—N2 | 1.471 (10) |
| C1—C2 | 1.438 (11) | C8—C9 | 1.382 (9) |
| C2—C3 | 1.342 (10) | C9—O1 | 1.368 (8) |
| C2—H3 | 0.9300 | C10—H11A | 0.9700 |
| C3—C8 | 1.458 (10) | C10—H11B | 0.9700 |
| C3—C10 | 1.494 (11) | C11—H12A | 0.9600 |
| C4—C8 | 1.370 (10) | C11—H12B | 0.9600 |
| C4—C5 | 1.383 (9) | C11—H12C | 0.9600 |
| C4—H5 | 0.9300 | N1—O3 | 1.169 (9) |
| C5—C6 | 1.398 (10) | N1—O4 | 1.210 (9) |
| C5—N1 | 1.488 (10) | N2—O5 | 1.160 (8) |
| C6—C7 | 1.375 (10) | N2—O6 | 1.201 (9) |
| C6—C11 | 1.501 (10) | | |
| O2—C1—O1 | 116.2 (8) | C9—C8—C3 | 117.3 (7) |
| O2—C1—C2 | 127.4 (8) | O1—C9—C8 | 122.8 (7) |
| O1—C1—C2 | 116.3 (7) | O1—C9—C7 | 116.4 (6) |
| C3—C2—C1 | 123.1 (7) | C8—C9—C7 | 120.9 (7) |
| C3—C2—H3 | 118.4 | C3—C10—Br | 108.9 (5) |
| C1—C2—H3 | 118.4 | C3—C10—H11A | 109.9 |
| C2—C3—C8 | 119.2 (7) | Br—C10—H11A | 109.9 |
| C2—C3—C10 | 120.9 (7) | C3—C10—H11B | 109.9 |
| C8—C3—C10 | 120.0 (7) | Br—C10—H11B | 109.9 |
| C8—C4—C5 | 121.6 (7) | H11A—C10—H11B | 108.3 |
| C8—C4—H5 | 119.2 | C6—C11—H12A | 109.5 |
| C5—C4—H5 | 119.2 | C6—C11—H12B | 109.5 |
| C4—C5—C6 | 122.9 (7) | H12A—C11—H12B | 109.5 |
| C4—C5—N1 | 116.5 (7) | C6—C11—H12C | 109.5 |
| C6—C5—N1 | 120.7 (7) | H12A—C11—H12C | 109.5 |
| C7—C6—C5 | 114.4 (7) | H12B—C11—H12C | 109.5 |
| C7—C6—C11 | 120.8 (7) | O3—N1—O4 | 125.5 (8) |
| C5—C6—C11 | 124.8 (7) | O3—N1—C5 | 118.9 (7) |
| C6—C7—C9 | 123.3 (7) | O4—N1—C5 | 115.6 (7) |
| C6—C7—N2 | 120.2 (7) | O5—N2—O6 | 123.2 (8) |
| C9—C7—N2 | 116.5 (6) | O5—N2—C7 | 119.7 (8) |
| C4—C8—C9 | 116.9 (7) | O6—N2—C7 | 117.0 (7) |
| C4—C8—C3 | 125.8 (7) | C9—O1—C1 | 121.2 (6) |
| O2—C1—C2—C3 | 179.1 (8) | C3—C8—C9—O1 | 1.3 (10) |
| O1—C1—C2—C3 | -2.9 (11) | C4—C8—C9—C7 | 0.4 (11) |

supplementary materials

| | | | |
|--------------|------------|--------------|------------|
| C1—C2—C3—C8 | 2.1 (11) | C3—C8—C9—C7 | -179.7 (6) |
| C1—C2—C3—C10 | -176.6 (7) | C6—C7—C9—O1 | 177.6 (6) |
| C8—C4—C5—C6 | -3.6 (11) | N2—C7—C9—O1 | -4.9 (9) |
| C8—C4—C5—N1 | 177.2 (7) | C6—C7—C9—C8 | -1.5 (11) |
| C4—C5—C6—C7 | 2.5 (11) | N2—C7—C9—C8 | 176.0 (7) |
| N1—C5—C6—C7 | -178.4 (6) | C2—C3—C10—Br | 104.6 (7) |
| C4—C5—C6—C11 | -175.8 (7) | C8—C3—C10—Br | -74.1 (7) |
| N1—C5—C6—C11 | 3.3 (11) | C4—C5—N1—O3 | 41.9 (11) |
| C5—C6—C7—C9 | 0.0 (10) | C6—C5—N1—O3 | -137.3 (8) |
| C11—C6—C7—C9 | 178.4 (7) | C4—C5—N1—O4 | -136.2 (8) |
| C5—C6—C7—N2 | -177.3 (6) | C6—C5—N1—O4 | 44.6 (10) |
| C11—C6—C7—N2 | 1.0 (11) | C6—C7—N2—O5 | -81.1 (10) |
| C5—C4—C8—C9 | 2.0 (11) | C9—C7—N2—O5 | 101.4 (9) |
| C5—C4—C8—C3 | -177.9 (7) | C6—C7—N2—O6 | 101.1 (10) |
| C2—C3—C8—C4 | 178.7 (7) | C9—C7—N2—O6 | -76.4 (10) |
| C10—C3—C8—C4 | -2.6 (11) | C8—C9—O1—C1 | -2.2 (10) |
| C2—C3—C8—C9 | -1.2 (10) | C7—C9—O1—C1 | 178.7 (6) |
| C10—C3—C8—C9 | 177.5 (7) | O2—C1—O1—C9 | -178.9 (7) |
| C4—C8—C9—O1 | -178.6 (6) | C2—C1—O1—C9 | 2.9 (10) |

supplementary materials

Fig. 1

