

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,4-Bis(pyridin-3-ylmethoxy)benzene

Jin-Sheng Gao,* Ying Liu, Shuang Zhang, Dian-Fa Zuo and Guang-Feng Hou

College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China

Correspondence e-mail: hgf1000@163.com


Received 4 September 2009; accepted 8 September 2009

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.041; wR factor = 0.118; data-to-parameter ratio = 16.8.

The asymmetric unit of the centrosymmetric title compound, $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2$, contains one half-molecule. The central benzene ring forms a dihedral angle of $66.8(1)^\circ$ with two outer aromatic rings. In the crystal structure, weak intermolecular $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds link molecules into sheets parallel to (104).

Related literature

For general background to bridging molecules with pyridyl substituents at the terminal positions, see: McMorran & Steel (1998); Zaman *et al.* (2005). For details of the synthesis, see: Gao *et al.* (2004). For a related structure, see: Gao *et al.* (2006).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_2$
 $M_r = 292.33$
 Monoclinic, $P2_1/c$
 $a = 6.852(5)$ Å

$b = 5.688(3)$ Å
 $c = 18.861(12)$ Å
 $\beta = 90.60(3)^\circ$
 $V = 735.0(8)$ Å³

$Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹

$T = 291$ K
 $0.22 \times 0.20 \times 0.19$ mm

Data collection

Rigaku RAXIS-RAPID diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.981$, $T_{\max} = 0.984$

6855 measured reflections
 1684 independent reflections
 1213 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.030$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.118$
 $S = 1.09$
 1684 reflections

100 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.24$ e Å⁻³
 $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1}\cdots\text{N1}^i$	0.93	2.57	3.437 (3)	155

Symmetry code: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the Specialized Research Funds for Technological Innovative Talent in Harbin (RC2009XK018007) and Heilongjiang University for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2613).

References

- Gao, C.-M., Cao, D. & Zhu, L. (2004). *Photogr. Sci. Photochem.* **22**, 103–107.
 Gao, J.-S., Liu, Y., Hou, G.-F., Yu, Y.-H. & Yan, P.-F. (2006). *Acta Cryst.* **E62**, o5645–o5646.
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
 McMorran, D. A. & Steel, P. J. (1998). *Angew. Chem. Int. Ed.* **37**, 3295–3297.
 Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
 Rigaku/MSK (2002). *CrystalClear*. Rigaku/MSK Inc., The Woodlands, Texas, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zaman, Md. B., Udachin, K., Ripmeester, J. A., Smith, M. D. & zur Loye, H.-C. (2005). *Inorg. Chem.* **44**, 5047–5059.

supplementary materials

Acta Cryst. (2009). E65, o2457 [doi:10.1107/S1600536809036265]

1,4-Bis(pyridin-3-ylmethoxy)benzene

J.-S. Gao, Y. Liu, S. Zhang, D.-F. Zuo and G.-F. Hou

Comment

The bridging molecules with pyridyl substituents at the terminal positions are hoped to construct interesting supramolecular architectures by intermolecular hydrogen bonding and coordination with metals. McMorran' group have reported the synthesis of a quadruply stranded helicate that encapsulates a hexafluorophosphate anion by the reaction of 1,4-bis(3-pyridylmethoxy)benzene with palladium chlorate (McMorran *et al.*, 1998). Zaman's group have designed a long rigid organic ligand, 1,4-bis[(3-pyridyl)ethynyl]benzene, which reacted with metal salts to form interpenetrating two-dimensional and three-dimensional cross-zigzag chains and metallocyclic chain structures (Zaman *et al.*, 2005). As an extension of our work about bipyridyl aromatic ligands, we report the crystal structure of the title compound here.

In the title compound (Fig. 1), all bond lengths and angles are normal and correpond to those observed in the related compound (Gao *et al.*, 2006). The 1,4-bis(3-pyridylmethoxy)benzene molecule is centrosymmetric. The planes of two terminal pyridyl groups rotate drastically and make dihedral angles of $66.8(1)^\circ$ with the plane of the central benzene ring.

In the crystal structure, the adjacent 1,4-bis(3-pyridylmethoxy)benzene molecules are linked into two-dimensional supramolecular sheets by intermolecular C—H \cdots N hydrogen bonds (Table 1, Figure 2).

Experimental

The 1,4-bis(3-pyridylmethoxy)benzene was synthesized by the reaction of *p*-benzenediol and 3-chloromethylpyridine hydrochloride under nitrogen atmosphere and alkaline condition (Gao *et al.*, 2004; Gao *et al.*, 2006). Colourless block-shaped crystals of title compound were obtained by slow evaporation of an methanol solution after three days.

Refinement

All H atoms were placed in calculated positions with C—H = 0.93 Å (aromatic), C—H = 0.97 Å (methylene), and treated as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title compound, showing the atomic numbering and displacement ellipsoids at the 30% probability level [symmetry code: (i) $-x, 1 - y, -z$].




Fig. 2. A portion of the crystal packing showing the two-dimensional hydrogen bonded (dashed lines) sheet.

1,4-Bis(pyridin-3-ylmethoxy)benzene

Crystal data

$C_{18}H_{16}N_2O_2$	$F_{000} = 308$
$M_r = 292.33$	$D_x = 1.321 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4889 reflections
$a = 6.852 (5) \text{ \AA}$	$\theta = 3.7\text{--}27.5^\circ$
$b = 5.688 (3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 18.861 (12) \text{ \AA}$	$T = 291 \text{ K}$
$\beta = 90.60 (3)^\circ$	Block, colourless
$V = 735.0 (8) \text{ \AA}^3$	$0.22 \times 0.20 \times 0.19 \text{ mm}$
$Z = 2$	

Data collection

Rigaku RAXIS-RAPID diffractometer	1684 independent reflections
Radiation source: fine-focus sealed tube	1213 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.030$
$T = 291 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scan	$\theta_{\text{min}} = 3.7^\circ$
Absorption correction: multi-scan (ABSCOR; Higashi, 1995)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.981$, $T_{\text{max}} = 0.984$	$k = -6 \rightarrow 7$
6855 measured reflections	$l = -24 \rightarrow 24$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.041$	H-atom parameters constrained
$wR(F^2) = 0.118$	$w = 1/[\sigma^2(F_o^2) + (0.0606P)^2 + 0.0635P]$
$S = 1.09$	where $P = (F_o^2 + 2F_c^2)/3$
1684 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
100 parameters	$\Delta\rho_{\text{max}} = 0.24 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\text{min}} = -0.14 \text{ e \AA}^{-3}$
	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations

between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.8460 (2)	-0.0288 (3)	0.20865 (8)	0.0485 (4)
H1	0.9650	-0.0611	0.2307	0.058*
C2	0.7721 (2)	-0.1906 (3)	0.16191 (8)	0.0530 (4)
H2	0.8404	-0.3281	0.1524	0.064*
C3	0.5951 (2)	-0.1469 (3)	0.12921 (8)	0.0484 (4)
H3	0.5422	-0.2545	0.0972	0.058*
C4	0.49744 (18)	0.0588 (2)	0.14460 (7)	0.0367 (3)
C5	0.5857 (2)	0.2119 (3)	0.19208 (7)	0.0428 (4)
H5	0.5213	0.3517	0.2022	0.051*
C6	0.30072 (19)	0.1158 (3)	0.11378 (7)	0.0433 (4)
H6A	0.2437	-0.0233	0.0922	0.052*
H6B	0.2144	0.1704	0.1507	0.052*
C7	0.15874 (17)	0.3913 (2)	0.03246 (7)	0.0368 (3)
C8	-0.02824 (18)	0.3034 (3)	0.04198 (7)	0.0408 (3)
H8	-0.0477	0.1710	0.0700	0.049*
C9	-0.18583 (18)	0.4142 (3)	0.00955 (7)	0.0408 (3)
H9	-0.3112	0.3563	0.0163	0.049*
N1	0.75686 (17)	0.1729 (2)	0.22431 (7)	0.0510 (4)
O1	0.32526 (13)	0.29466 (19)	0.06180 (5)	0.0509 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0308 (7)	0.0572 (9)	0.0571 (9)	0.0002 (6)	-0.0101 (6)	0.0171 (7)
C2	0.0439 (9)	0.0449 (8)	0.0700 (10)	0.0117 (7)	-0.0063 (7)	0.0064 (8)
C3	0.0487 (8)	0.0432 (8)	0.0532 (8)	0.0035 (7)	-0.0108 (6)	-0.0038 (7)
C4	0.0308 (6)	0.0395 (7)	0.0396 (7)	-0.0005 (5)	-0.0056 (5)	0.0083 (6)
C5	0.0371 (7)	0.0388 (7)	0.0525 (8)	0.0016 (6)	-0.0069 (6)	0.0008 (6)
C6	0.0334 (7)	0.0482 (8)	0.0482 (8)	-0.0010 (6)	-0.0093 (6)	0.0109 (6)
C7	0.0264 (6)	0.0465 (8)	0.0374 (6)	0.0056 (5)	-0.0046 (5)	0.0038 (6)
C8	0.0308 (7)	0.0475 (8)	0.0438 (7)	-0.0007 (6)	-0.0045 (5)	0.0111 (6)
C9	0.0250 (6)	0.0530 (8)	0.0444 (7)	-0.0016 (6)	-0.0030 (5)	0.0073 (6)
N1	0.0394 (7)	0.0538 (8)	0.0596 (8)	-0.0056 (6)	-0.0145 (6)	0.0004 (6)
O1	0.0269 (5)	0.0683 (7)	0.0573 (6)	0.0058 (4)	-0.0048 (4)	0.0267 (5)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.334 (2)	C6—O1	1.4239 (17)
-------	-----------	-------	-------------

supplementary materials

C1—C2	1.368 (2)	C6—H6A	0.9700
C1—H1	0.9300	C6—H6B	0.9700
C2—C3	1.378 (2)	C7—C9 ⁱ	1.374 (2)
C2—H2	0.9300	C7—O1	1.3773 (17)
C3—C4	1.380 (2)	C7—C8	1.389 (2)
C3—H3	0.9300	C8—C9	1.3869 (19)
C4—C5	1.3839 (19)	C8—H8	0.9300
C4—C6	1.4979 (19)	C9—C7 ⁱ	1.374 (2)
C5—N1	1.3336 (19)	C9—H9	0.9300
C5—H5	0.9300		
N1—C1—C2	123.64 (13)	O1—C6—H6A	110.1
N1—C1—H1	118.2	C4—C6—H6A	110.1
C2—C1—H1	118.2	O1—C6—H6B	110.1
C1—C2—C3	119.03 (14)	C4—C6—H6B	110.1
C1—C2—H2	120.5	H6A—C6—H6B	108.4
C3—C2—H2	120.5	C9 ⁱ —C7—O1	115.88 (11)
C2—C3—C4	119.04 (14)	C9 ⁱ —C7—C8	119.64 (12)
C2—C3—H3	120.5	O1—C7—C8	124.47 (13)
C4—C3—H3	120.5	C9—C8—C7	119.64 (14)
C3—C4—C5	117.37 (13)	C9—C8—H8	120.2
C3—C4—C6	122.56 (13)	C7—C8—H8	120.2
C5—C4—C6	120.04 (13)	C7 ⁱ —C9—C8	120.71 (12)
N1—C5—C4	124.50 (14)	C7 ⁱ —C9—H9	119.6
N1—C5—H5	117.7	C8—C9—H9	119.6
C4—C5—H5	117.7	C5—N1—C1	116.41 (13)
O1—C6—C4	108.04 (11)	C7—O1—C6	117.28 (10)

Symmetry codes: (i) $-x, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1 \cdots N1 ⁱⁱ	0.93	2.57	3.437 (3)	155

Symmetry codes: (ii) $-x+2, y-1/2, -z+1/2$.

Fig. 1



Fig. 2

