

(E)-1-Ethyl-4-oxo-N'-(4-pyridylmethyl-ene)-1,4-dihydroquinoline-3-carbohydrazide

Fernanda da C. Santos,^a Pedro Netto Batalha,^a Anna Claudia Cunha,^a Rafael A. Alão,^a Vitor Francisco Ferreira,^a Maria Cecilia B. V. de Souza^a and Sauli Santos Jr^{b*}

^aInstituto de Química, Universidade Federal Fluminense, Niterói - Rio de Janeiro, Brazil, and ^bCoordenação de Física, Campus de Jataí, Universidade Federal de Goiás, Jataí - Goiás, Brazil

Correspondence e-mail: sauli@pq.cnpq.br


Received 15 July 2009; accepted 9 September 2009

Key indicators: single-crystal X-ray study; *T* = 293 K; mean $\sigma(\text{C}-\text{C}) = 0.002 \text{ \AA}$; *R* factor = 0.041; *wR* factor = 0.102; data-to-parameter ratio = 11.6.

In the title compound, C₁₈H₁₆N₄O₂, the plane defined by the ethyl C atoms and the attached N atom is inclined to the adjacent pyridine ring at an angle of 67.87 (16)°. The dihedral angle between the two heterocyclic rings is 3.33 (16)°. The molecular conformation is stabilized by an intramolecular N—H···O hydrogen bond and the crystal structure by intermolecular C—H···O hydrogen bonds, forming a one-dimensional structure.

Related literature

For the biological properties of oxoquinoline derivatives, see: Van Bambeke *et al.* (2005); Canuto *et al.* (2007); Lucero *et al.* (2006). For their potential use in the treatment of fungal and viral infections, see: Brideau *et al.* (2002); Souza *et al.* (2008) and in cancer chemotherapy, see: Chu *et al.* (1992). For acylhydrazones and their antileishmanial activity, see: Bernardino *et al.* (2006); Cunha *et al.* (2003). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

C₁₈H₁₆N₄O₂
M_r = 320.35
 Monoclinic, *P*2₁/*n*
a = 7.6460 (12) Å
b = 19.205 (2) Å
c = 10.7050 (9) Å
 β = 99.722 (10)°
V = 1549.4 (3) Å³
Z = 4
 Mo *K*α radiation
 μ = 0.09 mm⁻¹
T = 293 K
 0.5 × 0.2 × 0.2 mm

Data collection

Nonius KappaCCD diffractometer
 Absorption correction: none
 7914 measured reflections
 2586 independent reflections
 2001 reflections with *I* > 2σ(*I*)
*R*_{int} = 0.019

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.102$
S = 1.07
 2586 reflections
 222 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.13 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.15 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N3—H3N···O1	0.92 (2)	1.85 (2)	2.639 (2)	143.1 (17)
C10—H10A···O2 ⁱ	0.97	2.46	3.398 (2)	163

Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: COLLECT (Nonius, 2000); cell refinement: HKL SCALEPACK (Otwinowski & Minor 1997); data reduction: HKL DENZO (Otwinowski & Minor 1997) and SCALEPACK; program(s) used to solve structure: SIR2002 (Burla *et al.*, 2003); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2008); software used to prepare material for publication: WinGX publication routines (Farrugia, 1999).

The authors acknowledge the financial support from FAPERJ, CAPES and CNPq; they also thank Dra Maria G.F. Vaz, Marilene M. do Canto-Cavaleiro and Vanessa P. Rodrigues for useful discussions.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2340).

References

Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
 Bernardino, A. M. R., Gomes, A. O., Charret, K. S., Freitas, A. C. C., Machado, G. M. C., Canto-Cavaleiro, M. M., Leon, L. L. & Amaral, V. F. (2006). *Eur. J. Med. Chem.* **41**, 80–87.
 Brideau, R. J., Knechtel, M. L., Huang, A., Vaillancourt, V. A., Vera, E. E., Oien, N. L., Hopkins, T. A., Wieber, J. L., Wilkinson, K. F., Rush, B. D., Schwende, F. J. & Wathen, M. W. (2002). *Antiviral Res.* **54**, 19–28.
 Burla, M. C., Camalli, M., Carrozzini, B., Cascarano, G. L., Giacovazzo, C., Polidori, G. & Spagna, R. (2003). *J. Appl. Cryst.* **36**, 1103.
 Canuto, C. V. B. S., Gomes, C. R. B., Marques, I. P., Faro, L. V., Santos, F. C., Frugulhetti, I. C. P. P., Souza, T. M. L., Cunha, A. C., Romeiro, G. A., Ferreira, V. F. & Souza, M. C. B. (2007). *Lett. Drug Des. Discov.* **4**, 404–409.
 Chu, D. T., Hallas, R., Clement, J. J., Alder, J., McDonald, E. & Plattner, J. J. (1992). *Drugs Exp. Clin. Res.* **18**, 275–282.

- Cunha, A. C., Figueiredo, J. M., Tributino, J. L. M., Miranda, A. L. P., Castro, H. C., Zingali, R. B., Fraga, C. A. M., Souza, M. C. B. V., Ferreira, V. F. & Barreiro, E. J. (2003). *Bioorg. Med. Chem.* **11**, 2051–2059.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Lucero, B. A., Gomes, C. R. B., Frugulhetti, I. C. P. P., Faro, L. V., Alvarenga, L., Souza, M. C. B. V., Souza, T. M. L. & Ferreira, V. F. (2006). *Bioorg. Med. Chem. Lett.* **16**, 1010–1013.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Souza, T. M. L., Cirne-Santos, C. C., Rodrigues, D. Q., Abreu, C. M., Tanuri, A., Ferreira, V. F., Marques, I. P., Souza, M. C. B. V., Fontes, C. F. L., Frugulhetti, I. C. P. P. & Bou-Habib, D. C. (2008). *Curr. HIV Res.* **6**, 209–217.
- Van Bambeke, F., Michot, J. M., Van Eldere, J. & Tulkens, P. M. (2005). *Clin. Microbiol. Infect.* **11**, 256–280.

supplementary materials

Acta Cryst. (2009). E65, o2476-o2477 [doi:10.1107/S160053680903654X]

(E)-1-Ethyl-4-oxo-*N'*-(4-pyridylmethylene)-1,4-dihydroquinoline-3-carbohydrazide

F. da C. Santos, P. N. Batalha, A. C. Cunha, R. A. Alão, V. F. Ferreira, M. C. B. V. de Souza and S. Santos Jr

Comment

Since the discovery of nalidixic acid, the parent compound of the 1,4-dihydro-4-oxoquinoline antibiotics, the molecular structures of oxoquinolines have been extensively modified to improve their pharmacological properties and pharmacokinetic profiles (Van Bambeke et al., 2005). 1,4-Dihydro-4-oxoquinolines have a broad antimicrobial spectrum, are orally and parenterally active and, apart from a few exceptions, are non-toxic compounds. Therefore, they are important agents against microbial pathogens. They have also been considered for the treatment of fungal and viral (Brideau et al., 2002; Souza et al., 2008) infections and for cancer chemotherapy (Chu et al., 1992). Our research group has designed and synthesized new oxoquinoline derivatives, such as ribonucleosides (Canuto et al., 2007) and acyclonucleosides (Lucero et al., 2006). These compounds exhibited interesting activity against HIV-1 and HSV viruses, respectively.

Continuing our interest in the chemistry of oxoquinolinic molecules, we decided to prepare a congener series of new acylhydrazones (Cunha et al., 2003) and test them for antileishmanial activity. These studies showed that the hydrazone group plays an important role in antileishmanial activity (Bernadino et al., 2006). Among these compounds, (E)-1-ethyl-*N'*-[(pyrid-4'-yl)methylene]-1,4-dihydro-4-oxoquinoline-3-carbohydrazide, was found to exhibit a significant activity against *Leishmania amazonensis*.

As an extension of our work on the structural characterization of oxoquinolinic derivatives (Canuto et al., 2007; Lucero et al. 2006), the crystal structure of the title compound is reported here. In the title compound (Fig. 1), the dihedral angle between the C1,C6-C9/N1 ring and the C14-C18/N4 pyridine ring is 3.33 (16)°. The plane defined by N1,C10,C11 is inclined to the C1,C6-C9/N1 ring at an angle of 67.87 (16)°. The torsion angles C8—C12—N3—N2, C12—N3—N2—C13 and N3—N2—C13—C14 are 0.79 (14), 1.38 (16) and 1.53 (14)°, respectively. All the bond lengths are within normal ranges (Allen et al., 1987). The molecular structure is stabilized by an intramolecular N—H···O hydrogen bond (Table 1). The crystal structure is stabilized by intermolecular C—H···O hydrogen bonds, forming a one-dimensional structure (Table 1 and Fig. 2).

Experimental

A solution of 1-ethyl-1,4-dihydro-4-oxoquinoline ethyl carboxylate (3.70 mmol) and 3.7 ml of 80% hydrazine monohydrate in 10 ml of dimethylformamide was stirred under reflux for two hours. The reaction mixture was poured into ice, giving rise to a white solid that was collected by filtration, washed with cold ethyl acetate and dried under vacuum, leading to the pure desired 1-ethyl-1,4-dihydro-4-oxoquinoline-3-carbohydrazide, in 90% yield. This carbohydrazide (1.00 mmol) and pyridine-4-carbaldehyde (1.10 mmol) in 5 ml of dimethylformamide and a catalytic amount of 35% HCl were stirred at room temperature. The reaction mixture was poured into ice, leading to a white solid that was collected by filtration, giving (E)-1-ethyl-*N'*-[(pyrid-4'-yl)methylene]-1,4-dihydro-4-oxoquinoline-3-carbohydrazide in 83% yield after purification.

Refinement

The N-bound H atom was located in a difference map and refined freely [to N—H = 0.92 (2) Å]. The other H atoms were positioned with idealized geometry and refined using a riding model, with C—H = 0.93 Å for aryl, 0.97 Å for methylene, and 0.96 Å for methyl H atoms. The constraints $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and $1.5U_{\text{eq}}(\text{C})$ (methyl C) were applied.

Figures




Fig. 1. The molecular structure of the title compound showing the atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen atoms are shown as spheres of arbitrary radius.




Fig. 2. The molecular packing, with hydrogen bonds drawn as dashed lines.

(E)-1-Ethyl-4-oxo-N'-(4-pyridylmethylene)-1,4-dihydroquinoline-3-carbohydrazide

Crystal data

$\text{C}_{18}\text{H}_{16}\text{N}_4\text{O}_2$

$M_r = 320.35$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 7.6460$ (12) Å

$b = 19.205$ (2) Å

$c = 10.7050$ (9) Å

$\beta = 99.722$ (10)°

$V = 1549.4$ (3) Å³

$Z = 4$

$F_{000} = 672$

$D_x = 1.373$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 7914 reflections

$\theta = 5.2$ – 25 °

$\mu = 0.09$ mm⁻¹

$T = 293$ K

Prism, yellow

$0.5 \times 0.2 \times 0.2$ mm

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 9 pixels mm⁻¹

$T = 293$ K

2586 independent reflections

2001 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.019$

$\theta_{\text{max}} = 25.0$ °

$\theta_{\text{min}} = 5.2$ °

φ scans and ω scans with κ offsets $h = -9 \rightarrow 9$
 Absorption correction: none $k = -22 \rightarrow 20$
 7914 measured reflections $l = -12 \rightarrow 8$

Refinement

Refinement on F^2 Secondary atom site location: difference Fourier map
 Least-squares matrix: full Hydrogen site location: inferred from neighbouring sites
 $R[F^2 > 2\sigma(F^2)] = 0.041$ H atoms treated by a mixture of independent and constrained refinement
 $wR(F^2) = 0.102$ $w = 1/[\sigma^2(F_o^2) + (0.0352P)^2 + 0.5415P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $S = 1.07$ $(\Delta/\sigma)_{\max} < 0.001$
 2586 reflections $\Delta\rho_{\max} = 0.13 \text{ e } \text{\AA}^{-3}$
 222 parameters $\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$
 Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.9168 (2)	0.11495 (9)	0.16391 (15)	0.0361 (4)
C2	0.8969 (2)	0.07582 (10)	0.05163 (16)	0.0455 (4)
H2	0.8539	0.0305	0.0514	0.055*
C3	0.9396 (3)	0.10304 (10)	-0.05772 (17)	0.0516 (5)
H3	0.9240	0.0768	-0.1318	0.062*
C4	1.0065 (3)	0.17027 (11)	-0.05650 (17)	0.0521 (5)
H4	1.0374	0.1886	-0.1302	0.063*
C5	1.0280 (2)	0.21026 (10)	0.05098 (16)	0.0471 (4)
H5	1.0740	0.2550	0.0500	0.056*
C6	0.9804 (2)	0.18344 (9)	0.16269 (15)	0.0374 (4)
C7	0.8708 (2)	0.08354 (9)	0.27920 (15)	0.0371 (4)
C8	0.8931 (2)	0.12883 (8)	0.38879 (15)	0.0365 (4)
C9	0.9529 (2)	0.19544 (9)	0.37882 (15)	0.0401 (4)
H9	0.9638	0.2235	0.4505	0.048*
C10	1.0520 (3)	0.29759 (10)	0.27524 (18)	0.0538 (5)
H10A	1.1500	0.3026	0.2291	0.065*
H10B	1.0931	0.3118	0.3622	0.065*
C11	0.9018 (3)	0.34469 (11)	0.2170 (2)	0.0750 (7)
H11A	0.9448	0.3915	0.2143	0.113*

supplementary materials

H11B	0.8090	0.3432	0.2671	0.113*
H11C	0.8563	0.3293	0.1324	0.113*
C12	0.8472 (2)	0.10900 (9)	0.51400 (16)	0.0416 (4)
C13	0.6773 (2)	-0.04576 (9)	0.61567 (16)	0.0429 (4)
H13	0.6722	-0.0691	0.5391	0.051*
C14	0.6223 (2)	-0.08221 (9)	0.72279 (15)	0.0388 (4)
C15	0.5717 (2)	-0.15131 (9)	0.71190 (17)	0.0494 (5)
H15	0.5690	-0.1747	0.6356	0.059*
C16	0.5249 (3)	-0.18542 (10)	0.81527 (19)	0.0544 (5)
H16	0.4916	-0.2319	0.8057	0.065*
C17	0.5729 (3)	-0.08884 (11)	0.93622 (18)	0.0551 (5)
H17	0.5739	-0.0667	1.0136	0.066*
C18	0.6209 (2)	-0.05045 (10)	0.83932 (17)	0.0479 (4)
H18	0.6521	-0.0038	0.8514	0.057*
N1	0.99728 (18)	0.22334 (7)	0.27329 (12)	0.0402 (4)
N2	0.73182 (18)	0.01676 (7)	0.62429 (13)	0.0419 (4)
N3	0.7828 (2)	0.04327 (8)	0.51697 (14)	0.0445 (4)
N4	0.5248 (2)	-0.15582 (9)	0.92715 (15)	0.0541 (4)
O1	0.81667 (17)	0.02200 (6)	0.27864 (11)	0.0522 (3)
O2	0.8639 (2)	0.14902 (7)	0.60464 (12)	0.0636 (4)
H3N	0.776 (2)	0.0178 (10)	0.4436 (19)	0.057 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0335 (8)	0.0403 (10)	0.0342 (9)	0.0054 (7)	0.0048 (7)	-0.0024 (7)
C2	0.0512 (10)	0.0443 (10)	0.0408 (10)	0.0051 (8)	0.0074 (8)	-0.0066 (8)
C3	0.0632 (12)	0.0569 (12)	0.0346 (10)	0.0102 (10)	0.0078 (8)	-0.0070 (8)
C4	0.0603 (12)	0.0623 (13)	0.0366 (10)	0.0074 (10)	0.0163 (8)	0.0047 (9)
C5	0.0523 (11)	0.0506 (11)	0.0409 (10)	0.0003 (9)	0.0151 (8)	0.0035 (8)
C6	0.0360 (9)	0.0428 (10)	0.0343 (9)	0.0027 (7)	0.0088 (7)	-0.0018 (7)
C7	0.0364 (9)	0.0365 (9)	0.0388 (9)	0.0030 (7)	0.0075 (7)	-0.0011 (7)
C8	0.0391 (9)	0.0362 (9)	0.0354 (9)	0.0013 (7)	0.0101 (7)	-0.0018 (7)
C9	0.0456 (10)	0.0427 (10)	0.0342 (9)	-0.0017 (8)	0.0132 (7)	-0.0055 (7)
C10	0.0727 (13)	0.0478 (11)	0.0452 (10)	-0.0231 (10)	0.0227 (9)	-0.0063 (9)
C11	0.114 (2)	0.0441 (12)	0.0719 (15)	-0.0014 (13)	0.0312 (14)	0.0043 (11)
C12	0.0491 (10)	0.0379 (10)	0.0403 (10)	0.0021 (8)	0.0144 (8)	-0.0024 (8)
C13	0.0520 (10)	0.0398 (10)	0.0368 (9)	-0.0028 (8)	0.0071 (8)	-0.0011 (7)
C14	0.0380 (9)	0.0378 (9)	0.0394 (9)	0.0020 (7)	0.0029 (7)	0.0047 (7)
C15	0.0611 (12)	0.0424 (10)	0.0440 (10)	-0.0037 (9)	0.0066 (9)	0.0000 (8)
C16	0.0643 (13)	0.0409 (11)	0.0573 (12)	-0.0057 (9)	0.0084 (10)	0.0082 (9)
C17	0.0720 (13)	0.0519 (12)	0.0446 (11)	0.0025 (10)	0.0191 (9)	0.0019 (9)
C18	0.0584 (11)	0.0402 (10)	0.0457 (10)	-0.0006 (8)	0.0110 (8)	-0.0003 (8)
N1	0.0474 (8)	0.0392 (8)	0.0361 (8)	-0.0068 (6)	0.0128 (6)	-0.0033 (6)
N2	0.0500 (9)	0.0395 (9)	0.0381 (8)	-0.0008 (7)	0.0125 (6)	0.0029 (6)
N3	0.0613 (10)	0.0397 (9)	0.0353 (8)	-0.0056 (7)	0.0158 (7)	0.0001 (7)
N4	0.0598 (10)	0.0525 (10)	0.0521 (10)	0.0032 (8)	0.0149 (8)	0.0113 (8)
O1	0.0732 (9)	0.0387 (7)	0.0464 (7)	-0.0095 (6)	0.0147 (6)	-0.0056 (5)

O2 0.1082 (11) 0.0441 (8) 0.0471 (8) -0.0128 (7) 0.0379 (7) -0.0102 (6)

Geometric parameters (Å, °)

C1—C6	1.403 (2)	C10—H10B	0.9700
C1—C2	1.404 (2)	C11—H11A	0.9600
C1—C7	1.469 (2)	C11—H11B	0.9600
C2—C3	1.371 (2)	C11—H11C	0.9600
C2—H2	0.9300	C12—O2	1.227 (2)
C3—C4	1.388 (3)	C12—N3	1.357 (2)
C3—H3	0.9300	C13—N2	1.269 (2)
C4—C5	1.370 (2)	C13—C14	1.465 (2)
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.405 (2)	C14—C15	1.382 (2)
C5—H5	0.9300	C14—C18	1.390 (2)
C6—N1	1.398 (2)	C15—C16	1.384 (3)
C7—O1	1.2518 (19)	C15—H15	0.9300
C7—C8	1.447 (2)	C16—N4	1.326 (2)
C8—C9	1.369 (2)	C16—H16	0.9300
C8—C12	1.491 (2)	C17—N4	1.337 (2)
C9—N1	1.344 (2)	C17—C18	1.372 (3)
C9—H9	0.9300	C17—H17	0.9300
C10—N1	1.485 (2)	C18—H18	0.9300
C10—C11	1.511 (3)	N2—N3	1.3719 (19)
C10—H10A	0.9700	N3—H3N	0.92 (2)
C6—C1—C2	118.79 (15)	C10—C11—H11B	109.5
C6—C1—C7	121.70 (14)	H11A—C11—H11B	109.5
C2—C1—C7	119.52 (15)	C10—C11—H11C	109.5
C3—C2—C1	121.32 (17)	H11A—C11—H11C	109.5
C3—C2—H2	119.3	H11B—C11—H11C	109.5
C1—C2—H2	119.3	O2—C12—N3	123.61 (16)
C2—C3—C4	119.16 (17)	O2—C12—C8	122.81 (15)
C2—C3—H3	120.4	N3—C12—C8	113.58 (14)
C4—C3—H3	120.4	N2—C13—C14	121.97 (15)
C5—C4—C3	121.40 (17)	N2—C13—H13	119.0
C5—C4—H4	119.3	C14—C13—H13	119.0
C3—C4—H4	119.3	C15—C14—C18	116.92 (16)
C4—C5—C6	119.85 (17)	C15—C14—C13	120.43 (16)
C4—C5—H5	120.1	C18—C14—C13	122.65 (16)
C6—C5—H5	120.1	C14—C15—C16	119.58 (17)
N1—C6—C1	119.25 (14)	C14—C15—H15	120.2
N1—C6—C5	121.31 (16)	C16—C15—H15	120.2
C1—C6—C5	119.44 (15)	N4—C16—C15	123.99 (18)
O1—C7—C8	124.40 (15)	N4—C16—H16	118.0
O1—C7—C1	120.64 (14)	C15—C16—H16	118.0
C8—C7—C1	114.96 (14)	N4—C17—C18	124.58 (18)
C9—C8—C7	119.59 (14)	N4—C17—H17	117.7
C9—C8—C12	116.19 (14)	C18—C17—H17	117.7
C7—C8—C12	124.17 (15)	C17—C18—C14	119.08 (18)

supplementary materials

N1—C9—C8	124.96 (15)	C17—C18—H18	120.5
N1—C9—H9	117.5	C14—C18—H18	120.5
C8—C9—H9	117.5	C9—N1—C6	119.53 (14)
N1—C10—C11	112.13 (16)	C9—N1—C10	118.75 (14)
N1—C10—H10A	109.2	C6—N1—C10	121.59 (13)
C11—C10—H10A	109.2	C13—N2—N3	115.25 (14)
N1—C10—H10B	109.2	C12—N3—N2	121.35 (15)
C11—C10—H10B	109.2	C12—N3—H3N	116.4 (12)
H10A—C10—H10B	107.9	N2—N3—H3N	122.2 (12)
C10—C11—H11A	109.5	C16—N4—C17	115.84 (16)
C6—C1—C2—C3	0.6 (2)	C6—C1—C2—H2	-179.4
C7—C1—C2—C3	-179.53 (16)	C7—C1—C2—H2	0.5
C1—C2—C3—C4	1.0 (3)	C1—C2—C3—H3	-179.0
C2—C3—C4—C5	-1.1 (3)	H2—C2—C3—H3	1.0
C3—C4—C5—C6	-0.6 (3)	H2—C2—C3—C4	-179.0
C2—C1—C6—N1	178.70 (14)	C2—C3—C4—H4	178.9
C7—C1—C6—N1	-1.2 (2)	H3—C3—C4—H4	-1.1
C2—C1—C6—C5	-2.2 (2)	H3—C3—C4—C5	178.9
C7—C1—C6—C5	177.95 (14)	C3—C4—C5—H5	179.4
C4—C5—C6—N1	-178.71 (16)	H4—C4—C5—H5	-0.6
C4—C5—C6—C1	2.2 (3)	H4—C4—C5—C6	179.4
C6—C1—C7—O1	-178.98 (15)	H5—C5—C6—C1	-177.8
C2—C1—C7—O1	1.1 (2)	H5—C5—C6—N1	1.3
C6—C1—C7—C8	1.0 (2)	C7—C8—C9—H9	178.8
C2—C1—C7—C8	-178.88 (14)	C12—C8—C9—H9	1.4
O1—C7—C8—C9	-179.87 (16)	H9—C9—N1—C10	-2.9
C1—C7—C8—C9	0.1 (2)	H9—C9—N1—C6	-178.9
O1—C7—C8—C12	-2.7 (3)	H10A—C10—C11—H11A	-54.6
C1—C7—C8—C12	177.27 (14)	H10A—C10—C11—H11B	-174.6
C7—C8—C9—N1	-1.2 (3)	H10A—C10—C11—H11C	65.4
C12—C8—C9—N1	-178.55 (15)	H10B—C10—C11—H11A	63.1
C9—C8—C12—O2	-1.1 (3)	H10B—C10—C11—H11B	-56.9
C7—C8—C12—O2	-178.36 (17)	H10B—C10—C11—H11C	-176.9
C9—C8—C12—N3	177.80 (15)	N1—C10—C11—H11A	-175.7
C7—C8—C12—N3	0.6 (2)	N1—C10—C11—H11B	64.3
N2—C13—C14—C15	-176.91 (16)	N1—C10—C11—H11C	-55.7
N2—C13—C14—C18	1.9 (3)	H10A—C10—N1—C6	-43.9
C18—C14—C15—C16	-0.8 (3)	H10A—C10—N1—C9	140.2
C13—C14—C15—C16	178.15 (16)	H10B—C10—N1—C6	-161.6
C14—C15—C16—N4	0.2 (3)	H10B—C10—N1—C9	22.5
N4—C17—C18—C14	-0.6 (3)	C8—C12—N3—H3N	1.0 (13)
C15—C14—C18—C17	0.9 (3)	H13—C13—C14—C15	3.1
C13—C14—C18—C17	-177.93 (17)	H13—C13—C14—C18	-178.1
C8—C9—N1—C6	1.1 (2)	H13—C13—N2—N3	-1.5
C8—C9—N1—C10	177.08 (16)	C13—C14—C15—H15	-1.8
C1—C6—N1—C9	0.2 (2)	C18—C14—C15—H15	179.2
C5—C6—N1—C9	-178.95 (15)	C13—C14—C18—H18	2.1
C1—C6—N1—C10	-175.74 (15)	C15—C14—C18—H18	-179.1
C5—C6—N1—C10	5.1 (2)	C14—C15—C16—H16	-179.8

C11—C10—N1—C9	-98.68 (19)	H15—C15—C16—H16	0.2
C11—C10—N1—C6	77.3 (2)	H15—C15—C16—N4	-179.8
C14—C13—N2—N3	178.47 (14)	H16—C16—N4—C17	-179.7
O2—C12—N3—N2	-0.3 (3)	H17—C17—C18—C14	179.4
C8—C12—N3—N2	-179.21 (14)	H17—C17—C18—H18	-0.6
C13—N2—N3—C12	-178.62 (16)	N4—C17—C18—H18	179.4
C15—C16—N4—C17	0.3 (3)	H17—C17—N4—C16	180.0
C18—C17—N4—C16	0.0 (3)	C13—N2—N3—H3N	1.2 (14)
C6—C1—C2—H2	-179.43 (1)	O2—C12—N3—H3N	179.9 (13)
C7—C1—C2—H2	0.46 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3—H3N \cdots O1	0.92 (2)	1.85 (2)	2.639 (2)	143.1 (17)
C10—H10A \cdots O2 ⁱ	0.97	2.46	3.398 (2)	163

Symmetry codes: (i) $x+1/2, -y+1/2, z-1/2$.

Fig. 1




Fig. 2

