

3-(1*H*-Imidazol-1-yl)propanenitrile

Tim Peppel and Martin Köckerling*

Universität Rostock, Institut für Chemie, Abteilung Anorganische Chemie/Festkörperchemie, Albert-Einstein-Strasse 3a, D-18059 Rostock, Germany
Correspondence e-mail: Martin.Koeckerling@uni-rostock.de


Received 18 August 2009; accepted 3 September 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.032; wR factor = 0.086; data-to-parameter ratio = 13.9.

The title compound, $\text{C}_6\text{H}_7\text{N}_3$, has an ethylene group connecting an imidazole ring and a $-\text{CN}$ group. These groups are in a staggered conformation. The shortest intermolecular contact is found between the imidazole N atom and a $-\text{CH}_2-$ group of a neighboring molecule.

Related literature

For background and applications of ionic liquids, see: Hayashi *et al.* (2006); Kozlova *et al.* (2009a,b); Lombardo *et al.* (2007); Macaev *et al.* (2007); Sawa & Okamura (1969); Scheers *et al.* (2008); Visser *et al.* (2001); Wang *et al.* (2003); Wasserscheid & Keim (2000); Xu *et al.* (2007); Yamauchi & Masui (1976); Yang *et al.* (2006).



Experimental

Crystal data

$\text{C}_6\text{H}_7\text{N}_3$

$M_r = 121.15$

Monoclinic, $P2_1/c$

$a = 7.2712 (3)\text{ \AA}$

$b = 5.5917 (2)\text{ \AA}$

$c = 15.4625 (5)\text{ \AA}$

$\beta = 100.979 (1)^\circ$

$V = 617.17 (4)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.09\text{ mm}^{-1}$

$T = 173\text{ K}$

$0.45 \times 0.40 \times 0.30\text{ mm}$

Data collection

Bruker-Nonius X8 APEX diffractometer

Absorption correction: multi-scan (*SADABS*; Bruker, 2007)

$T_{\min} = 0.946$, $T_{\max} = 0.975$

11604 measured reflections

1542 independent reflections

1456 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.018$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.086$

$S = 1.04$

1542 reflections

111 parameters

All H-atom parameters refined

$\Delta\rho_{\text{max}} = 0.27\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.16\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}5-\text{H}5\text{A}\cdots \text{N}2^1$	0.97 (1)	2.66 (1)	3.366 (1)	135.7 (9)
Symmetry code: (i) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.				

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 1999); software used to prepare material for publication: *ct.exe* (Köckerling, 1996).

We thank Professor Dr Helmut Reinke (University of Rostock/Germany, Institute of Chemistry) for maintaining the X-ray equipment. Support from the DFG (priority program SPP 1191-Ionic Liquids, KO-1616/4-1 and 1616/4-2) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2269).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2007). *SAINT*, *SMART* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Hayashi, S., Saha, S. & Hamaguchi, H. (2006). *IEEE Trans. Magn.* **42**, 12–14.
Köckerling, M. (1996). *ct.exe*. Universität Rostock, Germany.
Kozlova, S. A., Verevkin, S. P., Heintz, A., Peppel, T. & Köckerling, M. (2009a). *J. Chem. Eng. Data*, **54**, 1524–1528.
Kozlova, S. A., Verevkin, S. P., Heintz, A., Peppel, T. & Köckerling, M. (2009b). *J. Chem. Thermodyn.* **41**, 330–333.
Lombardo, M., Pasi, F., Trombini, C., Seddon, K. R. & Pittner, W. R. (2007). *Green Chem.* **9**, 321–322.
Macaev, F., Gavrilov, K., Muntyanu, V., Styngach, E., Vlad, L., Bets, L., Pogrebnai, S. & Barba, A. (2007). *Chem. Nat. Compd.* **43**, 136–139.
Sawa, N. & Okamura, S. (1969). *Nippon Kagaku Zasshi*, **90**, 704–707.
Scheers, J., Johansson, P. & Jacobsson, P. (2008). *J. Electrochem. Soc.* **155**, A628–A634.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Visser, A. E., Swatloski, R. P., Reichert, W. M., Mayton, R., Sheff, S., Wierzbicki, A., Davis, J. H. Jr & Rogers, R. D. (2001). *Chem. Commun.* pp. 135–136.
Wang, P., Zakeeruddin, S. M., Comte, P., Exnar, I. & Grätzel, M. (2003). *J. Am. Chem. Soc.* **125**, 1166–1167.
Wasserscheid, P. & Keim, W. (2000). *Angew. Chem. Int. Ed.* **39**, 3772–3789.
Xu, J.-M., Qian, C., Liu, B.-K., Wu, Q. & Lin, X.-F. (2007). *Tetrahedron*, **63**, 986–990.
Yamauchi, M. & Masui, M. (1976). *Chem. Pharm. Bull.* **24**, 1480–1484.
Yang, L., Xu, L.-W., Zhou, W., Li, L. & Xia, C.-G. (2006). *Tetrahedron Lett.* **47**, 7723–7726.

supplementary materials

Acta Cryst. (2009). E65, o2480 [doi:10.1107/S1600536809035685]

3-(1*H*-Imidazol-1-yl)propanenitrile

T. Peppel and M. Köckerling

Comment

Ionic Liquids (ILs) are more and more attracting remarkable attention as new solvents and reaction media for organic and inorganic synthesis and catalysis in order to replace volatile organic solvents (Wasserscheid & Keim, 2000). Imidazolium based ILs have emerged as leading candidates since they have very low vapor pressures, are moisture and air stable, and are highly solvating for both molecular and ionic species. Furthermore, ILs are finding use in separation processes (Visser *et al.*, 2001), in battery applications (Scheers *et al.*, 2008), as electrolytes in solar cells (Wang *et al.*, 2003), or as part of metal catalysts and magnetic liquids (Lombardo *et al.*, 2007; Hayashi *et al.*, 2006; Kozlova *et al.*, 2009a; Kozlova *et al.*, 2009b). Ionic Liquids and imidazole compounds incorporating nitrile functionalities have shown applications in heterocyclic and terpene chemistry (Macaev *et al.*, 2007; Yamauchi & Masui, 1976), in Michael addition or aza-Michael reactions (Xu *et al.*, 2007; Yang *et al.*, 2006).

In this context we present the molecular and single-crystal structure of 3-(1*H*-imidazol-1-yl)propanenitrile, an important precursor of nitrile functionalized ILs. Although all other sources report this compound to be a liquid at room temperature, there is one single indication in the literature that it is a solid melting at 33–35 °C (Sawa & Okamura, 1969). A freshly distilled sample crystallizes spontaneously within days, forming big block-shaped colourless crystals, which are barely deliquescent and melting at 37 °C.

All the bond lengths are within the expected ranges. The molecular structure of the title compound is shown in Figure 1. The C6–N3 bond length of 1.141 (1) Å indicates the triple-bond nitrile character. This nitrile group is connected through an ethylene group with the planar imidazolyl ring. The ethylene group has a staggered conformation with an N1–C4–C5–C6 torsion angle of -65.43 (9)°.

The shortest intermolecular contacts are found between one of the H atoms bonded at C5 and the N2 atom of the symmetry equivalent neighboring molecule. The H5A…N2[#] distance measures 2.66 (1) Å, the corresponding C5…N2[#] distance 3.366 (1) Å, and the C5–H5a…N2[#] angle 135.6° (symmetry code #: x, 1/2-y, z+1/2). Therefore the contact can be considered as a weak hydrogen bond. Figure 2 shows the orientation of two molecules, which are attached through this weak hydrogen bond.

In the crystal the molecules are arranged in rows, such that this weak hydrogen bond is oriented approximately parallel to the crystallographic c direction. Due to the c glide plane, every second row, stacked along a is shifted by c/2. The packing diagram, Figure 3, shows this assembly.

Experimental

1*H*-Imidazole (50.0 g, 0.7 mol) and acrylonitrile (117.0 g, 2.2 mol) were refluxed in 150 ml of ethanol overnight. Excess acrylonitrile and the solvent were evaporated and the residue was distilled in vacuum, yielding a colourless supercooled liquid, which crystallized at room temperature after several days. Yield: 75.0 g (84%), mp 37 °C.

supplementary materials

Refinement

The positions of the hydrogen atoms were located from a difference Fourier map and refined isotropically.

Figures




Fig. 1. Molecular structure of 3-(1*H*-imidazol-1-yl)propanenitrile with atom labelling scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Two neighboring C₆H₇N₃ molecules with the short N2···H(C5) contact.




Fig. 3. Packing diagram of 3-(1*H*-imidazol-1-yl)propanenitrile in a view down the crystallographic *b* direction.

3-(1*H*-Imidazol-1-yl)propanenitrile

Crystal data

C ₆ H ₇ N ₃	$D_x = 1.304 \text{ Mg m}^{-3}$
$M_r = 121.15$	Melting point: 310 K
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.2712 (3) \text{ \AA}$	Cell parameters from 8906 reflections
$b = 5.5917 (2) \text{ \AA}$	$\theta = 2.7\text{--}28.3^\circ$
$c = 15.4625 (5) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\beta = 100.979 (1)^\circ$	$T = 173 \text{ K}$
$V = 617.17 (4) \text{ \AA}^3$	Block, colourless
$Z = 4$	$0.45 \times 0.40 \times 0.30 \text{ mm}$
$F_{000} = 256$	

Data collection

Bruker–Nonius X8 Apex diffractometer	1542 independent reflections
Radiation source: fine-focus sealed tube	1456 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.018$
$T = 173 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 3.5^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2007)	$h = -9\text{--}9$

$T_{\min} = 0.946$, $T_{\max} = 0.975$
11604 measured reflections

$k = -7 \rightarrow 7$
 $l = -20 \rightarrow 16$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	All H-atom parameters refined
$R[F^2 > 2\sigma(F^2)] = 0.032$	$w = 1/[\sigma^2(F_o^2) + (0.0486P)^2 + 0.1206P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.086$	$(\Delta/\sigma)_{\max} < 0.001$
$S = 1.04$	$\Delta\rho_{\max} = 0.27 \text{ e \AA}^{-3}$
1542 reflections	$\Delta\rho_{\min} = -0.16 \text{ e \AA}^{-3}$
111 parameters	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Primary atom site location: structure-invariant direct methods	Extinction coefficient: 0.064 (15)
Secondary atom site location: difference Fourier map	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.76561 (9)	0.2518 (1)	0.30688 (4)	0.0207 (2)
C1	0.6541 (1)	0.1964 (2)	0.22892 (5)	0.0254 (2)
H1	0.579 (2)	0.053 (2)	0.2217 (7)	0.033 (3)*
N2	0.6623 (1)	0.3580 (1)	0.16800 (4)	0.0302 (2)
C2	0.7866 (1)	0.5266 (2)	0.20919 (6)	0.0281 (2)
H2	0.817 (2)	0.670 (2)	0.1771 (8)	0.037 (3)*
C3	0.8519 (1)	0.4648 (1)	0.29465 (5)	0.0249 (2)
H3	0.944 (2)	0.535 (2)	0.3408 (8)	0.035 (3)*
C4	0.7876 (1)	0.1115 (1)	0.38757 (5)	0.0239 (2)
H4A	0.919 (2)	0.109 (2)	0.4147 (7)	0.030 (3)*
H4B	0.744 (2)	-0.051 (2)	0.3700 (7)	0.031 (3)*
C5	0.6708 (1)	0.2069 (2)	0.45232 (5)	0.0254 (2)
H5A	0.683 (2)	0.100 (2)	0.5022 (8)	0.037 (3)*
H5B	0.538 (2)	0.217 (2)	0.4243 (7)	0.035 (3)*

supplementary materials

C6	0.7298 (1)	0.4436 (2)	0.48692 (5)	0.0244 (2)
N3	0.7779 (1)	0.6275 (1)	0.51425 (5)	0.0333 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0242 (3)	0.0212 (3)	0.0176 (3)	0.0012 (2)	0.0061 (2)	0.0007 (2)
C1	0.0293 (4)	0.0279 (4)	0.0194 (4)	0.0001 (3)	0.0057 (3)	-0.0026 (3)
N2	0.0345 (4)	0.0371 (4)	0.0196 (3)	0.0033 (3)	0.0064 (3)	0.0029 (3)
C2	0.0290 (4)	0.0307 (4)	0.0271 (4)	0.0027 (3)	0.0113 (3)	0.0075 (3)
C3	0.0246 (4)	0.0249 (4)	0.0257 (4)	-0.0015 (3)	0.0065 (3)	0.0023 (3)
C4	0.0318 (4)	0.0213 (4)	0.0188 (3)	0.0026 (3)	0.0058 (3)	0.0025 (3)
C5	0.0321 (4)	0.0265 (4)	0.0190 (4)	-0.0056 (3)	0.0084 (3)	-0.0011 (3)
C6	0.0265 (4)	0.0288 (4)	0.0186 (3)	0.0011 (3)	0.0060 (3)	-0.0002 (3)
N3	0.0386 (4)	0.0303 (4)	0.0313 (4)	-0.0002 (3)	0.0078 (3)	-0.0047 (3)

Geometric parameters (\AA , $^\circ$)


N1—C1	1.354 (1)	C6—N3	1.141 (1)
N1—C3	1.376 (1)	C1—H1	0.97 (1)
N1—C4	1.4565 (9)	C2—H2	0.99 (1)
C1—N2	1.315 (1)	C3—H3	0.97 (1)
N2—C2	1.376 (1)	C4—H4A	0.97 (1)
C2—C3	1.361 (1)	C4—H4B	0.99 (1)
C4—C5	1.527 (1)	C5—H5A	0.97 (1)
C5—C6	1.461 (1)	C5—H5B	0.98 (1)
C1—N1—C3	106.68 (6)	H2—C2—N2	120.9 (7)
C1—N1—C4	126.06 (7)	H3—C3—C2	132.8 (7)
C3—N1—C4	127.27 (7)	H3—C3—N1	121.4 (7)
N2—C1—N1	112.30 (7)	H4A—C4—H4B	110.4 (9)
C1—N2—C2	104.76 (7)	H4A—C4—N1	108.5 (7)
C3—C2—N2	110.64 (7)	H4B—C4—N1	106.5 (7)
C2—C3—N1	105.63 (7)	H4A—C4—C5	110.2 (7)
N1—C4—C5	112.92 (6)	H4B—C4—C5	108.2 (7)
C6—C5—C4	113.28 (7)	H5A—C5—H5B	109 (1)
N3—C6—C5	179.24 (9)	H5A—C5—C6	107.0 (7)
H1—C1—N2	126.2 (7)	H5B—C5—C6	107.9 (7)
H1—C1—N1	121.5 (7)	H5A—C5—C4	109.2 (7)
H2—C2—C3	128.4 (7)	H5B—C5—C4	110.7 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
C5—H5A—N2 ⁱ	0.97 (1)	2.66 (1)	3.366 (1)	135.7 (9)

Symmetry codes: (i) $x, -y+1/2, z+1/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

