

Isopropyl 2-(5-fluoro-3-methylsulfinyl-1-benzofuran-2-yl)acetate

Hong Dae Choi,^a Pil Ja Seo,^a Byeng Wha Son^b and Uk Lee^{b*}

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea
Correspondence e-mail: uklee@pknu.ac.kr


Received 9 September 2009; accepted 13 September 2009

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.036; wR factor = 0.098; data-to-parameter ratio = 17.4.

In the title compound, $\text{C}_{14}\text{H}_{15}\text{FO}_4\text{S}$, the O atom and the methyl group of the methylsulfinyl substituent are located on opposite sides of the plane of the benzofuran fragment which is essentially planar with a mean deviation of $0.008(1)\text{ \AA}$ from its least-squares plane. The crystal structure stabilized by three different intermolecular non-classical C—H···O hydrogen bonds. The crystal structure also exhibits aromatic π – π interactions between the benzene rings of adjacent benzofuran ring systems [centroid–centroid distance = $3.688(2)\text{ \AA}$].

Related literature

For the crystal structures of similar alkyl 2-(5-fluoro-3-methylsulfinyl-1-benzofuran-2-yl) acetate derivatives, see: Choi *et al.* (2009*a,b*). For the pharmacological activity of benzofuran compounds, see: Howlett *et al.* (1999); Twyman & Allsop (1999).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{15}\text{FO}_4\text{S}$

$M_r = 298.32$

Monoclinic, $P2_1/c$
 $a = 11.6332(6)\text{ \AA}$
 $b = 14.9522(7)\text{ \AA}$
 $c = 8.2333(4)\text{ \AA}$
 $\beta = 102.277(1)^\circ$
 $V = 1399.36(12)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.25\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.25 \times 0.20 \times 0.16\text{ mm}$

Data collection

Bruker SMART CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2000)
 $T_{\min} = 0.940$, $T_{\max} = 0.961$

12229 measured reflections
3173 independent reflections
2476 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.098$
 $S = 1.09$
3173 reflections

182 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.39\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.39\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C5—H5···O3 ⁱ	0.93	2.50	3.370 (2)	155
C6—H6···O2 ⁱⁱ	0.93	2.54	3.369 (2)	149
C9—H9B···O4 ⁱⁱⁱ	0.97	2.26	3.228 (2)	176

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 2$; (iii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2158).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2001). *SAINT* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2009*a*). *Acta Cryst. E65*, o1826.
Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2009*b*). *Acta Cryst. E65*, o2096.
Farrugia, L. J. (1997). *J. Appl. Cryst. 30*, 565.
Howlett, D. R., Perry, A. E., Godfrey, F., Swatton, J. E., Jennings, K. H., Spitzfaden, C., Wadsworth, H., Wood, S. J. & Markwell, R. E. (1999). *Biochem. J. 340*, 283–289.
Sheldrick, G. M. (2000). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Twyman, L. J. & Allsop, D. (1999). *Tetrahedron Lett. 40*, 9383–9384.

supplementary materials

Acta Cryst. (2009). E65, o2488 [doi:10.1107/S1600536809037003]

Isopropyl 2-(5-fluoro-3-methylsulfinyl-1-benzofuran-2-yl)acetate

H. D. Choi, P. J. Seo, B. W. Son and U. Lee

Comment

Molecules involving benzofuran skeleton have attracted particular interest in view of their biological and pharmacological properties (Howlett *et al.*, 1999; Twyman & Allsop, 1999). As a part of our ongoing studies of the effect of side chain substituents on the solid state structures of alkyl 2-(5-fluoro-3-methylsulfinyl-1-benzofuran-2-yl) acetate analogues (Choi *et al.*, 2009*a,b*), we report the crystal structure of the title compound (Fig. 1).

The benzofuran unit is essentially planar, with a mean deviation of 0.008 (1) Å from the least-squares plane defined by the nine constituent atoms. The crystal packing (Fig. 2) is stabilized by three intermolecular non-classical C—H···O hydrogen bonds; the first between an H atom of the benzofuran ring and the oxygen of the C=O unit, with a C5—H5···O3ⁱ, the second between an H atom of the benzofuran ring and the oxygen of the isopropoxy group, with a C6—H6···O2ⁱⁱ, the third between a methylene H atom and the oxygen of the S=O unit, with a C9—H9B···O4ⁱⁱⁱ, respectively (Table 1). The crystal packing (Fig. 2) is further stabilized by aromatic π···π interactions between the benzene rings of neighboring molecules, with a Cg···Cgⁱ distance of 3.688 (2) Å (Cg is the centroid of the C2–C7 benzene ring).

Experimental

77% 3-chloroperoxybenzoic acid (247 mg, 1.1 mmol) was added in small portions to a stirred solution of isopropyl 2-(5-fluoro-3-methylsulfonyl-1-benzofuran-2-yl) acetate (282 mg, 1.0 mmol) in dichloromethane (30 ml) at 273 K. After being stirred for 4 h at room temperature, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated in vacuum. The residue was purified by column chromatography (hexane–ethyl acetate, 1:2 v/v) to afford the title compound as a colorless solid [yield 83%, m.p. 391–392 K; R_f = 0.67 (hexane–ethyl acetate, 1:2 v/v)]. Single crystals suitable for X-ray diffraction were prepared by evaporation of a solution of the title compound in acetone at room temperature. Spectroscopic analysis: EI-MS 298 [M^+].

Refinement

All H atoms were geometrically positioned and refined using a riding model, with C—H = 0.93 Å for the aryl, 0.97 Å for the methine, 0.98 Å for the methylene, and 0.96 Å for the methyl H atoms. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for the aryl, methine, and methylene H atoms, and $1.5U_{\text{eq}}(\text{C})$ for the methyl H atoms.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. The C—H···O and $\pi\cdots\pi$ interactions (dotted lines) in the title compound. Cg denotes the benzene ring centroid. [Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + 1, -y + 1, -z + 2$; (iii) $x, -y, z + 1/2$; (iv) $x, -y + 1/2, z - 1/2$.]

Isopropyl 2-(5-fluoro-3-methylsulfinyl-1-benzofuran-2-yl)acetate

Crystal data

C ₁₄ H ₁₅ FO ₄ S	$F_{000} = 624$
$M_r = 298.32$	$D_x = 1.416 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 5216 reflections
$a = 11.6332 (6) \text{ \AA}$	$\theta = 2.3\text{--}27.4^\circ$
$b = 14.9522 (7) \text{ \AA}$	$\mu = 0.25 \text{ mm}^{-1}$
$c = 8.2333 (4) \text{ \AA}$	$T = 173 \text{ K}$
$\beta = 102.2770 (10)^\circ$	Block, colorless
$V = 1399.36 (12) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.16 \text{ mm}$
Z = 4	

Data collection

Bruker SMART CCD diffractometer	3173 independent reflections
Radiation source: fine-focus sealed tube	2476 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.043$
Detector resolution: 10.0 pixels mm^{-1}	$\theta_{\text{max}} = 27.5^\circ$
$T = 173 \text{ K}$	$\theta_{\text{min}} = 1.8^\circ$
φ and ω scans	$h = -14\rightarrow 15$
Absorption correction: multi-scan (SADABS; Sheldrick, 2000)	$k = -19\rightarrow 19$
$T_{\text{min}} = 0.940, T_{\text{max}} = 0.961$	$l = -10\rightarrow 10$
12229 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.036$	H-atom parameters constrained
$wR(F^2) = 0.098$	$w = 1/[\sigma^2(F_o^2) + (0.0423P)^2 + 0.5417P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\max} = 0.001$
3173 reflections	$\Delta\rho_{\max} = 0.39 \text{ e \AA}^{-3}$
182 parameters	$\Delta\rho_{\min} = -0.39 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S	0.16889 (4)	0.32823 (3)	0.59899 (5)	0.02546 (13)
F	0.53377 (11)	0.32123 (8)	0.19202 (14)	0.0427 (3)
O1	0.47612 (10)	0.43326 (8)	0.79555 (14)	0.0242 (3)
O2	0.19830 (10)	0.54133 (8)	1.03615 (14)	0.0264 (3)
O3	0.17889 (11)	0.53806 (9)	0.75830 (15)	0.0329 (3)
O4	0.17181 (12)	0.23235 (9)	0.54858 (18)	0.0393 (3)
C1	0.31410 (14)	0.36896 (11)	0.6411 (2)	0.0222 (3)
C2	0.40529 (14)	0.36553 (11)	0.5455 (2)	0.0222 (3)
C3	0.41420 (16)	0.33514 (12)	0.3875 (2)	0.0264 (4)
H3	0.3516	0.3078	0.3155	0.032*
C4	0.52168 (16)	0.34834 (12)	0.3458 (2)	0.0289 (4)
C5	0.61801 (16)	0.38849 (12)	0.4473 (2)	0.0284 (4)
H5	0.6880	0.3950	0.4111	0.034*
C6	0.60937 (14)	0.41885 (12)	0.6032 (2)	0.0257 (4)
H6	0.6722	0.4464	0.6744	0.031*
C7	0.50247 (14)	0.40604 (11)	0.6470 (2)	0.0224 (3)
C8	0.36106 (14)	0.40933 (11)	0.7872 (2)	0.0229 (4)
C9	0.31082 (15)	0.43478 (12)	0.9330 (2)	0.0252 (4)
H9A	0.3743	0.4524	1.0239	0.030*
H9B	0.2725	0.3831	0.9689	0.030*
C10	0.22306 (14)	0.51050 (12)	0.8949 (2)	0.0234 (4)
C11	0.11677 (16)	0.61749 (12)	1.0234 (2)	0.0298 (4)

supplementary materials

H11	0.0555	0.6121	0.9218	0.036*
C12	0.06204 (17)	0.61224 (13)	1.1728 (2)	0.0340 (4)
H12A	0.0191	0.5572	1.1699	0.051*
H12B	0.0094	0.6618	1.1718	0.051*
H12C	0.1226	0.6143	1.2721	0.051*
C13	0.1857 (2)	0.70278 (14)	1.0176 (3)	0.0504 (6)
H13A	0.2183	0.7028	0.9199	0.076*
H13B	0.2482	0.7064	1.1146	0.076*
H13C	0.1344	0.7533	1.0147	0.076*
C14	0.11297 (18)	0.39058 (15)	0.4135 (2)	0.0397 (5)
H14A	0.1529	0.3725	0.3282	0.060*
H14B	0.1258	0.4532	0.4356	0.060*
H14C	0.0302	0.3794	0.3773	0.060*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S	0.0205 (2)	0.0281 (2)	0.0274 (2)	-0.00173 (17)	0.00402 (16)	-0.00285 (18)
F	0.0481 (7)	0.0526 (8)	0.0320 (6)	-0.0002 (6)	0.0189 (5)	-0.0125 (5)
O1	0.0211 (6)	0.0286 (6)	0.0221 (6)	-0.0011 (5)	0.0031 (5)	-0.0028 (5)
O2	0.0283 (6)	0.0295 (7)	0.0213 (6)	0.0066 (5)	0.0046 (5)	-0.0010 (5)
O3	0.0323 (7)	0.0439 (8)	0.0226 (6)	0.0096 (6)	0.0062 (5)	0.0042 (6)
O4	0.0339 (7)	0.0291 (7)	0.0529 (9)	-0.0065 (6)	0.0048 (6)	-0.0116 (6)
C1	0.0203 (8)	0.0202 (8)	0.0253 (8)	0.0012 (6)	0.0030 (6)	0.0000 (7)
C2	0.0222 (8)	0.0196 (8)	0.0241 (8)	0.0022 (6)	0.0033 (6)	0.0002 (7)
C3	0.0281 (9)	0.0243 (9)	0.0258 (8)	0.0002 (7)	0.0037 (7)	-0.0046 (7)
C4	0.0359 (10)	0.0275 (9)	0.0254 (9)	0.0052 (8)	0.0113 (8)	-0.0022 (7)
C5	0.0256 (9)	0.0279 (9)	0.0339 (9)	0.0044 (7)	0.0108 (7)	0.0028 (8)
C6	0.0200 (8)	0.0263 (9)	0.0298 (9)	0.0015 (7)	0.0029 (7)	0.0005 (7)
C7	0.0241 (8)	0.0213 (8)	0.0217 (8)	0.0028 (7)	0.0042 (6)	0.0000 (6)
C8	0.0198 (8)	0.0233 (8)	0.0251 (8)	0.0009 (6)	0.0037 (6)	0.0026 (7)
C9	0.0263 (9)	0.0283 (9)	0.0205 (8)	0.0009 (7)	0.0037 (7)	0.0005 (7)
C10	0.0215 (8)	0.0274 (9)	0.0214 (8)	-0.0046 (7)	0.0052 (6)	-0.0015 (7)
C11	0.0299 (9)	0.0320 (10)	0.0269 (9)	0.0098 (8)	0.0049 (7)	0.0021 (8)
C12	0.0301 (9)	0.0382 (11)	0.0359 (10)	0.0079 (8)	0.0119 (8)	0.0019 (9)
C13	0.0648 (15)	0.0322 (11)	0.0636 (15)	0.0043 (10)	0.0345 (13)	0.0087 (11)
C14	0.0302 (10)	0.0517 (13)	0.0329 (10)	0.0040 (9)	-0.0030 (8)	0.0067 (9)

Geometric parameters (\AA , $^\circ$)

S—O4	1.4949 (14)	C6—C7	1.380 (2)
S—C1	1.7595 (17)	C6—H6	0.9300
S—C14	1.7894 (19)	C8—C9	1.491 (2)
F—C4	1.365 (2)	C9—C10	1.512 (2)
O1—C8	1.3731 (19)	C9—H9A	0.9700
O1—C7	1.384 (2)	C9—H9B	0.9700
O2—C10	1.338 (2)	C11—C12	1.502 (2)
O2—C11	1.472 (2)	C11—C13	1.513 (3)
O3—C10	1.205 (2)	C11—H11	0.9800

C1—C8	1.353 (2)	C12—H12A	0.9600
C1—C2	1.450 (2)	C12—H12B	0.9600
C2—C7	1.394 (2)	C12—H12C	0.9600
C2—C3	1.402 (2)	C13—H13A	0.9600
C3—C4	1.380 (2)	C13—H13B	0.9600
C3—H3	0.9300	C13—H13C	0.9600
C4—C5	1.384 (3)	C14—H14A	0.9600
C5—C6	1.385 (3)	C14—H14B	0.9600
C5—H5	0.9300	C14—H14C	0.9600
O4—S—C1	108.00 (8)	C10—C9—H9A	109.0
O4—S—C14	106.75 (10)	C8—C9—H9B	109.0
C1—S—C14	98.47 (9)	C10—C9—H9B	109.0
C8—O1—C7	106.03 (12)	H9A—C9—H9B	107.8
C10—O2—C11	117.35 (13)	O3—C10—O2	124.43 (16)
C8—C1—C2	107.15 (14)	O3—C10—C9	125.76 (15)
C8—C1—S	121.23 (13)	O2—C10—C9	109.79 (14)
C2—C1—S	131.57 (13)	O2—C11—C12	106.27 (14)
C7—C2—C3	119.14 (15)	O2—C11—C13	108.45 (16)
C7—C2—C1	104.64 (14)	C12—C11—C13	112.95 (17)
C3—C2—C1	136.21 (16)	O2—C11—H11	109.7
C4—C3—C2	115.75 (16)	C12—C11—H11	109.7
C4—C3—H3	122.1	C13—C11—H11	109.7
C2—C3—H3	122.1	C11—C12—H12A	109.5
F—C4—C3	117.80 (16)	C11—C12—H12B	109.5
F—C4—C5	117.32 (16)	H12A—C12—H12B	109.5
C3—C4—C5	124.87 (16)	C11—C12—H12C	109.5
C4—C5—C6	119.52 (16)	H12A—C12—H12C	109.5
C4—C5—H5	120.2	H12B—C12—H12C	109.5
C6—C5—H5	120.2	C11—C13—H13A	109.5
C7—C6—C5	116.35 (16)	C11—C13—H13B	109.5
C7—C6—H6	121.8	H13A—C13—H13B	109.5
C5—C6—H6	121.8	C11—C13—H13C	109.5
C6—C7—O1	124.87 (15)	H13A—C13—H13C	109.5
C6—C7—C2	124.37 (16)	H13B—C13—H13C	109.5
O1—C7—C2	110.75 (14)	S—C14—H14A	109.5
C1—C8—O1	111.43 (14)	S—C14—H14B	109.5
C1—C8—C9	132.54 (16)	H14A—C14—H14B	109.5
O1—C8—C9	115.98 (14)	S—C14—H14C	109.5
C8—C9—C10	112.79 (14)	H14A—C14—H14C	109.5
C8—C9—H9A	109.0	H14B—C14—H14C	109.5
O4—S—C1—C8	-126.67 (15)	C3—C2—C7—C6	0.0 (3)
C14—S—C1—C8	122.54 (15)	C1—C2—C7—C6	-178.66 (16)
O4—S—C1—C2	50.55 (18)	C3—C2—C7—O1	178.64 (14)
C14—S—C1—C2	-60.23 (18)	C1—C2—C7—O1	0.01 (18)
C8—C1—C2—C7	-0.17 (18)	C2—C1—C8—O1	0.27 (19)
S—C1—C2—C7	-177.69 (14)	S—C1—C8—O1	178.10 (11)
C8—C1—C2—C3	-178.44 (19)	C2—C1—C8—C9	177.57 (17)
S—C1—C2—C3	4.0 (3)	S—C1—C8—C9	-4.6 (3)

supplementary materials


C7—C2—C3—C4	0.1 (2)	C7—O1—C8—C1	-0.26 (18)
C1—C2—C3—C4	178.20 (18)	C7—O1—C8—C9	-178.05 (14)
C2—C3—C4—F	-178.87 (15)	C1—C8—C9—C10	-69.4 (2)
C2—C3—C4—C5	0.0 (3)	O1—C8—C9—C10	107.82 (16)
F—C4—C5—C6	178.67 (16)	C11—O2—C10—O3	-3.7 (2)
C3—C4—C5—C6	-0.2 (3)	C11—O2—C10—C9	178.11 (14)
C4—C5—C6—C7	0.3 (3)	C8—C9—C10—O3	13.5 (2)
C5—C6—C7—O1	-178.66 (15)	C8—C9—C10—O2	-168.35 (14)
C5—C6—C7—C2	-0.2 (3)	C10—O2—C11—C12	154.84 (15)
C8—O1—C7—C6	178.81 (16)	C10—O2—C11—C13	-83.45 (19)
C8—O1—C7—C2	0.15 (18)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
C5—H5 ⁱ …O3 ⁱ	0.93	2.50	3.370 (2)	155
C6—H6 ⁱⁱ …O2 ⁱⁱ	0.93	2.54	3.369 (2)	149
C9—H9B ⁱⁱⁱ …O4 ⁱⁱⁱ	0.97	2.26	3.228 (2)	176

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+1, -y+1, -z+2$; (iii) $x, -y+1/2, z+1/2$.

Fig. 1



supplementary materials

Fig. 2

