

1-(1,3-Benzothiazol-2-yl)-3-(4-chlorobenzoyl)thiourea

M. Sukeri M. Yusof,^{a*} Zakaria S. Aishah,^a Wan M. Khairul^a and Bohari M. Yamin^b

^aDepartment of Chemical Sciences, Faculty of Science and Technology, Universiti Malaysia Terengganu, Mengabang Telipot, 21030 Kuala Terengganu, Malaysia, and

^bSchool of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

Correspondence e-mail: mohdsukeri@umt.edu.my


Received 25 August 2009; accepted 16 September 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.039; wR factor = 0.109; data-to-parameter ratio = 19.0.

The title compound, $\text{C}_{15}\text{H}_{10}\text{ClN}_3\text{OS}_2$, adopts a *cis-trans* configuration across the thiourea C–N bonds with respect to the positions of the benzothiazole and 4-chlorobenzoyl groups relative to thiono S atom. An intramolecular N–H···O hydrogen bond is present. In the crystal structure, molecules are linked by a weak intermolecular N–H···S hydrogen bond, forming centrosymmetric dimers.

Related literature

For the biological activity of thiadiazoles, see: Shukla & Srivastava (2008); Göblyös *et al.* (2005); Terzioglu & Gürsoy (2003); Rana *et al.* (2008). For their potential as insecticides and fungicides, see: Jian *et al.* (2005). For C–S and C–O bond lengths, see: Saeed & Flörke (2006); Yamin & Yusof (2003). For the structures of other benzoylthiourea derivatives, see: Dillen *et al.* (2006); Khawar Rauf *et al.* (2006); Weiqun *et al.* (2004).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{10}\text{ClN}_3\text{OS}_2$

$M_r = 347.83$

Monoclinic, $P2_1/c$

$a = 11.726(2)\text{ \AA}$

$b = 17.934(4)\text{ \AA}$

$c = 7.2617(16)\text{ \AA}$

$\beta = 96.848(4)^\circ$

$V = 1516.1(6)\text{ \AA}^3$

$Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.53\text{ mm}^{-1}$

$T = 298\text{ K}$
 $0.55 \times 0.42 \times 0.40\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.759$, $T_{\max} = 0.816$

11030 measured reflections
3772 independent reflections
2891 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.109$
 $S = 1.05$
3772 reflections

199 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.30\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.22\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N2–H2A···O1	0.86	1.88	2.6056 (19)	141
N1–H1A···S1 ⁱ	0.86	2.75	3.5377 (17)	152

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2009).

The authors would like to thank the Malaysian Government, Universiti Kebangsaan Malaysia, Universiti Malaysia Terengganu and the Ministry of Science, Technology and Innovation for the research grants e-science fund 52022.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2003).

References

- Bruker (2000). *SADABS, SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dillen, J., Woldu, M. G. & Koch, K. R. (2006). *Acta Cryst. E62*, o5225–o5227.
- Göblyös, A., Vries, H., Brussee, J. & Ijzerman, A. P. (2005). *J. Med. Chem.* **48**, 1145–1151.
- Jian, F., Zhao, P., Hou, Y. & Lu, L. (2005). *Struct. Chem.* **16**, 123–128.
- Khawar Rauf, M., Badshah, A. & Flörke, U. (2006). *Acta Cryst. E62*, o2452–o2453.
- Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
- Rana, A., Siddiqui, N., Khan, S. A., Haque, S. E. & Bhat, M. A. (2008). *Eur. J. Med. Chem.* **43**, 1114–1122.
- Saeed, A. & Flörke, U. (2006). *Acta Cryst. E62*, o2924–o2925.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Shukla, D. K. & Srivastava, S. D. (2008). *Indian J. Chem. Sect. B*, **47**, 463–469.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
- Terzioglu, N. & Gürsoy, A. (2003). *Eur. J. Med. Chem.* **38**, 781–786.
- Weiqun, Z., Baolong, L., Liming, Z., Jiangang, D., Yong, Z., Lude, L. & Xujie, Y. (2004). *J. Mol. Struct.* **690**, 145–150.
- Yamin, B. M. & Yusof, M. S. M. (2003). *Acta Cryst. E59*, o151–o152.

supplementary materials

Acta Cryst. (2009). E65, o2519 [doi:10.1107/S160053680903743X]

1-(1,3-Benzothiazol-2-yl)-3-(4-chlorobenzoyl)thiourea

M. S. M. Yusof, Z. S. Aishah, W. M. Khairul and B. M. Yamin

Comment

For the past few decades, heterocycles featuring thiadiazoles which consist of sulfur and nitrogen have been developed consistently to act as drugs as well as to play an active role in numerous biological activities (Shukla & Srivastava 2008; Göblyös *et al.* 2005). These derivatives have been shown to exhibit anticancer, antitubercular and anticonvulsant activities (Terzioglu & Gürsoy 2003; Rana *et al.* 2008). In agriculture, thiadiazole derivatives have a potential as insecticide and fungicide (Jian *et al.* 2005). The title compound (I), adopts *cis-trans* configuration with respect to the positions of the benzothiazole and 4-chlorobenzoyl groups relative to the thiono S atom, across their C—N bonds (Fig 1). The central carbonyl thiourea moiety (S1/C8/N1/N2/C7/O1), phenyl ring (C1—C6) and benzothiazole (S2/N3/C9—C15) groups are all planar, with a maximum deviation of 0.021 (1) Å for atom C10 from the least-squares plane. The central carbonyl thiourea fragment makes dihedral angles of 24.09 (7)° and 4.58 (4)° with the phenyl ring and benzothiazole group, respectively. The two aryl rings are inclined to each other at an angle of 28.42 (8)°. The C8—S1 and C7—O1 bond length show the expected double bond character of 1.6570 (17) Å and 1.2245 (19) Å (Saeed & Flörke 2006; Yamin & Yusof 2003). The N1—C8 is longer than N2—C8 by 0.045 Å, similar to other benzoylthiourea derivatives (Dillen *et al.* 2006; Khawar Rauf *et al.* 2006) which is probably due to the intramolecular hydrogen bonding interaction (Weioun *et al.* 2004).

There is an intramolecular hydrogen bond, N2—H2···O1 forming a pseudo-six-membered ring, O1···H2—N2—C8—N1—C7—O1 (Fig. 1). In the crystal structure, the molecules are linked by a weak intermolecular interaction N1—H1A···S1 (symmetry codes as in Table 1) forming dimers (Fig. 2).

Experimental

To a stirring acetone solution (75 ml) of 4-chlorobenzoyl chloride (2.0 g, 11.4 mmol) and ammoniumthiocyanate (0.87 g, 11.4 mmol), 2-aminobenzothiazole (1.17 g, 11.4 mmol) in 40 ml of acetone was added dropwise. The solution mixture was put at reflux for 1 h. The resulting solution was poured into a beaker containing some ice blocks. The light yellow precipitate was filtered off and washed with distilled water and cold ethanol before dried under vacuum. Good quality crystals were obtained by recrystallization from DMSO.

Refinement

After their location in the difference map, all H-atoms were fixed geometrically at ideal positions and allowed to ride on the parent C or N atoms with C—H = 0.93 Å and N—H = 0.86 Å with $U_{\text{iso}}(\text{H})= 1.2$ (CH and NH).

Figures




Fig. 1. The molecule of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. Dashed lines indicate hydrogen bonds.

supplementary materials



Fig. 2. Packing diagram of compound (I), viewed down the *b* axis. The dashed lines denote the N—H···S hydrogen bond.

1-(1,3-Benzothiazol-2-yl)-3-(4-chlorobenzoyl)thiourea

Crystal data

C ₁₅ H ₁₀ ClN ₃ OS ₂	$F_{000} = 712$
$M_r = 347.83$	$D_x = 1.524 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 906 reflections
$a = 11.726 (2) \text{ \AA}$	$\theta = 1.8\text{--}28.3^\circ$
$b = 17.934 (4) \text{ \AA}$	$\mu = 0.53 \text{ mm}^{-1}$
$c = 7.2617 (16) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 96.848 (4)^\circ$	Block, light yellow
$V = 1516.1 (6) \text{ \AA}^3$	$0.55 \times 0.42 \times 0.40 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	3772 independent reflections
Radiation source: fine-focus sealed tube	2891 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.022$
Detector resolution: 83.66 pixels mm^{-1}	$\theta_{\text{max}} = 28.3^\circ$
$T = 298 \text{ K}$	$\theta_{\text{min}} = 1.8^\circ$
ω scans	$h = -15 \rightarrow 15$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$k = -23 \rightarrow 20$
$T_{\text{min}} = 0.759$, $T_{\text{max}} = 0.816$	$l = -8 \rightarrow 9$
11030 measured reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H-atom parameters constrained
$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0606P)^2 + 0.1737P]$
$S = 1.05$	where $P = (F_o^2 + 2F_c^2)/3$
3772 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
199 parameters	$\Delta\rho_{\text{max}} = 0.30 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.22 \text{ e \AA}^{-3}$

Primary atom site location: structure-invariant direct methods Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.84415 (5)	0.83795 (3)	0.39677 (9)	0.07175 (19)
S1	0.49665 (4)	0.38829 (3)	0.39437 (8)	0.05844 (16)
S2	0.58802 (4)	0.22891 (2)	0.41252 (6)	0.04563 (13)
O1	0.86117 (10)	0.46684 (7)	0.3645 (2)	0.0549 (3)
N3	0.80322 (12)	0.24282 (8)	0.3637 (2)	0.0474 (3)
N1	0.67029 (11)	0.48112 (8)	0.3894 (2)	0.0450 (3)
H1A	0.6183	0.5141	0.4005	0.054*
N2	0.71709 (11)	0.35696 (7)	0.3786 (2)	0.0434 (3)
H2A	0.7842	0.3750	0.3695	0.052*
C1	0.70106 (14)	0.63867 (10)	0.3181 (3)	0.0491 (4)
H1	0.6291	0.6194	0.2754	0.059*
C2	0.71738 (15)	0.71472 (10)	0.3236 (3)	0.0520 (4)
H2	0.6570	0.7469	0.2855	0.062*
C3	0.82429 (16)	0.74219 (10)	0.3862 (3)	0.0489 (4)
C4	0.91544 (15)	0.69616 (11)	0.4420 (3)	0.0551 (5)
H4	0.9873	0.7159	0.4832	0.066*
C5	0.89835 (14)	0.61965 (10)	0.4358 (3)	0.0507 (4)
H5	0.9594	0.5877	0.4722	0.061*
C6	0.79104 (13)	0.59043 (9)	0.3759 (2)	0.0418 (4)
C7	0.77928 (13)	0.50832 (10)	0.3753 (2)	0.0420 (4)
C8	0.63429 (13)	0.40769 (9)	0.3881 (2)	0.0417 (4)
C9	0.71104 (14)	0.27973 (9)	0.3813 (2)	0.0404 (4)
C10	0.78156 (15)	0.16707 (10)	0.3728 (2)	0.0459 (4)
C11	0.86167 (18)	0.11049 (11)	0.3579 (3)	0.0612 (5)
H11	0.9372	0.1220	0.3421	0.073*
C12	0.82695 (19)	0.03736 (12)	0.3672 (3)	0.0685 (6)
H12	0.8797	-0.0007	0.3568	0.082*
C13	0.7148 (2)	0.01951 (11)	0.3915 (3)	0.0657 (6)
H13	0.6932	-0.0303	0.3964	0.079*
C14	0.63489 (17)	0.07448 (10)	0.4087 (3)	0.0574 (5)
H14	0.5599	0.0624	0.4266	0.069*

supplementary materials

C15 0.66891 (15) 0.14852 (9) 0.3988 (2) 0.0447 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0907 (4)	0.0384 (3)	0.0882 (4)	-0.0049 (2)	0.0194 (3)	-0.0016 (2)
S1	0.0365 (2)	0.0450 (3)	0.0947 (4)	-0.00093 (18)	0.0117 (2)	-0.0050 (2)
S2	0.0387 (2)	0.0442 (2)	0.0534 (3)	-0.00318 (16)	0.00325 (18)	0.00143 (18)
O1	0.0381 (6)	0.0427 (7)	0.0842 (10)	0.0047 (5)	0.0088 (6)	-0.0007 (6)
N3	0.0428 (7)	0.0402 (8)	0.0602 (10)	-0.0005 (6)	0.0099 (7)	-0.0024 (6)
N1	0.0356 (7)	0.0381 (7)	0.0618 (9)	0.0029 (5)	0.0076 (6)	-0.0010 (6)
N2	0.0352 (6)	0.0381 (7)	0.0567 (9)	-0.0015 (5)	0.0045 (6)	-0.0020 (6)
C1	0.0366 (8)	0.0454 (9)	0.0649 (12)	0.0015 (7)	0.0046 (8)	0.0055 (8)
C2	0.0449 (9)	0.0453 (10)	0.0664 (12)	0.0083 (7)	0.0085 (8)	0.0083 (8)
C3	0.0570 (10)	0.0360 (8)	0.0557 (11)	-0.0018 (7)	0.0150 (8)	0.0002 (7)
C4	0.0428 (9)	0.0491 (10)	0.0727 (13)	-0.0069 (8)	0.0042 (8)	-0.0027 (9)
C5	0.0361 (8)	0.0456 (10)	0.0696 (12)	0.0023 (7)	0.0037 (8)	0.0028 (8)
C6	0.0355 (7)	0.0401 (9)	0.0503 (10)	0.0012 (6)	0.0071 (7)	0.0008 (7)
C7	0.0353 (7)	0.0438 (9)	0.0469 (9)	0.0007 (6)	0.0043 (7)	0.0009 (7)
C8	0.0391 (8)	0.0404 (8)	0.0455 (9)	0.0001 (6)	0.0043 (7)	-0.0014 (7)
C9	0.0387 (8)	0.0399 (8)	0.0419 (9)	-0.0021 (6)	0.0018 (6)	-0.0021 (6)
C10	0.0486 (9)	0.0411 (9)	0.0476 (10)	0.0009 (7)	0.0034 (7)	-0.0031 (7)
C11	0.0588 (11)	0.0462 (10)	0.0798 (14)	0.0061 (8)	0.0135 (10)	-0.0040 (9)
C12	0.0754 (14)	0.0442 (11)	0.0844 (16)	0.0115 (10)	0.0034 (12)	-0.0080 (10)
C13	0.0804 (15)	0.0391 (10)	0.0735 (14)	-0.0046 (10)	-0.0076 (11)	0.0004 (9)
C14	0.0605 (11)	0.0462 (10)	0.0628 (12)	-0.0098 (9)	-0.0035 (9)	0.0046 (8)
C15	0.0482 (9)	0.0414 (9)	0.0427 (9)	-0.0020 (7)	-0.0018 (7)	0.0006 (7)

Geometric parameters (\AA , $^\circ$)

Cl1—C3	1.7334 (19)	C2—H2	0.9300
S1—C8	1.6570 (17)	C3—C4	1.373 (3)
S2—C15	1.7352 (18)	C4—C5	1.386 (3)
S2—C9	1.7438 (17)	C4—H4	0.9300
O1—C7	1.2245 (19)	C5—C6	1.384 (2)
N3—C9	1.287 (2)	C5—H5	0.9300
N3—C10	1.385 (2)	C6—C7	1.479 (2)
N1—C8	1.383 (2)	C10—C15	1.397 (2)
N1—C7	1.383 (2)	C10—C11	1.396 (3)
N1—H1A	0.8600	C11—C12	1.377 (3)
N2—C8	1.338 (2)	C11—H11	0.9300
N2—C9	1.387 (2)	C12—C13	1.385 (3)
N2—H2A	0.8600	C12—H12	0.9300
C1—C2	1.377 (2)	C13—C14	1.376 (3)
C1—C6	1.390 (2)	C13—H13	0.9300
C1—H1	0.9300	C14—C15	1.391 (2)
C2—C3	1.373 (3)	C14—H14	0.9300
C15—S2—C9	87.73 (8)	O1—C7—C6	122.10 (15)

C9—N3—C10	109.80 (14)	N1—C7—C6	115.95 (14)
C8—N1—C7	128.28 (14)	N2—C8—N1	115.16 (14)
C8—N1—H1A	115.9	N2—C8—S1	125.01 (13)
C7—N1—H1A	115.9	N1—C8—S1	119.83 (12)
C8—N2—C9	129.75 (14)	N3—C9—N2	117.86 (15)
C8—N2—H2A	115.1	N3—C9—S2	117.51 (13)
C9—N2—H2A	115.1	N2—C9—S2	124.62 (12)
C2—C1—C6	120.65 (16)	N3—C10—C15	114.96 (15)
C2—C1—H1	119.7	N3—C10—C11	125.47 (17)
C6—C1—H1	119.7	C15—C10—C11	119.57 (17)
C3—C2—C1	118.89 (16)	C12—C11—C10	118.9 (2)
C3—C2—H2	120.6	C12—C11—H11	120.6
C1—C2—H2	120.6	C10—C11—H11	120.6
C4—C3—C2	122.00 (17)	C11—C12—C13	121.1 (2)
C4—C3—Cl1	119.14 (15)	C11—C12—H12	119.4
C2—C3—Cl1	118.86 (14)	C13—C12—H12	119.4
C3—C4—C5	118.75 (17)	C14—C13—C12	120.87 (19)
C3—C4—H4	120.6	C14—C13—H13	119.6
C5—C4—H4	120.6	C12—C13—H13	119.6
C6—C5—C4	120.48 (16)	C13—C14—C15	118.47 (19)
C6—C5—H5	119.8	C13—C14—H14	120.8
C4—C5—H5	119.8	C15—C14—H14	120.8
C5—C6—C1	119.21 (16)	C14—C15—C10	121.07 (17)
C5—C6—C7	117.30 (14)	C14—C15—S2	128.91 (15)
C1—C6—C7	123.48 (15)	C10—C15—S2	110.01 (13)
O1—C7—N1	121.94 (16)		
C6—C1—C2—C3	0.3 (3)	C10—N3—C9—S2	0.4 (2)
C1—C2—C3—C4	0.5 (3)	C8—N2—C9—N3	177.61 (17)
C1—C2—C3—Cl1	−178.90 (15)	C8—N2—C9—S2	−4.0 (3)
C2—C3—C4—C5	−0.4 (3)	C15—S2—C9—N3	−0.44 (15)
Cl1—C3—C4—C5	179.00 (15)	C15—S2—C9—N2	−178.82 (15)
C3—C4—C5—C6	−0.5 (3)	C9—N3—C10—C15	−0.2 (2)
C4—C5—C6—C1	1.4 (3)	C9—N3—C10—C11	179.64 (19)
C4—C5—C6—C7	−179.30 (17)	N3—C10—C11—C12	−179.08 (18)
C2—C1—C6—C5	−1.3 (3)	C15—C10—C11—C12	0.8 (3)
C2—C1—C6—C7	179.44 (17)	C10—C11—C12—C13	−0.3 (3)
C8—N1—C7—O1	−2.2 (3)	C11—C12—C13—C14	−0.5 (3)
C8—N1—C7—C6	178.38 (16)	C12—C13—C14—C15	0.8 (3)
C5—C6—C7—O1	−24.7 (3)	C13—C14—C15—C10	−0.3 (3)
C1—C6—C7—O1	154.64 (18)	C13—C14—C15—S2	179.10 (15)
C5—C6—C7—N1	154.75 (16)	N3—C10—C15—C14	179.37 (16)
C1—C6—C7—N1	−25.9 (2)	C11—C10—C15—C14	−0.5 (3)
C9—N2—C8—N1	177.45 (16)	N3—C10—C15—S2	−0.1 (2)
C9—N2—C8—S1	−3.6 (3)	C11—C10—C15—S2	−179.97 (15)
C7—N1—C8—N2	2.7 (3)	C9—S2—C15—C14	−179.14 (18)
C7—N1—C8—S1	−176.34 (14)	C9—S2—C15—C10	0.28 (13)
C10—N3—C9—N2	178.94 (14)		


supplementary materials

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N2—H2A···O1	0.86	1.88	2.6056 (19)	141
N1—H1A···S1 ⁱ	0.86	2.75	3.5377 (17)	152
C5—H5···O1 ⁱⁱ	0.93	2.49	3.389 (2)	163

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+2, -y+1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

