

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-Phenyl-N-(3-phenylprop-2-ynyl)aniline

Tao Pang, Yi-chong Sun and Jian-ming Zhang*

Key Laboratory of Pesticides and Chemical Biology of the Ministry of Education, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: leileitaotao6666@163.com


Received 29 April 2009; accepted 18 September 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.055; wR factor = 0.141; data-to-parameter ratio = 9.8.

In the title compound, $\text{C}_{21}\text{H}_{17}\text{N}$, synthesized by a three-component coupling reaction in the presence of copper(I) iodide, the N-bound phenyl rings form a dihedral angle of 72.5 (1)° with each other. There are no remarkable interactions in the crystal structure.

Related literature

For the preparation of the title compound, see: Nilsson *et al.* (1992). For the biological activity of propargylamines and their use as synthetic intermediates, see: Bieber & da Silva (2004); Hattori *et al.* (1993); Huffman *et al.* (1995); Konishi *et al.* (1990).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{17}\text{N}$
 $M_r = 283.36$
 Monoclinic, $P2_1$
 $a = 11.376$ (1) Å
 $b = 5.7287$ (5) Å
 $c = 13.409$ (1) Å
 $\beta = 111.276$ (3)°

$V = 814.30$ (12) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 298$ K
 $0.23 \times 0.13 \times 0.10$ mm

Data collection

Bruker SMART CCD
 diffractometer
 Absorption correction: none
 5689 measured reflections

1953 independent reflections
 1448 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.136$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.141$
 $S = 0.91$
 1953 reflections
 199 parameters

1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.15$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.19$ e Å⁻³

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The author is grateful to Central China Normal University for support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LX2105).

References

- Bieber, L. W. & da Silva, M. F. (2004). *Tetrahedron Lett.* **45**, 8281–8283.
 Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Hattori, K., Miyata, M. & Yamamoto, H. (1993). *J. Am. Chem. Soc.* **115**, 1151–1152.
 Huffman, M. A., Yasuda, N., DeCamp, A. E. & Grabowski, E. J. J. (1995). *J. Org. Chem.* **60**, 1590–1594.
 Konishi, M., Ohkuma, H., Tsuno, T., Oki, T., VanDuyne, G. D. & Clardy, J. (1990). *J. Am. Chem. Soc.* **112**, 3715–3716.
 Nilsson, B. M., Vargas, H. M., Ringdahl, B. & Hacksell, U. (1992). *J. Med. Chem.* **35**, 285–294.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o2555 [doi:10.1107/S1600536809037866]

N-Phenyl-*N*-(3-phenylprop-2-ynyl)aniline

T. Pang, Y. Sun and J. Zhang

Comment

Propargylamines are compounds of interesting biological properties and important synthetic intermediates (Konishi *et al.*, 1990; Huffman *et al.*, 1995; Hattori *et al.*, 1993; Bieber *et al.*, 2004). The reaction which a three component procedure between terminal alkynes, formaldehyde and secondary amines has been extended to some less activated alkynes by the introduction of copper catalysts. Here we report the crystal structure of the title compound (Fig. 1).

In the molecule of the title compound, the *N*-bound two phenyl rings form a dihedral angle of 72.5 (1)° with each other.

Experimental

The title compound was synthesized according to the literature procedure of Nilsson *et al.* (1992). Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in chloroform : methanol (50 : 1) at room temperature.

Refinement

All H atoms were initially located in a difference map, but were constrained to an idealized geometry. Constrained bond lengths and isotropic displacement parameters: (C–H = 0.97 Å) and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ for methylene, and (C–H = 0.93 Å) and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ for aromatic H atoms.

Figures




Fig. 1. View of the molecule of (I) showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented by spheres of arbitrary radius.

N-Phenyl-*N*-(3-phenylprop-2-ynyl)aniline

Crystal data

C₂₁H₁₇N

$M_r = 283.36$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$F_{000} = 300$

$D_x = 1.156 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1607 reflections

supplementary materials

$a = 11.376 (1) \text{ \AA}$	$\theta = 2.9\text{--}22.6^\circ$
$b = 5.7287 (5) \text{ \AA}$	$\mu = 0.07 \text{ mm}^{-1}$
$c = 13.409 (1) \text{ \AA}$	$T = 298 \text{ K}$
$\beta = 111.276 (3)^\circ$	Block, colorless
$V = 814.30 (12) \text{ \AA}^3$	$0.23 \times 0.13 \times 0.10 \text{ mm}$
$Z = 2$	

Data collection

Bruker SMART CCD diffractometer	1953 independent reflections
Radiation source: fine-focus sealed tube	1448 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.136$
Detector resolution: $10.0 \text{ pixels mm}^{-1}$	$\theta_{\text{max}} = 27.0^\circ$
$T = 298 \text{ K}$	$\theta_{\text{min}} = 1.6^\circ$
φ and ω scans	$h = -13 \rightarrow 14$
Absorption correction: none	$k = -7 \rightarrow 7$
5689 measured reflections	$l = -17 \rightarrow 10$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.055$	H-atom parameters constrained
$wR(F^2) = 0.141$	$w = 1/[\sigma^2(F_o^2) + (0.074P)^2]$
$S = 0.91$	where $P = (F_o^2 + 2F_c^2)/3$
1953 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
199 parameters	$\Delta\rho_{\text{max}} = 0.15 \text{ e \AA}^{-3}$
1 restraint	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2341 (2)	0.1972 (5)	0.3674 (2)	0.0541 (7)
C2	0.1912 (3)	0.1537 (7)	0.4505 (2)	0.0716 (9)
H2	0.1255	0.2420	0.4565	0.086*
C3	0.2457 (4)	-0.0199 (7)	0.5240 (3)	0.0859 (11)
H3	0.2158	-0.0466	0.5789	0.103*
C4	0.3427 (3)	-0.1539 (8)	0.5183 (3)	0.0870 (11)
H4	0.3790	-0.2695	0.5688	0.104*
C5	0.3851 (3)	-0.1131 (7)	0.4356 (3)	0.0775 (9)
H5	0.4500	-0.2039	0.4298	0.093*
C6	0.3327 (2)	0.0596 (6)	0.3619 (2)	0.0606 (7)
H6	0.3635	0.0853	0.3074	0.073*
C7	0.1821 (2)	0.3552 (4)	0.1865 (2)	0.0473 (6)
C8	0.1297 (2)	0.1628 (5)	0.1253 (2)	0.0555 (7)
H8	0.0980	0.0420	0.1546	0.067*
C9	0.1235 (3)	0.1472 (6)	0.0206 (2)	0.0646 (8)
H9	0.0885	0.0158	-0.0201	0.078*
C10	0.1692 (3)	0.3265 (6)	-0.0233 (2)	0.0651 (8)
H10	0.1657	0.3167	-0.0936	0.078*
C11	0.2198 (3)	0.5184 (6)	0.0370 (3)	0.0690 (8)
H11	0.2496	0.6407	0.0072	0.083*
C12	0.2273 (3)	0.5334 (5)	0.1416 (2)	0.0606 (7)
H12	0.2632	0.6645	0.1821	0.073*
C13	0.0902 (3)	0.5331 (5)	0.3057 (2)	0.0637 (7)
H13A	0.0843	0.6719	0.2627	0.076*
H13B	0.1166	0.5817	0.3799	0.076*
C14	-0.0356 (3)	0.4259 (6)	0.2741 (2)	0.0611 (7)
C15	-0.1346 (3)	0.3302 (6)	0.2484 (2)	0.0627 (7)
C16	-0.2554 (3)	0.2165 (6)	0.2187 (2)	0.0589 (7)
C17	-0.2702 (4)	0.0104 (7)	0.2656 (3)	0.0808 (9)
H17	-0.2010	-0.0604	0.3170	0.097*
C18	-0.3885 (5)	-0.0912 (9)	0.2361 (4)	0.1097 (16)
H18	-0.3990	-0.2296	0.2681	0.132*
C19	-0.4903 (5)	0.0132 (13)	0.1594 (5)	0.1160 (19)
H19	-0.5697	-0.0548	0.1397	0.139*
C20	-0.4753 (3)	0.2131 (11)	0.1130 (4)	0.1032 (15)
H20	-0.5446	0.2817	0.0607	0.124*
C21	-0.3604 (3)	0.3163 (7)	0.1413 (3)	0.0760 (9)
H21	-0.3518	0.4551	0.1086	0.091*
N1	0.1865 (2)	0.3795 (4)	0.29364 (18)	0.0548 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0509 (14)	0.0643 (16)	0.0386 (12)	-0.0118 (12)	0.0061 (10)	-0.0006 (13)

supplementary materials

C2	0.0736 (18)	0.092 (2)	0.0501 (16)	-0.0029 (18)	0.0231 (14)	0.0022 (17)
C3	0.094 (2)	0.114 (3)	0.0463 (16)	-0.006 (2)	0.0216 (17)	0.0205 (19)
C4	0.082 (2)	0.101 (3)	0.061 (2)	0.004 (2)	0.0065 (17)	0.026 (2)
C5	0.0654 (18)	0.086 (2)	0.0677 (19)	0.0086 (16)	0.0085 (15)	0.0125 (19)
C6	0.0503 (14)	0.0747 (19)	0.0496 (15)	-0.0017 (14)	0.0096 (12)	0.0065 (14)
C7	0.0430 (12)	0.0519 (14)	0.0453 (13)	-0.0001 (11)	0.0139 (10)	0.0052 (12)
C8	0.0564 (15)	0.0556 (16)	0.0488 (14)	-0.0085 (12)	0.0120 (11)	0.0017 (13)
C9	0.0681 (18)	0.0638 (18)	0.0508 (15)	-0.0011 (14)	0.0082 (13)	-0.0048 (14)
C10	0.0657 (17)	0.083 (2)	0.0476 (15)	0.0153 (16)	0.0216 (13)	0.0090 (16)
C11	0.0769 (19)	0.0682 (19)	0.071 (2)	-0.0052 (16)	0.0381 (16)	0.0152 (18)
C12	0.0650 (16)	0.0546 (15)	0.0655 (18)	-0.0137 (14)	0.0275 (13)	-0.0015 (15)
C13	0.0715 (18)	0.0593 (16)	0.0621 (17)	0.0007 (15)	0.0265 (14)	-0.0023 (15)
C14	0.0622 (17)	0.0729 (18)	0.0523 (15)	0.0082 (15)	0.0257 (13)	-0.0003 (15)
C15	0.0634 (17)	0.080 (2)	0.0490 (15)	0.0096 (16)	0.0252 (13)	-0.0016 (15)
C16	0.0642 (17)	0.0692 (19)	0.0500 (15)	0.0063 (14)	0.0288 (13)	-0.0089 (15)
C17	0.092 (2)	0.082 (2)	0.072 (2)	-0.0031 (19)	0.0344 (18)	-0.008 (2)
C18	0.141 (4)	0.102 (3)	0.112 (4)	-0.045 (3)	0.077 (4)	-0.033 (3)
C19	0.087 (3)	0.168 (5)	0.110 (4)	-0.052 (3)	0.055 (3)	-0.059 (4)
C20	0.060 (2)	0.149 (5)	0.097 (3)	-0.002 (2)	0.024 (2)	-0.029 (3)
C21	0.0680 (19)	0.085 (2)	0.074 (2)	0.0091 (17)	0.0248 (16)	-0.0135 (18)
N1	0.0555 (12)	0.0624 (13)	0.0476 (12)	-0.0017 (10)	0.0201 (10)	-0.0001 (11)

Geometric parameters (Å, °)

C1—C2	1.391 (4)	C11—C12	1.377 (4)
C1—C6	1.395 (4)	C11—H11	0.9300
C1—N1	1.405 (4)	C12—H12	0.9300
C2—C3	1.378 (5)	C13—N1	1.459 (3)
C2—H2	0.9300	C13—C14	1.471 (4)
C3—C4	1.368 (5)	C13—H13A	0.9700
C3—H3	0.9300	C13—H13B	0.9700
C4—C5	1.379 (5)	C14—C15	1.185 (4)
C4—H4	0.9300	C15—C16	1.441 (5)
C5—C6	1.372 (5)	C16—C17	1.377 (5)
C5—H5	0.9300	C16—C21	1.390 (4)
C6—H6	0.9300	C17—C18	1.386 (6)
C7—C8	1.374 (4)	C17—H17	0.9300
C7—C12	1.377 (4)	C18—C19	1.376 (8)
C7—N1	1.425 (3)	C18—H18	0.9300
C8—C9	1.383 (4)	C19—C20	1.344 (7)
C8—H8	0.9300	C19—H19	0.9300
C9—C10	1.377 (5)	C20—C21	1.358 (5)
C9—H9	0.9300	C20—H20	0.9300
C10—C11	1.362 (5)	C21—H21	0.9300
C10—H10	0.9300		
C2—C1—C6	117.6 (3)	C7—C12—C11	120.3 (3)
C2—C1—N1	122.7 (3)	C7—C12—H12	119.8
C6—C1—N1	119.6 (2)	C11—C12—H12	119.8
C3—C2—C1	120.2 (3)	N1—C13—C14	114.0 (2)

C3—C2—H2	119.9	N1—C13—H13A	108.7
C1—C2—H2	119.9	C14—C13—H13A	108.7
C4—C3—C2	121.8 (3)	N1—C13—H13B	108.7
C4—C3—H3	119.1	C14—C13—H13B	108.7
C2—C3—H3	119.1	H13A—C13—H13B	107.6
C3—C4—C5	118.3 (3)	C15—C14—C13	177.1 (3)
C3—C4—H4	120.9	C14—C15—C16	178.9 (3)
C5—C4—H4	120.9	C17—C16—C21	118.7 (3)
C6—C5—C4	120.9 (3)	C17—C16—C15	121.6 (3)
C6—C5—H5	119.5	C21—C16—C15	119.7 (3)
C4—C5—H5	119.5	C16—C17—C18	119.8 (4)
C5—C6—C1	121.1 (3)	C16—C17—H17	120.1
C5—C6—H6	119.5	C18—C17—H17	120.1
C1—C6—H6	119.5	C19—C18—C17	119.9 (5)
C8—C7—C12	118.9 (2)	C19—C18—H18	120.1
C8—C7—N1	122.0 (2)	C17—C18—H18	120.1
C12—C7—N1	119.1 (2)	C20—C19—C18	120.2 (4)
C7—C8—C9	120.6 (3)	C20—C19—H19	119.9
C7—C8—H8	119.7	C18—C19—H19	119.9
C9—C8—H8	119.7	C19—C20—C21	120.8 (4)
C10—C9—C8	119.9 (3)	C19—C20—H20	119.6
C10—C9—H9	120.1	C21—C20—H20	119.6
C8—C9—H9	120.1	C20—C21—C16	120.6 (4)
C11—C10—C9	119.5 (3)	C20—C21—H21	119.7
C11—C10—H10	120.3	C16—C21—H21	119.7
C9—C10—H10	120.3	C1—N1—C7	120.1 (2)
C10—C11—C12	120.8 (3)	C1—N1—C13	118.9 (2)
C10—C11—H11	119.6	C7—N1—C13	114.6 (2)
C12—C11—H11	119.6		
C6—C1—C2—C3	0.1 (4)	C14—C15—C16—C21	90 (18)
N1—C1—C2—C3	-176.6 (3)	C21—C16—C17—C18	-0.8 (4)
C1—C2—C3—C4	0.1 (5)	C15—C16—C17—C18	178.9 (3)
C2—C3—C4—C5	-0.6 (6)	C16—C17—C18—C19	0.6 (5)
C3—C4—C5—C6	0.9 (6)	C17—C18—C19—C20	0.1 (6)
C4—C5—C6—C1	-0.8 (5)	C18—C19—C20—C21	-0.6 (6)
C2—C1—C6—C5	0.2 (4)	C19—C20—C21—C16	0.4 (6)
N1—C1—C6—C5	177.1 (3)	C17—C16—C21—C20	0.3 (5)
C12—C7—C8—C9	0.6 (4)	C15—C16—C21—C20	-179.3 (3)
N1—C7—C8—C9	178.2 (2)	C2—C1—N1—C7	-148.9 (3)
C7—C8—C9—C10	-0.5 (4)	C6—C1—N1—C7	34.4 (3)
C8—C9—C10—C11	-0.2 (4)	C2—C1—N1—C13	1.4 (4)
C9—C10—C11—C12	1.0 (4)	C6—C1—N1—C13	-175.3 (3)
C8—C7—C12—C11	0.1 (4)	C8—C7—N1—C1	50.4 (3)
N1—C7—C12—C11	-177.5 (3)	C12—C7—N1—C1	-132.0 (3)
C10—C11—C12—C7	-0.9 (4)	C8—C7—N1—C13	-101.1 (3)
N1—C13—C14—C15	9(6)	C12—C7—N1—C13	76.4 (3)
C13—C14—C15—C16	136 (16)	C14—C13—N1—C1	-74.1 (3)
C14—C15—C16—C17	-89 (18)	C14—C13—N1—C7	77.8 (3)

Fig. 1

