

N-[2-(Aminocarbonyl)phenyl]-4-hydroxy-2-methyl-2H-1,2-benzothiazine-3-carboxamide 1,1-dioxide

Muhammad Nadeem Arshad,^a Muhammad Zia-ur-Rehman^{b*} and Islam Ullah Khan^a

^aDepartment of Chemistry, Government College University, Lahore 54000, Pakistan,

and ^bApplied Chemistry Research Centre, PCSIR Laboratories Complex, Ferozpur Road, Lahore 54600, Pakistan

Correspondence e-mail: rehman_pcsir@hotmail.com


Received 24 September 2009; accepted 25 September 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$; R factor = 0.087; wR factor = 0.213; data-to-parameter ratio = 15.8.

In the title compound, $C_{17}H_{15}N_3O_5S$, the thiazine ring adopts a distorted half-chair conformation. The molecular structure is stabilized by intramolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonding. Pairs of molecules are bound together as centrosymmetric dimers through $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For the synthesis of related molecules, see: Braun (1923); Ahmad *et al.* (2008); Zia-ur-Rehman *et al.* (2005, 2009). For the biological activity of 1,2-benzothiazine 1,1-dioxides, see: Bihovsky *et al.* (2004); Turck *et al.* (1996); Zia-ur-Rehman *et al.* (2006). For similar molecules, see: Kojić-Prodić & Ružić-Toroš (1982); Siddiqui *et al.* (2009); Weast *et al.* (1984); Zia-ur-Rehman *et al.* (2007).



Experimental

Crystal data



$M_r = 373.38$

Monoclinic, $P2_1/c$

$a = 8.1377(6)\text{ \AA}$

$b = 7.0515(6)\text{ \AA}$

$c = 29.069(2)\text{ \AA}$

$\beta = 96.502(3)^\circ$

$V = 1657.3(2)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.23\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.39 \times 0.25 \times 0.11\text{ mm}$

Data collection

Bruker APEXII CCD area-detector

diffractometer

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2007)

$T_{\min} = 0.915$, $T_{\max} = 0.975$

16109 measured reflections

3753 independent reflections

3011 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.087$

$wR(F^2) = 0.213$

$S = 1.09$

3753 reflections

237 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.38\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.40\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots\cdot A$	$D\cdots\cdot A$	$D-\text{H}\cdots\cdot A$
O1—H1 \cdots O4	0.82	1.85	2.569 (5)	145
N2—H2 \cdots O5	0.86	1.92	2.607 (5)	136
N2—H2 \cdots N1	0.86	2.28	2.728 (5)	113
N3—H3A \cdots O5 ⁱ	0.86	2.22	2.941 (6)	141

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors are grateful to the PCSIR Laboratories Complex, Lahore, for the provision of facilities necessary to complete this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5073).

References

- Ahmad, M., Siddiqui, H. L., Zia-ur-Rehman, M., Ashiq, M. I. & Tizzard, G. J. (2008). *Acta Cryst. E64*, o788.
Bihovsky, R., Tao, M., Mallamo, J. P. & Wells, G. J. (2004). *Bioorg. Med. Chem. Lett.* **14**, 1035–1038.
Braun, J. (1923). *Chem. Ber.* **56**, 2332–2343.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Kojić-Prodić, B. & Ružić-Toroš, Ž. (1982). *Acta Cryst. B38*, 2948–2951.
Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
Sheldrick, G. M. (2007). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Siddiqui, W. A., Ali, M., Zia-ur-Rehman, M., Sharif, S. & Tizzard, G. J. (2009). *Acta Cryst. E65*, o900–o901.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.
Turck, D., Busch, U., Heinzel, G., Narjes, H. & Nehmiz, G. (1996). *J. Clin. Pharmacol.* **36**, 79–84.
Weast, R. C., Astle, M. J. & Beyer, W. H. (1984). *Handbook of Chemistry and Physics*, 65th ed. Boca Raton, Florida: CRC Press.
Zia-ur-Rehman, M., Choudary, J. A. & Ahmad, S. (2005). *Bull. Korean Chem. Soc.* **26**, 1771–1775.
Zia-ur-Rehman, M., Choudary, J. A., Ahmad, S. & Siddiqui, H. L. (2006). *Chem. Pharm. Bull.* **54**, 1175–1178.
Zia-ur-Rehman, M., Choudary, J. A., Elsegood, M. R. J., Siddiqui, H. L. & Ahmad, S. (2007). *Acta Cryst. E63*, o900–o901.
Zia-ur-Rehman, M., Choudary, J. A., Elsegood, M. R. J., Siddiqui, H. L. & Khan, K. M. (2009). *Eur. J. Med. Chem.* **44**, 1311–1316.

Experimental

Crystal data



$M_r = 373.38$

Monoclinic, $P2_1/c$

$a = 8.1377(6)\text{ \AA}$

$b = 7.0515(6)\text{ \AA}$

$c = 29.069(2)\text{ \AA}$

$\beta = 96.502(3)^\circ$

$V = 1657.3(2)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 0.23\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.39 \times 0.25 \times 0.11\text{ mm}$

supplementary materials

Acta Cryst. (2009). E65, o2596 [doi:10.1107/S1600536809038951]

N-[2-(Aminocarbonyl)phenyl]-4-hydroxy-2-methyl-2H-1,2-benzothiazine-3-carboxamide 1,1-dioxide

M. N. Arshad, M. Zia-ur-Rehman and I. U. Khan

Comment

Owing to the verstaile applications of 1,2-benzothiazine 1,1-dioxides, considerable attention has been given to their synthesis since their very first synthesis (Braun, 1923). Among these, Piroxicam (Zia-ur-Rehman *et al.*, 2005), and Meloxicam (Turck *et al.*, 1996) are familiar for their analgesic action and are being used world wide as non-steroidal anti-inflammatory drugs (NSAIDs). Some of the 3,4-dihydro-1,2-benzothiazine-3-carboxylate 1,1-dioxide α -ketomide and P(2)—P(3) peptide mimetic aldehyde compounds act as potent calpain I inhibitors (Bihovsky *et al.*, 2004) while 1,2-benzothiazin-3-yl-quinazolin-4(3H)-ones possess anti-bacterial properties (Zia-ur-Rehman *et al.*, 2006).

In continuation of our work on the synthesis (Zia-ur-Rehman *et al.*, 2006, biological activity (Zia-ur-Rehman *et al.*, 2009) and crystal structures (Zia-ur-Rehman *et al.*, 2007; Ahmad *et al.*, 2008, Siddiqui *et al.*, 2009) of various 1,2-benzothiazine-1,1-dioxides, we herein report the crystal structure of the title compound (**I**) (Scheme and figure 1). Like its already reported dimethylsulfoxide solvate analogue (Zia-ur-Rehman *et al.*, 2007), thiazine ring involving two double bonds, exhibits sofa conformation; with S1/C1/C2/C7 relatively planar and N1 showing significant departure from plane due to its pyramidal geometry. The enolic hydrogen on O1 is involved in intramolecular hydrogen bonding [O1—H1 \cdots O4] with the carbonyl oxygen at C9 giving rise to a six-membered hydrogen bond ring (Table 1). Atom H2 forms hydrogen bonds with both N1 and O5 giving rise to five and six-membered hydrogen bond rings respectively. The C1—S1 [1.755 Å] bond is shorter than a normal C—S single bond (1.81–2.55 Å) (Weast *et al.*, 1984) due to partial double bond character and is in agreement with similar molecules (Kojić-Prodić & Ružić-Toroš, 1982). Each molecule is centrosymmetrically linked to its adjacent one forming a dimer through intermolecular [N—H3B \cdots O5] hydrogen bonds (Fig. 2).

Experimental

N-[2-(Aminocarbonyl)phenyl]-4-hydroxy-2-methyl-2*H*-1,2-benzothiazine-3-carboxamide 1,1-dioxide was synthesized according to a literature method (Zia-ur-Rehman *et al.*, 2006). Suitable crystals were obtained by dissoving the compound in chloroform followed by slow evaporation at room temperature. The compound was dissolved in a mixture of methanol and DMSO (80:20 v/v) at room temperature. Crystals were obtained by slow evaporation and dried under high vacuum.

Refinement

All hydrogen atoms were identified in the difference map. They were refined using a riding model with O—H = 0.84 Å, N—H = 0.86 Å, C_{methyl}—H 0.98 Å and C_{aromatic}—H = 0.95 Å. and U(H) set to 1.2Ueq of the parent atoms or set to 1.5Ueq(C_{methyl}).

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids at the 50% probability level.




Fig. 2. Perspective view of the three-dimensional crystal packing showing hydrogen-bonded interactions (dashed lines). H atoms not involved in hydrogen bonding have been omitted for clarity.

N-[2-(Aminocarbonyl)phenyl]-4-hydroxy-2-methyl-2H-1,2-benzothiazine-3-carboxamide 1,1-dioxide

Crystal data

C ₁₇ H ₁₅ N ₃ O ₅ S	$F_{000} = 776$
$M_r = 373.38$	$D_x = 1.496 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 6164 reflections
$a = 8.1377 (6) \text{ \AA}$	$\theta = 2.5\text{--}27.1^\circ$
$b = 7.0515 (6) \text{ \AA}$	$\mu = 0.23 \text{ mm}^{-1}$
$c = 29.069 (2) \text{ \AA}$	$T = 296 \text{ K}$
$\beta = 96.502 (3)^\circ$	Needles, colourless
$V = 1657.3 (2) \text{ \AA}^3$	$0.39 \times 0.25 \times 0.11 \text{ mm}$
$Z = 4$	

Data collection

Bruker APEXII CCD area-detector diffractometer	3753 independent reflections
Radiation source: fine-focus sealed tube	3011 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.039$
$T = 296 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ϕ and ω scans	$\theta_{\text{min}} = 2.5^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2007)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.915$, $T_{\text{max}} = 0.975$	$k = -9 \rightarrow 9$
16109 measured reflections	$l = -32 \rightarrow 37$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.087$	H-atom parameters constrained
$wR(F^2) = 0.213$	$w = 1/[\sigma^2(F_{\text{o}}^2) + (0.0335P)^2 + 9.427P]$
	where $P = (F_{\text{o}}^2 + 2F_{\text{c}}^2)/3$

$S = 1.09$	$(\Delta/\sigma)_{\max} < 0.001$
3753 reflections	$\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$
237 parameters	$\Delta\rho_{\min} = -0.40 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.35684 (14)	0.5031 (2)	0.33289 (4)	0.0343 (3)
O2	0.2977 (5)	0.6850 (5)	0.34450 (12)	0.0427 (9)
O3	0.5306 (4)	0.4668 (7)	0.33922 (13)	0.0538 (11)
O4	-0.1641 (4)	0.3152 (6)	0.38428 (12)	0.0480 (10)
O5	0.3670 (4)	0.3198 (6)	0.47838 (12)	0.0484 (10)
N1	0.2640 (5)	0.3427 (6)	0.36214 (13)	0.0310 (9)
N2	0.0743 (5)	0.2848 (6)	0.43247 (12)	0.0306 (8)
H2	0.1801	0.2826	0.4328	0.037*
N3	0.3804 (6)	0.4241 (8)	0.55160 (15)	0.0534 (13)
H3A	0.4817	0.4585	0.5513	0.064*
H3B	0.3319	0.4407	0.5761	0.064*
O1	-0.1513 (4)	0.3895 (6)	0.29823 (12)	0.0436 (9)
H1	-0.1968	0.3747	0.3218	0.065*
C1	0.2765 (6)	0.4493 (7)	0.27578 (16)	0.0329 (10)
C2	0.1069 (6)	0.4046 (7)	0.26901 (16)	0.0322 (10)
C3	0.0347 (8)	0.3766 (8)	0.22406 (17)	0.0445 (13)
H3	-0.0780	0.3516	0.2183	0.053*
C4	0.1301 (9)	0.3861 (9)	0.18781 (19)	0.0577 (17)
H4	0.0805	0.3685	0.1577	0.069*
C5	0.2966 (9)	0.4210 (10)	0.19540 (19)	0.0580 (17)
H5	0.3594	0.4221	0.1706	0.070*
C6	0.3718 (7)	0.4546 (9)	0.23979 (19)	0.0470 (14)
H6	0.4845	0.4802	0.2451	0.056*
C7	0.0124 (6)	0.3801 (7)	0.30885 (16)	0.0314 (10)
C8	0.0874 (6)	0.3479 (7)	0.35254 (16)	0.0308 (10)
C9	-0.0106 (6)	0.3133 (7)	0.39093 (15)	0.0315 (10)
C10	0.0150 (6)	0.2579 (6)	0.47583 (15)	0.0287 (9)

supplementary materials

C11	0.1236 (6)	0.2888 (7)	0.51660 (16)	0.0323 (10)
C12	0.0636 (7)	0.2643 (7)	0.55876 (17)	0.0402 (12)
H12	0.1350	0.2810	0.5858	0.048*
C13	-0.0980 (7)	0.2160 (8)	0.56205 (18)	0.0441 (13)
H13	-0.1358	0.2046	0.5909	0.053*
C14	-0.2039 (7)	0.1845 (8)	0.5223 (2)	0.0448 (13)
H14	-0.3134	0.1514	0.5245	0.054*
C15	-0.1485 (6)	0.2019 (7)	0.47945 (17)	0.0374 (11)
H15	-0.2197	0.1764	0.4528	0.045*
C16	0.2986 (6)	0.3449 (8)	0.51412 (16)	0.0368 (11)
C17	0.3393 (7)	0.1508 (9)	0.3632 (2)	0.0521 (15)
H17A	0.3025	0.0790	0.3882	0.078*
H17B	0.4576	0.1619	0.3678	0.078*
H17C	0.3065	0.0873	0.3344	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0273 (5)	0.0450 (7)	0.0309 (6)	-0.0003 (5)	0.0047 (4)	0.0006 (5)
O2	0.048 (2)	0.041 (2)	0.0391 (19)	-0.0070 (17)	0.0065 (16)	-0.0037 (16)
O3	0.0284 (18)	0.084 (3)	0.049 (2)	0.000 (2)	0.0067 (16)	0.010 (2)
O4	0.0289 (18)	0.079 (3)	0.0368 (19)	0.0027 (19)	0.0076 (14)	0.0050 (19)
O5	0.0350 (19)	0.075 (3)	0.0354 (19)	-0.0012 (19)	0.0067 (15)	-0.0072 (19)
N1	0.0278 (19)	0.038 (2)	0.0276 (19)	0.0065 (17)	0.0039 (15)	0.0006 (16)
N2	0.0260 (19)	0.039 (2)	0.0280 (19)	0.0001 (17)	0.0063 (15)	0.0053 (17)
N3	0.042 (3)	0.082 (4)	0.036 (2)	-0.006 (3)	0.0011 (19)	-0.009 (2)
O1	0.0307 (18)	0.065 (3)	0.0335 (18)	-0.0013 (18)	-0.0011 (14)	0.0024 (18)
C1	0.040 (3)	0.032 (2)	0.028 (2)	0.001 (2)	0.0079 (19)	0.0016 (18)
C2	0.037 (2)	0.031 (2)	0.029 (2)	-0.001 (2)	0.0064 (19)	0.0005 (19)
C3	0.063 (4)	0.042 (3)	0.029 (2)	-0.005 (3)	0.005 (2)	-0.002 (2)
C4	0.087 (5)	0.056 (4)	0.029 (3)	-0.013 (3)	0.004 (3)	-0.005 (3)
C5	0.076 (4)	0.071 (4)	0.031 (3)	-0.005 (4)	0.024 (3)	-0.004 (3)
C6	0.046 (3)	0.057 (4)	0.041 (3)	-0.003 (3)	0.016 (2)	-0.001 (3)
C7	0.030 (2)	0.034 (2)	0.030 (2)	0.0008 (19)	0.0067 (18)	-0.0015 (19)
C8	0.028 (2)	0.034 (2)	0.030 (2)	0.0032 (19)	0.0029 (17)	0.0000 (19)
C9	0.034 (2)	0.032 (2)	0.029 (2)	0.0000 (19)	0.0034 (18)	-0.0003 (19)
C10	0.032 (2)	0.026 (2)	0.028 (2)	0.0039 (18)	0.0082 (18)	0.0025 (18)
C11	0.039 (3)	0.028 (2)	0.031 (2)	0.004 (2)	0.0072 (19)	0.0029 (19)
C12	0.058 (3)	0.035 (3)	0.029 (2)	0.002 (2)	0.011 (2)	0.001 (2)
C13	0.063 (4)	0.039 (3)	0.035 (3)	0.000 (3)	0.023 (2)	0.003 (2)
C14	0.045 (3)	0.039 (3)	0.054 (3)	-0.002 (2)	0.021 (3)	0.005 (3)
C15	0.036 (3)	0.039 (3)	0.039 (3)	0.000 (2)	0.012 (2)	0.007 (2)
C16	0.034 (3)	0.044 (3)	0.031 (2)	0.006 (2)	-0.0011 (19)	0.003 (2)
C17	0.052 (3)	0.049 (3)	0.058 (4)	0.020 (3)	0.020 (3)	0.014 (3)

Geometric parameters (\AA , $^\circ$)

S1—O2	1.424 (4)	C4—C5	1.371 (9)
S1—O3	1.428 (4)	C4—H4	0.9300

S1—N1	1.648 (4)	C5—C6	1.384 (8)
S1—C1	1.755 (5)	C5—H5	0.9300
O4—C9	1.242 (6)	C6—H6	0.9300
O5—C16	1.246 (6)	C7—C8	1.364 (6)
N1—C8	1.433 (6)	C8—C9	1.464 (6)
N1—C17	1.484 (7)	C10—C15	1.403 (6)
N2—C9	1.337 (6)	C10—C11	1.413 (6)
N2—C10	1.412 (5)	C11—C12	1.380 (6)
N2—H2	0.8600	C11—C16	1.488 (7)
N3—C16	1.333 (6)	C12—C13	1.372 (8)
N3—H3A	0.8600	C12—H12	0.9300
N3—H3B	0.8600	C13—C14	1.378 (8)
O1—C7	1.335 (5)	C13—H13	0.9300
O1—H1	0.8200	C14—C15	1.379 (7)
C1—C6	1.372 (7)	C14—H14	0.9300
C1—C2	1.407 (7)	C15—H15	0.9300
C2—C3	1.385 (7)	C17—H17A	0.9600
C2—C7	1.471 (6)	C17—H17B	0.9600
C3—C4	1.379 (8)	C17—H17C	0.9600
C3—H3	0.9300		
O2—S1—O3	119.2 (3)	O1—C7—C2	114.2 (4)
O2—S1—N1	108.0 (2)	C8—C7—C2	122.3 (4)
O3—S1—N1	108.4 (2)	C7—C8—N1	121.2 (4)
O2—S1—C1	108.6 (2)	C7—C8—C9	120.8 (4)
O3—S1—C1	109.9 (2)	N1—C8—C9	117.9 (4)
N1—S1—C1	101.3 (2)	O4—C9—N2	123.4 (4)
C8—N1—C17	115.4 (4)	O4—C9—C8	120.3 (4)
C8—N1—S1	113.0 (3)	N2—C9—C8	116.3 (4)
C17—N1—S1	115.1 (3)	C15—C10—N2	121.8 (4)
C9—N2—C10	129.2 (4)	C15—C10—C11	119.3 (4)
C9—N2—H2	115.4	N2—C10—C11	118.9 (4)
C10—N2—H2	115.4	C12—C11—C10	118.4 (5)
C16—N3—H3A	120.0	C12—C11—C16	120.9 (5)
C16—N3—H3B	120.0	C10—C11—C16	120.8 (4)
H3A—N3—H3B	120.0	C13—C12—C11	122.1 (5)
C7—O1—H1	109.5	C13—C12—H12	119.0
C6—C1—C2	122.0 (5)	C11—C12—H12	119.0
C6—C1—S1	122.2 (4)	C12—C13—C14	119.7 (5)
C2—C1—S1	115.8 (3)	C12—C13—H13	120.2
C3—C2—C1	118.0 (5)	C14—C13—H13	120.2
C3—C2—C7	121.5 (5)	C13—C14—C15	120.4 (5)
C1—C2—C7	120.5 (4)	C13—C14—H14	119.8
C4—C3—C2	119.9 (6)	C15—C14—H14	119.8
C4—C3—H3	120.1	C14—C15—C10	120.2 (5)
C2—C3—H3	120.1	C14—C15—H15	119.9
C5—C4—C3	121.2 (5)	C10—C15—H15	119.9
C5—C4—H4	119.4	O5—C16—N3	120.8 (5)
C3—C4—H4	119.4	O5—C16—C11	121.6 (4)
C4—C5—C6	120.4 (5)	N3—C16—C11	117.6 (4)

supplementary materials


C4—C5—H5	119.8	N1—C17—H17A	109.5
C6—C5—H5	119.8	N1—C17—H17B	109.5
C1—C6—C5	118.5 (5)	H17A—C17—H17B	109.5
C1—C6—H6	120.7	N1—C17—H17C	109.5
C5—C6—H6	120.7	H17A—C17—H17C	109.5
O1—C7—C8	123.6 (4)	H17B—C17—H17C	109.5
O2—S1—N1—C8	58.6 (4)	O1—C7—C8—C9	2.6 (8)
O3—S1—N1—C8	-171.0 (3)	C2—C7—C8—C9	-176.4 (4)
C1—S1—N1—C8	-55.4 (4)	C17—N1—C8—C7	-96.6 (6)
O2—S1—N1—C17	-165.8 (4)	S1—N1—C8—C7	38.8 (6)
O3—S1—N1—C17	-35.4 (4)	C17—N1—C8—C9	82.0 (5)
C1—S1—N1—C17	80.2 (4)	S1—N1—C8—C9	-142.6 (4)
O2—S1—C1—C6	106.1 (5)	C10—N2—C9—O4	-2.0 (8)
O3—S1—C1—C6	-25.9 (5)	C10—N2—C9—C8	176.5 (4)
N1—S1—C1—C6	-140.4 (5)	C7—C8—C9—O4	-0.9 (8)
O2—S1—C1—C2	-72.5 (4)	N1—C8—C9—O4	-179.6 (5)
O3—S1—C1—C2	155.5 (4)	C7—C8—C9—N2	-179.5 (5)
N1—S1—C1—C2	41.0 (4)	N1—C8—C9—N2	1.9 (7)
C6—C1—C2—C3	-3.9 (8)	C9—N2—C10—C15	19.3 (8)
S1—C1—C2—C3	174.7 (4)	C9—N2—C10—C11	-160.5 (5)
C6—C1—C2—C7	173.4 (5)	C15—C10—C11—C12	-0.6 (7)
S1—C1—C2—C7	-8.0 (6)	N2—C10—C11—C12	179.2 (4)
C1—C2—C3—C4	2.4 (8)	C15—C10—C11—C16	178.9 (4)
C7—C2—C3—C4	-174.9 (5)	N2—C10—C11—C16	-1.3 (7)
C2—C3—C4—C5	0.7 (10)	C10—C11—C12—C13	-1.7 (8)
C3—C4—C5—C6	-2.4 (11)	C16—C11—C12—C13	178.8 (5)
C2—C1—C6—C5	2.2 (9)	C11—C12—C13—C14	2.2 (8)
S1—C1—C6—C5	-176.3 (5)	C12—C13—C14—C15	-0.2 (8)
C4—C5—C6—C1	0.9 (10)	C13—C14—C15—C10	-2.1 (8)
C3—C2—C7—O1	-20.1 (7)	N2—C10—C15—C14	-177.3 (5)
C1—C2—C7—O1	162.7 (5)	C11—C10—C15—C14	2.5 (7)
C3—C2—C7—C8	159.0 (5)	C12—C11—C16—O5	160.7 (5)
C1—C2—C7—C8	-18.2 (7)	C10—C11—C16—O5	-18.7 (8)
O1—C7—C8—N1	-178.8 (4)	C12—C11—C16—N3	-19.5 (7)
C2—C7—C8—N1	2.1 (7)	C10—C11—C16—N3	161.1 (5)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1···O4	0.82	1.85	2.569 (5)	145
N2—H2···O5	0.86	1.92	2.607 (5)	136
N2—H2···N1	0.86	2.28	2.728 (5)	113
N3—H3A···O5 ⁱ	0.86	2.22	2.941 (6)	141

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Fig. 1



supplementary materials

Fig. 2

