

Carbonyl[4-(2,3-dimethylphenylamino)-pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I)

Gertruida J. S. Venter,* Gideon Steyl and Andreas Roodt

Department of Chemistry, University of the Free State, PO Box 339, Bloemfontein 9300, South Africa

Correspondence e-mail: truidie@hotmail.com


Received 5 August 2009; accepted 30 September 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.004$ Å; disorder in main residue; R factor = 0.033; wR factor = 0.076; data-to-parameter ratio = 22.9.

In the title compound, $[Rh(C_{13}H_{16}NO)(C_{18}H_{15}P)(CO)]$, the coordination geometry of the Rh^I atom is square-planar, formed by the coordinating N and O atoms of the bidentate enaminoketonate ligand, one C atom from the carbonyl group and a P atom from triphenylphosphine. The complex displays a 0.591 (3):0.409 (3) ratio disorder of the phenyl unit of the monoanionic N,O -bidentate ligand. Intramolecular hydrogen bonding is observed between a C—H group of the triphenylphosphine unit and the O atom of the enaminoketonate ligand.

Related literature

For related derivatives of the 4-phenylaminopent-3-en-2-onate ligand, see: Da Silva *et al.* (1993); Gordon *et al.* (2002); Shaheen *et al.* (2006). For related dicarbonyl rhodium(I) complexes with a bidentate ligand, see: Cornils & Herrmann (1996); Trzeciak & Ziólkowski (1994); van Rooy *et al.* (1995). For related carbonyl rhodium(I) complexes with a phosphine and a bidentate ligand, see: Bonati & Wilkinson (1964); Damoense *et al.* (1994); Lamprecht *et al.* (1997); Leipoldt *et al.* (1978); Purcell *et al.* (1995); Varshavsky *et al.* (2001). For background information, see: Tolman (1977).



Experimental

Crystal data

$[Rh(C_{13}H_{16}NO)(C_{18}H_{15}P)(CO)]$	$V = 2770.88$ (11) Å ³
$M_r = 595.46$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 14.9077$ (3) Å	$\mu = 0.70$ mm ⁻¹
$b = 11.6202$ (3) Å	$T = 100$ K
$c = 16.0256$ (4) Å	$0.25 \times 0.15 \times 0.13$ mm
$\beta = 93.521$ (1)°	

Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer	30367 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2001)	6985 independent reflections
$T_{min} = 0.844$, $T_{max} = 0.914$	5783 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.040$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$	305 parameters
$wR(F^2) = 0.076$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{max} = 0.60$ e Å ⁻³
6985 reflections	$\Delta\rho_{min} = -0.69$ e Å ⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C326—H326 \cdots O12	0.95	2.36	3.177 (3)	143

Table 2

Comparative geometrical parameters for similar $[Rh(N,O-bid)(CO)-(PPh_3)]$ complexes (Å, °).

Parameters	(I)	(II)	(III)
Rh1—N11	2.069 (2)	2.045 (4)	2.045 (3)
Rh1—O12	2.028 (2)	2.044 (3)	2.045 (2)
Rh1—P13	2.2635 (6)	2.275 (1)	2.281 (2)
Rh1—C14	1.807 (2)	1.784 (5)	1.804 (3)
C14—O14	1.152 (3)	1.142 (7)	1.148 (4)
N11 \cdots O12	2.885 (3)	2.826 (6)	2.841 (3)
N11—Rh1—O12	89.54 (8)	87.4 (1)	87.95 (8)
O12—Rh1—P13	84.97 (5)	89.7 (1)	89.91 (5)
P13—Rh1—C14	91.87 (7)	90.3 (2)	89.48 (9)
N11—Rh1—C14	93.6 (1)	92.6 (2)	92.6 (1)
N11—C2—C4—O12	4.1 (2)	1.2 (4)	1.5 (2)
θ_E^a	156.39 (3)	156.0 (2)	156.23 (4)

Notes: (I) This work. (II) N,O -bid = 4-aminopent-3-en-2-onato (Damoense *et al.*, 1994). (III) N,O -bid = 4-amino-1,1,1-trifluoropent-3-en-2-onato (Varshavsky *et al.*, 2001). (a) Tolman (1977)

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT-Plus (Bruker, 2007); data reduction: SAINT-Plus; program(s) used to solve structure: SIR92 (Altomare *et al.*, 1994); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg & Putz, 1999); software used to prepare material for publication: WinGX (Farrugia, 1999).

Financial assistance from the University of the Free State is gratefully acknowledged, while Mr Leo Kirsten is thanked for the XRD data collections. We also express our gratitude towards SASOL and the South African National Research Foundation (SA-NRF/THRIP) for financial support of this

project. Part of this material is based on work supported by the SA-NRF/THRIP under grant No. GUN 2068915. Opinions, findings, conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the SA-NRF.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2218).

References

- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Bonati, F. & Wilkinson, G. (1964). *J. Chem. Soc.* pp. 3156–3160.
- Brandenburg, K. & Putz, H. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT-Plus*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cornils, B. & Herrmann, W. A. (1996). In *Applied Homogeneous Catalysis with Organometallic Compounds: A Comprehensive Handbook*. Weinheim: VCH.
- Damoense, L. J., Purcell, W., Roodt, A. & Leipoldt, J. G. (1994). *Rhodium Express*, **5**, 10–13.
- Da Silva, M. A. V. R., Da Silva, M. D. M. C. R., Paiva, J. P. A., Nogueira, I. M. C. S., Damas, A. M., Barkley, J. V., Harding, M. M., Akello, M. J. & Pilcher, G. (1993). *J. Chem. Soc. Perkin Trans. 2*, pp. 1765–1769.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gordon, J. C., Shukla, P., Cowley, A. H., Jones, J. N., Keogh, D. W. & Scott, B. L. (2002). *Chem. Commun.* pp. 2710–2711.
- Lamprecht, D., Lamprecht, G. J., Botha, J. M., Umakoshi, K. & Sasaki, Y. (1997). *Acta Cryst.* **C53**, 1403–1405.
- Leipoldt, J. G., Basson, S. S., Bok, L. D. C. & Gerber, T. I. A. (1978). *Inorg. Chim. Acta*, **26**, L35–L37.
- Purcell, W., Basson, S. S., Leipoldt, J. G., Roodt, A. & Preston, H. (1995). *Inorg. Chim. Acta*, **234**, 153–156.
- Rooy, A. van, Orij, E. N., Kamer, P. C. J. & van Leeuwen, P. W. M. N. (1995). *Organometallics*, **14**, 34–43.
- Shaheen, F., Marchio, L., Badshah, A. & Khosa, M. K. (2006). *Acta Cryst.* **E62**, o873–o874.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Tolman, C. A. (1977). *Chem. Rev.* **77**, 313–348.
- Trzeciak, A. M. & Ziólkowski, J. J. (1994). *J. Organomet. Chem.* **464**, 107–111.
- Varshavsky, Y. S., Galding, M. R., Cherkasova, T. G., Podkorytov, I. S., Nikol'skii, A. B., Trzeciak, A. M., Olejnik, Z., Lis, T. & Ziólkowski, J. J. (2001). *J. Organomet. Chem.* **628**, 195–210.

supplementary materials

Acta Cryst. (2009). E65, m1321-m1322 [doi:10.1107/S1600536809039816]

Carbonyl[4-(2,3-dimethylphenylamino)pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I)

G. J. S. Venter, G. Steyl and A. Roodt

Comment

Rhodium(I) dicarbonyl complexes of the type $[\text{Rh}(L,L')(\text{CO})_2]$ containing chelating mono-anionic bidentate (L,L') ligands coordinated to rhodium *via* (O,O) donor atoms have been studied as catalyst precursors (Cornils & Herrmann, 1996; Trzeciak & Ziólkowski, 1994; van Rooy *et al.*, 1995). In this study the investigation of these β -diketonato complexes is followed by complexes containing bidentate β -enaminoketonato ligands such as 4-(phenylamino)pent-3-en-2-onato (Phony) (Shaheen *et al.*, 2006) coordinated to rhodium *via* (N,O) donor atoms. Dicarbonyl complexes of the $[\text{Rh}(N,O\text{-bid})(\text{CO})_2]$ -type (Varshavsky *et al.*, 2001) react with phosphorus ligands to form $[\text{Rh}(N,O\text{-bid})(\text{CO})(\text{PZ}_3)]$ complexes (Damoense *et al.*, 1994; Varshavsky *et al.*, 2001). According to Bonati & Wilkinson (1964), only one CO group will be substituted by triphenylphosphine, with the product being one of two possible isomers. Since N atom has a larger *trans*-influence than O atom, the CO group *trans* to the N atom will be substituted. This is evident in the title compound (Fig. 1), where $[\text{Rh}(2,3\text{-diMe-Phony})(\text{CO})(\text{PPh}_3)]$ is formed by the substitution of the carbonyl ligand in the dicarbonyl rhodium(I) complex $[\text{Rh}(2,3\text{-diMe-Phony})(\text{CO})_2]$ by PPh_3 .

Bond distances involving Rh atom in the title complex differ significantly from the distances in related complexes (Table 2). The Rh—N bond distance in the title complex is longer than those in similar complexes while the Rh—O bond distance is shorter. This is due to the steric influence of the phenyl group connected to N atom in the title compound, as opposed to H atom in the related complexes. The Rh—C and the carbonyl C—O bond distances do not differ substantially from the distances in the related complexes (Table 2). The N—Rh—O bite angle is slightly larger than those observed in similar complexes found in literature. The effective cone angle, θ_E (Tolman, 1977), of $156.39(3)^\circ$ is similar to the angles in the related compounds. The title complex displays a disorder of the phenyl ring in a 59:41% ratio.

Experimental

To a 5 ml acetone solution of $[\text{Rh}(2,3\text{-diMe-Phony})(\text{CO})_2]$ (0.0204 g, 56.48 μmol) was added PPh_3 (0.0151 g, 57.57 μmol) resulting in the immediate evolution of gas. Crystallization from acetone produced yellow crystals in quantitative yield (0.0334 g). IR (KBr): ν_{CO} 1966.93 cm^{-1} .

Refinement

The methyl and aromatic H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H = 0.98 and 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ and $1.2U_{\text{eq}}(\text{C})$, respectively. The methyl groups were generated to fit the difference electron density and the groups were then refined as rigid rotors.

Figures




Fig. 1. Molecular structure of the title compound. Displacement ellipsoids are drawn at the 50% probability level. H atoms have been omitted for clarity.




Fig. 2. Crystal packing of the title compound.

Carbonyl[4-(2,3-dimethylphenylamino)pent-3-en-2-onato- κ^2N,O](triphenylphosphine- κP)rhodium(I)

Crystal data

[Rh(C₁₃H₁₆NO)(C₁₈H₁₅P)(CO)]

$M_r = 595.46$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 14.9077$ (3) Å

$b = 11.6202$ (3) Å

$c = 16.0256$ (4) Å

$\beta = 93.521$ (1)°

$V = 2770.88$ (11) Å³

$Z = 4$

$F_{000} = 1224$

$D_x = 1.427$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8984 reflections

$\theta = 2.2$ – 28.4 °

$\mu = 0.70$ mm⁻¹

$T = 100$ K

Cuboid, yellow

$0.25 \times 0.15 \times 0.13$ mm

Data collection

Bruker X8 APEXII 4K Kappa CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 100$ K

ω and φ scans

Absorption correction: multi-scan (SADABS; Bruker, 2001)

$T_{\min} = 0.844$, $T_{\max} = 0.914$

30367 measured reflections

6985 independent reflections

5783 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

$\theta_{\max} = 28.5$ °

$\theta_{\min} = 1.4$ °

$h = -19 \rightarrow 19$

$k = -15 \rightarrow 14$

$l = -21 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.033$	H-atom parameters constrained
$wR(F^2) = 0.076$	$w = 1/[\sigma^2(F_o^2) + (0.0235P)^2 + 2.8578P]$
$S = 1.04$	where $P = (F_o^2 + 2F_c^2)/3$
6985 reflections	$(\Delta/\sigma)_{\max} = 0.005$
305 parameters	$\Delta\rho_{\max} = 0.60 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.69 \text{ e } \text{\AA}^{-3}$
	Extinction correction: none

Special details

Experimental. The intensity data was collected on a Bruker X8 APEXII 4 K Kappa CCD diffractometer using an exposure time of 60 s/frame. A total of 1033 frames were collected with a frame width of 0.5° covering up to $\theta = 28.41^\circ$ with 99.4% completeness accomplished.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Rh1	0.258727 (11)	0.605541 (15)	1.062396 (10)	0.01882 (6)	
N11	0.26580 (15)	0.43296 (18)	1.03172 (15)	0.0361 (5)	
O12	0.36554 (10)	0.63702 (14)	0.99328 (9)	0.0221 (3)	
O14	0.09801 (11)	0.57318 (15)	1.15945 (10)	0.0285 (4)	
P13	0.26175 (4)	0.79721 (5)	1.08760 (3)	0.01818 (12)	
C1	0.3204 (2)	0.2591 (2)	0.9641 (2)	0.0576 (10)	
H1A	0.3099	0.218	1.0161	0.086*	
H1B	0.3782	0.2352	0.9439	0.086*	
H1C	0.2722	0.2409	0.9219	0.086*	
C2	0.32179 (18)	0.3869 (2)	0.98045 (17)	0.0325 (6)	
C3	0.38454 (16)	0.4508 (2)	0.93713 (15)	0.0267 (5)	
H3	0.4176	0.4093	0.8982	0.032*	
C4	0.40317 (14)	0.5655 (2)	0.94473 (14)	0.0222 (5)	
C5	0.47221 (16)	0.6203 (2)	0.89254 (16)	0.0318 (6)	
H5A	0.4419	0.6684	0.8492	0.048*	
H5B	0.5068	0.5601	0.8662	0.048*	
H5C	0.5129	0.668	0.9283	0.048*	
C11A	0.22412 (17)	0.3512 (2)	1.08695 (16)	0.0249 (5)	0.591 (3)
C12A	0.13769 (16)	0.3169 (2)	1.06046 (13)	0.0249 (5)	0.591 (3)
C13A	0.09120 (13)	0.2400 (2)	1.10833 (15)	0.0249 (5)	0.591 (3)
C14A	0.13113 (16)	0.1974 (2)	1.18269 (14)	0.0249 (5)	0.591 (3)
H14A	0.0994	0.1449	1.2154	0.03*	0.591 (3)
C15A	0.21756 (17)	0.2318 (2)	1.20919 (14)	0.0249 (5)	0.591 (3)
H15A	0.2449	0.2027	1.26	0.03*	0.591 (3)

supplementary materials

C16A	0.26405 (14)	0.3086 (3)	1.16132 (17)	0.0249 (5)	0.591 (3)
H16A	0.3231	0.3321	1.1794	0.03*	0.591 (3)
C18A	-0.0011 (3)	0.1950 (4)	1.0797 (2)	0.0249 (5)	0.591 (3)
H18A	0.0043	0.1423	1.0325	0.037*	0.591 (3)
H18B	-0.0401	0.2596	1.0622	0.037*	0.591 (3)
H18C	-0.0271	0.1539	1.1258	0.037*	0.591 (3)
C17A	0.0953 (3)	0.3664 (4)	0.9799 (3)	0.0249 (5)	0.591 (3)
H17A	0.0902	0.4501	0.9853	0.037*	0.591 (3)
H17B	0.0354	0.333	0.9688	0.037*	0.591 (3)
H17C	0.1329	0.348	0.9337	0.037*	0.591 (3)
C11B	0.1803 (2)	0.3671 (3)	1.0474 (2)	0.0243 (7)	0.409 (3)
C12B	0.18861 (19)	0.3040 (3)	1.1212 (2)	0.0243 (7)	0.409 (3)
C13B	0.1158 (2)	0.2418 (3)	1.14741 (19)	0.0243 (7)	0.409 (3)
C14B	0.0346 (2)	0.2428 (3)	1.0998 (2)	0.0243 (7)	0.409 (3)
H14B	-0.0152	0.2003	1.1178	0.029*	0.409 (3)
C15B	0.0263 (2)	0.3059 (3)	1.0261 (2)	0.0243 (7)	0.409 (3)
H15B	-0.0292	0.3065	0.9936	0.029*	0.409 (3)
C16B	0.0991 (3)	0.3680 (3)	0.99988 (19)	0.0243 (7)	0.409 (3)
H16B	0.0934	0.4111	0.9495	0.029*	0.409 (3)
C17B	0.2760 (4)	0.3064 (6)	1.1737 (4)	0.0243 (7)	0.409 (3)
H17D	0.3093	0.2351	1.165	0.036*	0.409 (3)
H17E	0.2636	0.3132	1.2328	0.036*	0.409 (3)
H17F	0.3119	0.3724	1.1574	0.036*	0.409 (3)
C18B	0.1219 (4)	0.1738 (5)	1.2287 (3)	0.0243 (7)	0.409 (3)
H18D	0.0624	0.144	1.2399	0.036*	0.409 (3)
H18E	0.1436	0.2241	1.2747	0.036*	0.409 (3)
H18F	0.1638	0.1094	1.2238	0.036*	0.409 (3)
C14	0.16135 (15)	0.58652 (19)	1.12283 (14)	0.0215 (5)	
C311	0.19823 (15)	0.8522 (2)	1.17253 (14)	0.0212 (5)	
C312	0.23991 (16)	0.8923 (2)	1.24725 (15)	0.0261 (5)	
H312	0.3036	0.898	1.2531	0.031*	
C313	0.18891 (18)	0.9239 (2)	1.31310 (16)	0.0336 (6)	
H313	0.2177	0.9501	1.3642	0.04*	
C314	0.09599 (18)	0.9173 (2)	1.30437 (17)	0.0372 (6)	
H314	0.0611	0.9395	1.3493	0.045*	
C315	0.05421 (17)	0.8787 (2)	1.23056 (18)	0.0364 (6)	
H315	-0.0095	0.8745	1.2248	0.044*	
C316	0.10419 (16)	0.8460 (2)	1.16490 (16)	0.0289 (5)	
H316	0.0748	0.8192	1.1143	0.035*	
C321	0.22426 (16)	0.8831 (2)	0.99645 (14)	0.0239 (5)	
C322	0.17703 (18)	0.9858 (2)	1.00150 (15)	0.0329 (6)	
H322	0.1615	1.014	1.0544	0.04*	
C323	0.1525 (2)	1.0472 (2)	0.92915 (17)	0.0409 (7)	
H323	0.1194	1.1167	0.9329	0.049*	
C324	0.1756 (2)	1.0084 (2)	0.85193 (16)	0.0383 (6)	
H324	0.1591	1.0511	0.8028	0.046*	
C325	0.22301 (19)	0.9068 (2)	0.84678 (16)	0.0362 (6)	
H325	0.2393	0.8796	0.7939	0.043*	
C326	0.24695 (16)	0.8442 (2)	0.91864 (14)	0.0280 (5)	

H326	0.2791	0.7741	0.9144	0.034*
C331	0.37540 (15)	0.8504 (2)	1.11391 (13)	0.0224 (5)
C332	0.40227 (19)	0.9603 (2)	1.09345 (18)	0.0389 (7)
H332	0.3623	1.0096	1.0618	0.047*
C333	0.4886 (2)	0.9988 (3)	1.1196 (2)	0.0510 (8)
H333	0.5073	1.074	1.1053	0.061*
C334	0.54615 (18)	0.9282 (3)	1.16566 (18)	0.0431 (8)
H334	0.6043	0.955	1.1839	0.052*
C335	0.52026 (16)	0.8193 (3)	1.18548 (16)	0.0353 (6)
H335	0.5605	0.7705	1.2172	0.042*
C336	0.43533 (15)	0.7797 (2)	1.15942 (14)	0.0264 (5)
H336	0.4181	0.7035	1.1729	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Rh1	0.01877 (9)	0.01664 (9)	0.02180 (9)	-0.00199 (7)	0.00727 (6)	-0.00079 (7)
N11	0.0465 (13)	0.0177 (10)	0.0478 (13)	-0.0072 (9)	0.0327 (11)	-0.0047 (9)
O12	0.0197 (8)	0.0216 (8)	0.0256 (8)	0.0005 (6)	0.0075 (6)	0.0035 (6)
O14	0.0241 (9)	0.0315 (10)	0.0309 (9)	-0.0077 (7)	0.0108 (7)	-0.0045 (7)
P13	0.0172 (3)	0.0182 (3)	0.0191 (3)	-0.0007 (2)	0.0002 (2)	-0.0006 (2)
C1	0.083 (2)	0.0238 (15)	0.073 (2)	-0.0050 (15)	0.055 (2)	-0.0111 (15)
C2	0.0384 (14)	0.0220 (13)	0.0393 (14)	-0.0005 (11)	0.0201 (12)	-0.0043 (11)
C3	0.0271 (12)	0.0248 (13)	0.0297 (12)	0.0038 (10)	0.0134 (10)	-0.0004 (10)
C4	0.0169 (10)	0.0277 (12)	0.0223 (11)	0.0033 (9)	0.0038 (9)	0.0054 (9)
C5	0.0247 (12)	0.0372 (15)	0.0349 (13)	-0.0004 (11)	0.0128 (10)	0.0068 (11)
C11A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C12A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C13A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C14A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C15A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C16A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C18A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C17A	0.0258 (8)	0.0232 (8)	0.0259 (8)	-0.0024 (6)	0.0018 (6)	0.0011 (6)
C11B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C12B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C13B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C14B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C15B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C16B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C17B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C18B	0.0281 (12)	0.0204 (11)	0.0246 (11)	0.0016 (9)	0.0036 (9)	0.0003 (8)
C14	0.0238 (11)	0.0192 (12)	0.0216 (11)	-0.0045 (9)	0.0027 (9)	-0.0031 (9)
C311	0.0201 (11)	0.0193 (11)	0.0242 (11)	0.0002 (9)	0.0009 (9)	-0.0020 (9)
C312	0.0232 (12)	0.0279 (13)	0.0267 (12)	0.0035 (10)	-0.0017 (9)	-0.0048 (10)
C313	0.0363 (14)	0.0400 (16)	0.0242 (12)	0.0017 (12)	-0.0009 (11)	-0.0116 (11)
C314	0.0320 (14)	0.0462 (17)	0.0345 (14)	0.0044 (12)	0.0106 (11)	-0.0135 (12)
C315	0.0201 (12)	0.0439 (17)	0.0459 (16)	-0.0004 (11)	0.0072 (11)	-0.0112 (13)

supplementary materials

C316	0.0235 (12)	0.0312 (14)	0.0319 (13)	-0.0013 (10)	0.0007 (10)	-0.0085 (11)
C321	0.0280 (12)	0.0199 (12)	0.0231 (11)	-0.0001 (10)	-0.0034 (9)	-0.0003 (9)
C322	0.0483 (16)	0.0230 (13)	0.0267 (12)	0.0066 (12)	-0.0043 (11)	-0.0046 (10)
C323	0.0585 (19)	0.0243 (14)	0.0386 (15)	0.0148 (13)	-0.0064 (14)	0.0002 (11)
C324	0.0529 (18)	0.0312 (15)	0.0298 (13)	0.0095 (13)	-0.0065 (12)	0.0074 (11)
C325	0.0469 (16)	0.0383 (16)	0.0230 (12)	0.0103 (13)	-0.0008 (11)	0.0022 (11)
C326	0.0340 (14)	0.0248 (13)	0.0246 (12)	0.0081 (10)	-0.0023 (10)	0.0009 (10)
C331	0.0200 (11)	0.0286 (12)	0.0190 (10)	-0.0059 (9)	0.0035 (9)	-0.0027 (9)
C332	0.0405 (16)	0.0334 (16)	0.0423 (15)	-0.0137 (12)	-0.0020 (13)	0.0053 (12)
C333	0.0503 (19)	0.0486 (19)	0.0550 (19)	-0.0328 (16)	0.0110 (15)	-0.0046 (16)
C334	0.0240 (13)	0.067 (2)	0.0387 (15)	-0.0154 (14)	0.0085 (12)	-0.0198 (15)
C335	0.0194 (12)	0.0592 (19)	0.0276 (12)	0.0007 (12)	0.0031 (10)	-0.0165 (12)
C336	0.0212 (11)	0.0368 (14)	0.0215 (11)	-0.0014 (10)	0.0050 (9)	-0.0095 (10)

Geometric parameters (Å, °)

Rh1—C14	1.807 (2)	C13B—C18B	1.522 (6)
Rh1—O12	2.0280 (15)	C14B—C15B	1.39
Rh1—N11	2.069 (2)	C14B—H14B	0.95
Rh1—P13	2.2635 (6)	C15B—C16B	1.39
N11—C2	1.320 (3)	C15B—H15B	0.95
N11—C11A	1.463 (3)	C16B—H16B	0.95
N11—C11B	1.522 (3)	C17B—H17D	0.98
O12—C4	1.290 (3)	C17B—H17E	0.98
O14—C14	1.152 (3)	C17B—H17F	0.98
P13—C311	1.821 (2)	C18B—H18D	0.98
P13—C331	1.828 (2)	C18B—H18E	0.98
P13—C321	1.828 (2)	C18B—H18F	0.98
C1—C2	1.508 (4)	C311—C312	1.395 (3)
C1—H1A	0.98	C311—C316	1.401 (3)
C1—H1B	0.98	C312—C313	1.388 (3)
C1—H1C	0.98	C312—H312	0.95
C2—C3	1.410 (3)	C313—C314	1.386 (4)
C3—C4	1.366 (3)	C313—H313	0.95
C3—H3	0.95	C314—C315	1.378 (4)
C4—C5	1.507 (3)	C314—H314	0.95
C5—H5A	0.98	C315—C316	1.379 (3)
C5—H5B	0.98	C315—H315	0.95
C5—H5C	0.98	C316—H316	0.95
C11A—C12A	1.39	C321—C326	1.388 (3)
C11A—C16A	1.39	C321—C322	1.390 (3)
C12A—C13A	1.39	C322—C323	1.391 (4)
C12A—C17A	1.515 (5)	C322—H322	0.95
C13A—C14A	1.39	C323—C324	1.381 (4)
C13A—C18A	1.516 (4)	C323—H323	0.95
C14A—C15A	1.39	C324—C325	1.380 (4)
C14A—H14A	0.95	C324—H324	0.95
C15A—C16A	1.39	C325—C326	1.390 (3)
C15A—H15A	0.95	C325—H325	0.95

C16A—H16A	0.95	C326—H326	0.95
C18A—H18A	0.98	C331—C332	1.384 (4)
C18A—H18B	0.98	C331—C336	1.388 (3)
C18A—H18C	0.98	C332—C333	1.403 (4)
C17A—H17A	0.98	C332—H332	0.95
C17A—H17B	0.98	C333—C334	1.369 (5)
C17A—H17C	0.98	C333—H333	0.95
C11B—C12B	1.39	C334—C335	1.366 (4)
C11B—C16B	1.39	C334—H334	0.95
C12B—C13B	1.39	C335—C336	1.387 (3)
C12B—C17B	1.506 (7)	C335—H335	0.95
C13B—C14B	1.39	C336—H336	0.95
C14—Rh1—O12	176.48 (9)	C14B—C15B—H15B	120
C14—Rh1—N11	93.62 (9)	C15B—C16B—C11B	120
O12—Rh1—N11	89.53 (7)	C15B—C16B—H16B	120
C14—Rh1—P13	91.86 (7)	C11B—C16B—H16B	120
O12—Rh1—P13	84.97 (5)	C12B—C17B—H17D	109.5
N11—Rh1—P13	174.49 (6)	C12B—C17B—H17E	109.5
C2—N11—C11A	114.9 (2)	H17D—C17B—H17E	109.5
C2—N11—C11B	117.8 (2)	C12B—C17B—H17F	109.5
C2—N11—Rh1	125.77 (17)	H17D—C17B—H17F	109.5
C11A—N11—Rh1	117.14 (17)	H17E—C17B—H17F	109.5
C11B—N11—Rh1	113.14 (19)	C13B—C18B—H18D	109.5
C4—O12—Rh1	126.75 (15)	C13B—C18B—H18E	109.5
C311—P13—C331	103.07 (10)	H18D—C18B—H18E	109.5
C311—P13—C321	104.96 (11)	C13B—C18B—H18F	109.5
C331—P13—C321	103.48 (11)	H18D—C18B—H18F	109.5
C311—P13—Rh1	118.23 (8)	H18E—C18B—H18F	109.5
C331—P13—Rh1	112.44 (8)	O14—C14—Rh1	178.1 (2)
C321—P13—Rh1	113.14 (8)	C312—C311—C316	118.8 (2)
C2—C1—H1A	109.5	C312—C311—P13	122.29 (17)
C2—C1—H1B	109.5	C316—C311—P13	118.70 (17)
H1A—C1—H1B	109.5	C313—C312—C311	120.4 (2)
C2—C1—H1C	109.5	C313—C312—H312	119.8
H1A—C1—H1C	109.5	C311—C312—H312	119.8
H1B—C1—H1C	109.5	C314—C313—C312	120.0 (2)
N11—C2—C3	123.8 (2)	C314—C313—H313	120
N11—C2—C1	120.3 (2)	C312—C313—H313	120
C3—C2—C1	115.8 (2)	C315—C314—C313	120.1 (2)
C4—C3—C2	127.4 (2)	C315—C314—H314	120
C4—C3—H3	116.3	C313—C314—H314	120
C2—C3—H3	116.3	C314—C315—C316	120.5 (2)
O12—C4—C3	126.1 (2)	C314—C315—H315	119.7
O12—C4—C5	113.6 (2)	C316—C315—H315	119.7
C3—C4—C5	120.3 (2)	C315—C316—C311	120.2 (2)
C4—C5—H5A	109.5	C315—C316—H316	119.9
C4—C5—H5B	109.5	C311—C316—H316	119.9
H5A—C5—H5B	109.5	C326—C321—C322	119.0 (2)
C4—C5—H5C	109.5	C326—C321—P13	117.40 (18)

supplementary materials

H5A—C5—H5C	109.5	C322—C321—P13	123.59 (18)
H5B—C5—H5C	109.5	C321—C322—C323	120.0 (2)
C12A—C11A—C16A	120	C321—C322—H322	120
C12A—C11A—N11	114.94 (19)	C323—C322—H322	120
C16A—C11A—N11	125.06 (19)	C324—C323—C322	120.8 (2)
C11A—C12A—C13A	120	C324—C323—H323	119.6
C11A—C12A—C17A	118.8 (2)	C322—C323—H323	119.6
C13A—C12A—C17A	121.2 (2)	C325—C324—C323	119.3 (2)
C12A—C13A—C14A	120	C325—C324—H324	120.3
C12A—C13A—C18A	121.9 (2)	C323—C324—H324	120.3
C14A—C13A—C18A	118.0 (2)	C324—C325—C326	120.3 (2)
C15A—C14A—C13A	120	C324—C325—H325	119.9
C15A—C14A—H14A	120	C326—C325—H325	119.9
C13A—C14A—H14A	120	C321—C326—C325	120.6 (2)
C14A—C15A—C16A	120	C321—C326—H326	119.7
C14A—C15A—H15A	120	C325—C326—H326	119.7
C16A—C15A—H15A	120	C332—C331—C336	119.0 (2)
C15A—C16A—C11A	120	C332—C331—P13	122.3 (2)
C15A—C16A—H16A	120	C336—C331—P13	118.61 (18)
C11A—C16A—H16A	120	C331—C332—C333	119.8 (3)
C12B—C11B—C16B	120	C331—C332—H332	120.1
C12B—C11B—N11	112.0 (2)	C333—C332—H332	120.1
C16B—C11B—N11	128.0 (2)	C334—C333—C332	120.2 (3)
C11B—C12B—C13B	120	C334—C333—H333	119.9
C11B—C12B—C17B	119.6 (3)	C332—C333—H333	119.9
C13B—C12B—C17B	120.4 (3)	C335—C334—C333	120.3 (3)
C14B—C13B—C12B	120	C335—C334—H334	119.9
C14B—C13B—C18B	118.6 (3)	C333—C334—H334	119.9
C12B—C13B—C18B	121.4 (3)	C334—C335—C336	120.2 (3)
C13B—C14B—C15B	120	C334—C335—H335	119.9
C13B—C14B—H14B	120	C336—C335—H335	119.9
C15B—C14B—H14B	120	C335—C336—C331	120.5 (3)
C16B—C15B—C14B	120	C335—C336—H336	119.7
C16B—C15B—H15B	120	C331—C336—H336	119.7
C14—Rh1—N11—C2	174.7 (3)	C16B—C11B—C12B—C17B	-178.1 (4)
O12—Rh1—N11—C2	-3.8 (3)	N11—C11B—C12B—C17B	-0.6 (4)
C14—Rh1—N11—C11A	-23.0 (2)	C11B—C12B—C13B—C14B	0
O12—Rh1—N11—C11A	158.5 (2)	C17B—C12B—C13B—C14B	178.1 (4)
C14—Rh1—N11—C11B	15.6 (2)	C11B—C12B—C13B—C18B	-178.8 (4)
O12—Rh1—N11—C11B	-162.8 (2)	C17B—C12B—C13B—C18B	-0.7 (5)
N11—Rh1—O12—C4	8.13 (19)	C12B—C13B—C14B—C15B	0
P13—Rh1—O12—C4	-171.45 (18)	C18B—C13B—C14B—C15B	178.8 (4)
C14—Rh1—P13—C311	16.41 (11)	C13B—C14B—C15B—C16B	0
O12—Rh1—P13—C311	-165.13 (10)	C14B—C15B—C16B—C11B	0
C14—Rh1—P13—C331	136.38 (11)	C12B—C11B—C16B—C15B	0
O12—Rh1—P13—C331	-45.16 (9)	N11—C11B—C16B—C15B	-177.0 (4)
C14—Rh1—P13—C321	-106.82 (11)	C331—P13—C311—C312	-16.6 (2)
O12—Rh1—P13—C321	71.63 (10)	C321—P13—C311—C312	-124.6 (2)
C11A—N11—C2—C3	-164.1 (3)	Rh1—P13—C311—C312	108.17 (19)

C11B—N11—C2—C3	156.7 (3)	C331—P13—C311—C316	168.3 (2)
Rh1—N11—C2—C3	-1.5 (4)	C321—P13—C311—C316	60.2 (2)
C11A—N11—C2—C1	16.8 (4)	Rh1—P13—C311—C316	-67.0 (2)
C11B—N11—C2—C1	-22.4 (4)	C316—C311—C312—C313	0.9 (4)
Rh1—N11—C2—C1	179.4 (2)	P13—C311—C312—C313	-174.3 (2)
N11—C2—C3—C4	5.4 (5)	C311—C312—C313—C314	-0.9 (4)
C1—C2—C3—C4	-175.5 (3)	C312—C313—C314—C315	0.4 (4)
Rh1—O12—C4—C3	-7.5 (3)	C313—C314—C315—C316	0.1 (5)
Rh1—O12—C4—C5	171.45 (15)	C314—C315—C316—C311	-0.2 (4)
C2—C3—C4—O12	-0.5 (4)	C312—C311—C316—C315	-0.3 (4)
C2—C3—C4—C5	-179.4 (3)	P13—C311—C316—C315	175.1 (2)
C2—N11—C11A—C12A	-98.8 (3)	C311—P13—C321—C326	-166.46 (19)
C11B—N11—C11A—C12A	4.8 (3)	C331—P13—C321—C326	85.8 (2)
Rh1—N11—C11A—C12A	97.0 (2)	Rh1—P13—C321—C326	-36.2 (2)
C2—N11—C11A—C16A	81.7 (3)	C311—P13—C321—C322	15.4 (2)
C11B—N11—C11A—C16A	-174.7 (4)	C331—P13—C321—C322	-92.3 (2)
Rh1—N11—C11A—C16A	-82.5 (2)	Rh1—P13—C321—C322	145.7 (2)
C16A—C11A—C12A—C13A	0	C326—C321—C322—C323	0.6 (4)
N11—C11A—C12A—C13A	-179.5 (3)	P13—C321—C322—C323	178.7 (2)
C16A—C11A—C12A—C17A	178.9 (3)	C321—C322—C323—C324	-1.0 (4)
N11—C11A—C12A—C17A	-0.6 (3)	C322—C323—C324—C325	0.6 (5)
C11A—C12A—C13A—C14A	0	C323—C324—C325—C326	0.2 (5)
C17A—C12A—C13A—C14A	-178.9 (3)	C322—C321—C326—C325	0.1 (4)
C11A—C12A—C13A—C18A	-176.5 (3)	P13—C321—C326—C325	-178.1 (2)
C17A—C12A—C13A—C18A	4.6 (4)	C324—C325—C326—C321	-0.5 (4)
C12A—C13A—C14A—C15A	0	C311—P13—C331—C332	-83.3 (2)
C18A—C13A—C14A—C15A	176.7 (3)	C321—P13—C331—C332	25.9 (2)
C13A—C14A—C15A—C16A	0	Rh1—P13—C331—C332	148.32 (19)
C14A—C15A—C16A—C11A	0	C311—P13—C331—C336	93.63 (19)
C12A—C11A—C16A—C15A	0	C321—P13—C331—C336	-157.22 (18)
N11—C11A—C16A—C15A	179.5 (3)	Rh1—P13—C331—C336	-34.78 (19)
C2—N11—C11B—C12B	98.0 (3)	C336—C331—C332—C333	-0.7 (4)
C11A—N11—C11B—C12B	3.6 (3)	P13—C331—C332—C333	176.2 (2)
Rh1—N11—C11B—C12B	-101.2 (2)	C331—C332—C333—C334	-0.4 (4)
C2—N11—C11B—C16B	-84.8 (3)	C332—C333—C334—C335	0.9 (4)
C11A—N11—C11B—C16B	-179.2 (5)	C333—C334—C335—C336	-0.3 (4)
Rh1—N11—C11B—C16B	76.1 (3)	C334—C335—C336—C331	-0.8 (3)
C16B—C11B—C12B—C13B	0	C332—C331—C336—C335	1.3 (3)
N11—C11B—C12B—C13B	177.5 (3)	P13—C331—C336—C335	-175.74 (17)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C326—H326 \cdots O12	0.95	2.36	3.177 (3)	143

Table 2

Comparative geometrical parameters for similar $[Rh(N,O-bid)(CO)(PPh_3)]$ complexes (\AA , $^\circ$).

Parameters	(I)	(II)	(III)
------------	-----	------	-------

supplementary materials

Rh1—N11	2.069 (2)	2.045 (4)	2.045 (3)
Rh1—O12	2.028 (2)	2.044 (3)	2.045 (2)
Rh1—P13	2.2635 (6)	2.275 (1)	2.281 (2)
Rh1—C14	1.807 (2)	1.784 (5)	1.804 (3)
C14—O14	1.152 (3)	1.142 (7)	1.148 (4)
N11···O12	2.885 (3)	2.826 (6)	2.841 (3)
N11—Rh1—O12	89.54 (8)	87.4 (1)	87.95 (8)
O12—Rh1—P13	84.97 (5)	89.7 (1)	89.91 (5)
P13—Rh1—C14	91.87 (7)	90.3 (2)	89.48 (9)
N11—Rh1—C14	93.6 (1)	92.6 (2)	92.6 (1)
N11—C2—C4—O12	4.1 (2)	1.2 (4)	1.5 (2)
θ_E (Tolman, 1977)	156.39 (3)	156.0 (2)	156.23 (4)

(I) This work. (II) N,O-bid = 4-aminopent-3-en-2-onato (Damoense *et al.*, 1994). (III) N,O-bid = 4-amino-1,1,1-trifluoropent-3-en-2-onato (Varshavsky *et al.*, 2001).

Fig. 1




Fig. 2

