

Bis(5-amino-2-chlorobenzoato- κ O)-triphenylantimony(V)

Liyuan Wen, Handong Yin* and Chuanhua Wang

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: handongyin@163.com


Received 10 October 2009; accepted 22 October 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.038; wR factor = 0.087; data-to-parameter ratio = 14.0.

In the title compound, $[\text{Sb}(\text{C}_6\text{H}_5)_3(\text{C}_7\text{H}_5\text{ClNO}_2)_2]$, the Sb center has a distorted trigonal-bipyramidal geometry, with the O atoms of two carboxylate groups in axial positions and the C atoms of the phenyl groups in equatorial positions. Intra-molecular C—H \cdots O interactions occur. The molecules are connected by intermolecular N—H \cdots O, N—H \cdots N and C—H \cdots O hydrogen-bonding interactions and C—H \cdots π stacking interactions, forming a three-dimensional supramolecular framework

Related literature

For related structures, see: Yin *et al.* (2009); Wang *et al.* (2005).



Experimental

Crystal data

$[\text{Sb}(\text{C}_6\text{H}_5)_3(\text{C}_7\text{H}_5\text{ClNO}_2)_2]$

$M_r = 694.19$

Monoclinic, $C2/c$

$a = 30.683$ (3) Å

$b = 9.0128$ (12) Å

$c = 23.096$ (3) Å

$\beta = 106.161$ (2) $^\circ$

$V = 6134.6$ (12) Å³

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 1.11$ mm⁻¹

$T = 298$ K

$0.35 \times 0.33 \times 0.17$ mm

Data collection

Siemens SMART CCD area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.697$, $T_{\max} = 0.833$

15481 measured reflections

5409 independent reflections

3450 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.087$

$S = 1.04$

5409 reflections

386 parameters

4 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.86$ e Å⁻³

$\Delta\rho_{\min} = -0.49$ e Å⁻³

Table 1

Selected geometric parameters (Å, $^\circ$).

Sb1—C21	2.098 (5)	Sb1—O3	2.125 (3)
Sb1—C15	2.108 (5)	Sb1—O1	2.137 (3)
Sb1—C27	2.109 (5)		
C21—Sb1—C15	109.59 (19)	C27—Sb1—O3	89.15 (15)
C21—Sb1—C27	109.92 (19)	C21—Sb1—O1	87.45 (16)
C15—Sb1—C27	140.49 (19)	C15—Sb1—O1	87.74 (15)
C21—Sb1—O3	87.33 (16)	C27—Sb1—O1	94.27 (15)
C15—Sb1—O3	92.36 (15)	O3—Sb1—O1	174.49 (12)

Table 2

Hydrogen-bond geometry (Å, $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C32—H32 \cdots O2	0.93	2.39	3.105 (6)	134
C20—H20 \cdots O4	0.93	2.47	3.120 (6)	127
N1—H1B \cdots N2 ⁱ	0.865 (19)	2.43 (3)	3.211 (8)	150 (5)
N1—H1A \cdots O2 ⁱⁱ	0.86 (2)	2.43 (5)	3.114 (6)	137 (5)
N2—H2A \cdots N1 ⁱⁱⁱ	0.847 (19)	2.44 (2)	3.272 (7)	169 (5)
N2—H2B \cdots O4 ^{iv}	0.847 (19)	2.36 (3)	3.151 (6)	156 (5)
C14—H14 \cdots O4 ^v	0.93	2.58	3.369 (6)	143
C12—H12 \cdots O2 ^v	0.93	2.56	3.433 (6)	157
C23—H23 \cdots Cg1 ^{vi}	0.93	2.72	3.567 (7)	151

Symmetry codes: (i) $x, -y + 2, z + \frac{1}{2}$; (ii) $-x, -y + 2, -z + 1$; (iii) $-x, y - 1, -z + \frac{1}{2}$; (iv) $-x, y, -z + \frac{1}{2}$; (v) $x, -y + 1, z - \frac{1}{2}$; (vi) $-x + \frac{1}{2}, -y + \frac{3}{2}, -z + 1$. Cg1 is the centroid of the C27—C32 ring.

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

We acknowledge the National Natural Science Foundation of China (20771053) and the Natural Science Foundation of Shandong Province (Y2008B48) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2233).

References

- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Wang, G. C., Lu, Y. N., Xiao, J., Yu, L., Song, H. B., Li, J. S., Cui, J. R., Wang, R. Q. & Ran, F. X. (2005). *J. Organomet. Chem.* **690**, 151–156.
- Yin, H. D., Wen, L. Y., Cui, J. C. & Li, W. K. (2009). *Polyhedron*, **28**, 2919–2926.

supplementary materials

Acta Cryst. (2009). E65, m1443 [doi:10.1107/S1600536809043578]

Bis(5-amino-2-chlorobenzoato- κ O)triphenylantimony(V)

L. Wen, H. Yin and C. Wang

Comment

Recently, the chemistry of antimony complexes derived from carboxylates has become an active area of research due to the biological perspective and their versatile bonding modes, the striking structural possibilities ranging from discrete monomeric structures to supramolecular assemblies (Yin *et al.* 2009). As a part of our ongoing investigations in this field we have synthesized the title compound and determined its crystal structure. As is shown in Fig. 1, the central antimony atom is five-coordinated with a slightly distorted trigonal bipyramidal geometry. Around the central Sb1 atom, C15, C21 and C27 occupy the equatorial plane, while O1 and O3 lie in axial sites. The Sb—O bond distances (Sb1—O1 = 2.137 (3) Å; Sb1—O3 = 2.125 (3) Å) (Table 1) are comparable to those found in organoantimony arylhydroximates (Wang *et al.* 2005). The Sb—C bond distances (Sb1—C15 = 2.108 (5) Å; Sb1—C21 = 2.098 (5) Å; Sb1—C27 = 2.109 (5) Å) of the compound lie within the normal range for Sb—C (phenyl) bonds (2.10–2.13 Å). The supramolecular structure of the title compound results from intermolecular C—H \cdots O, N—H \cdots O and N—H \cdots N hydrogen-bonding interactions and C—H \cdots π stacking interactions (Fig. 2, Table 2) assembling the molecules into a three-dimensional supramolecular frameworks

Experimental

The reaction was carried out under nitrogen atmosphere. 5-Amino-2-chlorobenzoic acid (1 mmol) and sodium ethoxide (1.2 mmol) were added to a stirred solution of methanol (30 ml) in a Schlenk flask and stirred for 0.5 h. Triphenylantimony dichloride (0.5 mmol) was then added to the reactor and the reaction mixture was stirred for 12 h at room temperature. The resulting clear solution was evaporated under vacuum. The product was crystallized from a mixture of ether/n-hexane (1:1) to yield colourless blocks of the title compound (yield 80%). Anal. Calcd (%) for C₃₂H₂₅Cl₂N₂O₄Sb (Mr = 694.19): C, 55.37; H, 3.63; Cl, 10.21; N, 4.04. Found (%): C, 55.31; H, 3.76; Cl, 10.31; N, 4.15.

Refinement

The N-bound H atoms were located in a difference Fourier map. In the refinement process the N-H bond lengths were restrained to 0.85 (2) Å and isotropic displacement parameters of these H atoms were freely refined. Other H atoms were positioned geometrically, with C—H = 0.93 Å and refined with a riding model; $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the compound, showing 50% probability displacement ellipsoids. H atoms have been omitted for clarity.




Fig. 2. View of the three-dimensional supramolecular network structure in the title compound.

Bis(5-amino-2-chlorobenzoato- κO)triphenylantimony(V)

Crystal data

[Sb(C₆H₅)₃(C₇H₅ClNO₂)₂]

$M_r = 694.19$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 30.683\ (3)\ \text{\AA}$

$b = 9.0128\ (12)\ \text{\AA}$

$c = 23.096\ (3)\ \text{\AA}$

$\beta = 106.161\ (2)^\circ$

$V = 6134.6\ (12)\ \text{\AA}^3$

$Z = 8$

$F_{000} = 2784$

$D_x = 1.503\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3588 reflections

$\theta = 2.4\text{--}23.7^\circ$

$\mu = 1.11\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, colourless

$0.35 \times 0.33 \times 0.17\ \text{mm}$

Data collection

Siemens SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 298\ \text{K}$

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.697$, $T_{\max} = 0.833$

15481 measured reflections

5409 independent reflections

3450 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 25.0^\circ$

$\theta_{\min} = 1.4^\circ$

$h = -25 \rightarrow 36$

$k = -10 \rightarrow 10$

$l = -27 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.087$

$S = 1.04$

5409 reflections

386 parameters

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0252P)^2 + 12.9995P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.86\ \text{e \AA}^{-3}$

$\Delta\rho_{\min} = -0.49\ \text{e \AA}^{-3}$

4 restraints

Extinction correction: none

Primary atom site location: structure-invariant direct methods

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sb1	0.133450 (10)	0.75719 (4)	0.387976 (14)	0.03529 (11)
Cl1	0.19341 (5)	0.92982 (17)	0.61118 (7)	0.0683 (4)
Cl2	0.17565 (5)	0.4095 (2)	0.25676 (8)	0.0998 (7)
N1	0.02031 (18)	1.2629 (6)	0.5343 (2)	0.0590 (12)
N2	-0.00974 (17)	0.4432 (6)	0.0918 (2)	0.0567 (13)
O1	0.14690 (11)	0.8704 (3)	0.47239 (14)	0.0427 (8)
O2	0.08342 (11)	0.7652 (4)	0.48081 (14)	0.0478 (8)
O3	0.12400 (10)	0.6602 (3)	0.30150 (13)	0.0411 (8)
O4	0.06817 (10)	0.5306 (4)	0.32076 (15)	0.0445 (9)
C1	0.11308 (17)	0.8607 (5)	0.4962 (2)	0.0390 (12)
C2	0.10837 (16)	0.9804 (5)	0.5384 (2)	0.0382 (12)
C3	0.14220 (16)	1.0253 (5)	0.5887 (2)	0.0432 (13)
C4	0.13472 (19)	1.1428 (6)	0.6237 (2)	0.0531 (14)
H4	0.1570	1.1696	0.6585	0.064*
C5	0.09432 (19)	1.2197 (6)	0.6068 (2)	0.0571 (15)
H5	0.0898	1.2997	0.6299	0.069*
C6	0.06032 (18)	1.1792 (6)	0.5558 (2)	0.0469 (13)
C7	0.06747 (17)	1.0588 (5)	0.5228 (2)	0.0426 (12)
H7	0.0445	1.0290	0.4893	0.051*
C8	0.09252 (16)	0.5591 (5)	0.2885 (2)	0.0360 (11)
C9	0.08637 (16)	0.4847 (5)	0.2287 (2)	0.0367 (11)
C10	0.12097 (16)	0.4236 (6)	0.2091 (2)	0.0524 (14)
C11	0.11202 (19)	0.3647 (6)	0.1518 (3)	0.0648 (17)
H11	0.1353	0.3213	0.1392	0.078*
C12	0.06878 (19)	0.3698 (6)	0.1131 (2)	0.0560 (15)
H12	0.0635	0.3331	0.0741	0.067*
C13	0.03327 (16)	0.4285 (5)	0.1315 (2)	0.0388 (12)
C14	0.04275 (16)	0.4845 (5)	0.1899 (2)	0.0391 (12)
H14	0.0191	0.5231	0.2032	0.047*
C15	0.07540 (17)	0.8918 (5)	0.3575 (2)	0.0408 (12)
C16	0.0831 (2)	1.0412 (6)	0.3515 (3)	0.0618 (16)
H16	0.1126	1.0768	0.3595	0.074*
C17	0.0468 (3)	1.1380 (7)	0.3337 (3)	0.084 (2)
H17	0.0518	1.2384	0.3286	0.100*
C18	0.0039 (3)	1.0854 (8)	0.3236 (3)	0.080 (2)
H18	-0.0203	1.1514	0.3129	0.095*
C19	-0.00454 (19)	0.9387 (7)	0.3289 (3)	0.0665 (17)
H19	-0.0342	0.9047	0.3212	0.080*
C20	0.03162 (18)	0.8400 (6)	0.3460 (2)	0.0526 (14)
H20	0.0262	0.7394	0.3497	0.063*
C21	0.18995 (16)	0.8670 (5)	0.3742 (2)	0.0413 (12)
C22	0.22768 (18)	0.8915 (6)	0.4213 (3)	0.0553 (15)

supplementary materials

H22	0.2283	0.8595	0.4598	0.066*
C23	0.26495 (19)	0.9639 (7)	0.4118 (3)	0.0706 (18)
H23	0.2908	0.9784	0.4437	0.085*
C24	0.2638 (2)	1.0132 (8)	0.3563 (4)	0.085 (2)
H24	0.2887	1.0628	0.3503	0.102*
C25	0.2264 (2)	0.9912 (8)	0.3089 (3)	0.098 (3)
H25	0.2259	1.0255	0.2708	0.117*
C26	0.18914 (19)	0.9175 (7)	0.3177 (3)	0.0725 (19)
H26	0.1636	0.9022	0.2854	0.087*
C27	0.15298 (16)	0.5477 (5)	0.4274 (2)	0.0375 (12)
C28	0.18933 (18)	0.4839 (6)	0.4126 (2)	0.0545 (15)
H28	0.2048	0.5363	0.3898	0.065*
C29	0.2027 (2)	0.3410 (7)	0.4322 (3)	0.0680 (17)
H29	0.2274	0.2980	0.4228	0.082*
C30	0.1797 (2)	0.2636 (6)	0.4651 (3)	0.0632 (16)
H30	0.1886	0.1676	0.4779	0.076*
C31	0.14364 (19)	0.3273 (6)	0.4793 (3)	0.0584 (16)
H31	0.1280	0.2738	0.5015	0.070*
C32	0.12994 (16)	0.4711 (5)	0.4609 (2)	0.0447 (13)
H32	0.1056	0.5144	0.4713	0.054*
H1A	-0.0042 (13)	1.213 (6)	0.521 (3)	0.09 (3)*
H2A	-0.0141 (17)	0.386 (5)	0.0615 (16)	0.061 (19)*
H1B	0.0182 (19)	1.324 (5)	0.5623 (19)	0.08 (2)*
H2B	-0.0315 (12)	0.450 (6)	0.108 (2)	0.057 (19)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sb1	0.03935 (18)	0.03370 (18)	0.03511 (18)	-0.00248 (17)	0.01417 (13)	-0.00089 (18)
Cl1	0.0535 (9)	0.0761 (11)	0.0696 (11)	0.0103 (8)	0.0079 (7)	-0.0040 (8)
Cl2	0.0513 (10)	0.1624 (19)	0.0777 (13)	0.0366 (11)	0.0047 (8)	-0.0392 (12)
N1	0.059 (3)	0.057 (3)	0.064 (3)	0.010 (3)	0.023 (3)	0.000 (3)
N2	0.049 (3)	0.077 (4)	0.043 (3)	-0.002 (3)	0.010 (3)	-0.014 (3)
O1	0.047 (2)	0.047 (2)	0.038 (2)	0.0001 (16)	0.0193 (17)	-0.0075 (16)
O2	0.058 (2)	0.039 (2)	0.050 (2)	-0.0073 (18)	0.0201 (17)	-0.0049 (18)
O3	0.049 (2)	0.0379 (19)	0.038 (2)	-0.0100 (16)	0.0161 (16)	-0.0050 (16)
O4	0.042 (2)	0.055 (2)	0.040 (2)	-0.0035 (17)	0.0166 (17)	-0.0031 (17)
C1	0.050 (3)	0.031 (3)	0.038 (3)	0.003 (2)	0.014 (3)	0.001 (2)
C2	0.049 (3)	0.035 (3)	0.036 (3)	-0.005 (2)	0.021 (2)	0.001 (2)
C3	0.047 (3)	0.045 (3)	0.039 (3)	0.002 (2)	0.014 (3)	0.003 (2)
C4	0.063 (4)	0.055 (4)	0.041 (3)	-0.003 (3)	0.014 (3)	-0.007 (3)
C5	0.066 (4)	0.061 (4)	0.050 (4)	0.000 (3)	0.025 (3)	-0.013 (3)
C6	0.050 (3)	0.042 (3)	0.054 (4)	0.000 (3)	0.023 (3)	0.004 (3)
C7	0.054 (3)	0.040 (3)	0.034 (3)	-0.005 (2)	0.012 (2)	-0.001 (2)
C8	0.038 (3)	0.033 (3)	0.038 (3)	0.004 (2)	0.012 (2)	0.003 (2)
C9	0.039 (3)	0.036 (3)	0.039 (3)	0.000 (2)	0.018 (2)	0.001 (2)
C10	0.041 (3)	0.066 (4)	0.052 (4)	0.009 (3)	0.014 (3)	-0.012 (3)
C11	0.060 (4)	0.077 (4)	0.064 (4)	0.015 (3)	0.027 (3)	-0.027 (3)

C12	0.064 (4)	0.059 (4)	0.045 (4)	-0.003 (3)	0.016 (3)	-0.021 (3)
C13	0.046 (3)	0.037 (3)	0.037 (3)	-0.005 (2)	0.017 (3)	-0.006 (2)
C14	0.041 (3)	0.036 (3)	0.045 (3)	-0.002 (2)	0.019 (2)	-0.001 (2)
C15	0.050 (3)	0.040 (3)	0.034 (3)	0.002 (2)	0.014 (2)	0.000 (2)
C16	0.072 (4)	0.038 (3)	0.073 (5)	0.002 (3)	0.018 (3)	0.006 (3)
C17	0.118 (6)	0.037 (4)	0.092 (6)	0.025 (4)	0.024 (5)	0.005 (3)
C18	0.092 (5)	0.072 (5)	0.074 (5)	0.044 (4)	0.023 (4)	0.006 (4)
C19	0.056 (4)	0.070 (4)	0.077 (5)	0.019 (3)	0.023 (3)	0.008 (4)
C20	0.052 (4)	0.046 (3)	0.061 (4)	0.011 (3)	0.018 (3)	0.005 (3)
C21	0.045 (3)	0.037 (3)	0.047 (3)	-0.006 (2)	0.021 (3)	-0.006 (2)
C22	0.052 (4)	0.061 (4)	0.053 (4)	-0.010 (3)	0.013 (3)	-0.004 (3)
C23	0.045 (4)	0.085 (5)	0.078 (5)	-0.019 (3)	0.010 (3)	-0.020 (4)
C24	0.073 (5)	0.090 (5)	0.105 (6)	-0.037 (4)	0.044 (5)	-0.015 (5)
C25	0.099 (6)	0.126 (7)	0.080 (6)	-0.046 (5)	0.043 (5)	0.012 (5)
C26	0.063 (4)	0.101 (5)	0.057 (4)	-0.040 (4)	0.022 (3)	0.006 (4)
C27	0.042 (3)	0.033 (3)	0.034 (3)	0.002 (2)	0.004 (2)	0.001 (2)
C28	0.060 (4)	0.050 (3)	0.060 (4)	0.012 (3)	0.027 (3)	0.005 (3)
C29	0.066 (4)	0.064 (4)	0.073 (5)	0.027 (3)	0.018 (3)	0.003 (4)
C30	0.076 (4)	0.037 (3)	0.068 (4)	0.007 (3)	0.005 (3)	0.008 (3)
C31	0.057 (4)	0.044 (3)	0.070 (4)	-0.009 (3)	0.010 (3)	0.011 (3)
C32	0.044 (3)	0.041 (3)	0.049 (3)	-0.003 (2)	0.011 (3)	0.006 (3)

Geometric parameters (Å, °)

Sb1—C21	2.098 (5)	C13—C14	1.392 (6)
Sb1—C15	2.108 (5)	C14—H14	0.9300
Sb1—C27	2.109 (5)	C15—C20	1.376 (6)
Sb1—O3	2.125 (3)	C15—C16	1.381 (6)
Sb1—O1	2.137 (3)	C16—C17	1.383 (7)
C11—C3	1.739 (5)	C16—H16	0.9300
C12—C10	1.736 (5)	C17—C18	1.357 (8)
N1—C6	1.408 (7)	C17—H17	0.9300
N1—H1A	0.86 (2)	C18—C19	1.360 (8)
N1—H1B	0.865 (19)	C18—H18	0.9300
N2—C13	1.388 (6)	C19—C20	1.391 (7)
N2—H2A	0.847 (19)	C19—H19	0.9300
N2—H2B	0.847 (19)	C20—H20	0.9300
O1—C1	1.305 (5)	C21—C22	1.369 (6)
O2—C1	1.231 (5)	C21—C26	1.375 (7)
O3—C8	1.301 (5)	C22—C23	1.386 (7)
O4—C8	1.220 (5)	C22—H22	0.9300
C1—C2	1.488 (6)	C23—C24	1.348 (8)
C2—C3	1.385 (6)	C23—H23	0.9300
C2—C7	1.397 (6)	C24—C25	1.361 (9)
C3—C4	1.390 (7)	C24—H24	0.9300
C4—C5	1.378 (7)	C25—C26	1.385 (7)
C4—H4	0.9300	C25—H25	0.9300
C5—C6	1.387 (7)	C26—H26	0.9300
C5—H5	0.9300	C27—C32	1.372 (6)

supplementary materials

C6—C7	1.378 (6)	C27—C28	1.379 (6)
C7—H7	0.9300	C28—C29	1.388 (7)
C8—C9	1.499 (6)	C28—H28	0.9300
C9—C10	1.379 (6)	C29—C30	1.364 (8)
C9—C14	1.388 (6)	C29—H29	0.9300
C10—C11	1.381 (7)	C30—C31	1.366 (7)
C11—C12	1.378 (7)	C30—H30	0.9300
C11—H11	0.9300	C31—C32	1.391 (7)
C12—C13	1.380 (6)	C31—H31	0.9300
C12—H12	0.9300	C32—H32	0.9300
C21—Sb1—C15	109.59 (19)	C9—C14—C13	122.0 (5)
C21—Sb1—C27	109.92 (19)	C9—C14—H14	119.0
C15—Sb1—C27	140.49 (19)	C13—C14—H14	119.0
C21—Sb1—O3	87.33 (16)	C20—C15—C16	119.7 (5)
C15—Sb1—O3	92.36 (15)	C20—C15—Sb1	123.9 (4)
C27—Sb1—O3	89.15 (15)	C16—C15—Sb1	116.3 (4)
C21—Sb1—O1	87.45 (16)	C15—C16—C17	120.0 (6)
C15—Sb1—O1	87.74 (15)	C15—C16—H16	120.0
C27—Sb1—O1	94.27 (15)	C17—C16—H16	120.0
O3—Sb1—O1	174.49 (12)	C18—C17—C16	119.5 (6)
C6—N1—H1A	116 (4)	C18—C17—H17	120.3
C6—N1—H1B	108 (4)	C16—C17—H17	120.3
H1A—N1—H1B	112 (6)	C17—C18—C19	121.6 (6)
C13—N2—H2A	113 (4)	C17—C18—H18	119.2
C13—N2—H2B	116 (4)	C19—C18—H18	119.2
H2A—N2—H2B	116 (5)	C18—C19—C20	119.4 (6)
C1—O1—Sb1	112.2 (3)	C18—C19—H19	120.3
C8—O3—Sb1	113.8 (3)	C20—C19—H19	120.3
O2—C1—O1	122.4 (4)	C15—C20—C19	119.8 (5)
O2—C1—C2	119.8 (5)	C15—C20—H20	120.1
O1—C1—C2	117.5 (4)	C19—C20—H20	120.1
C3—C2—C7	118.5 (4)	C22—C21—C26	119.3 (5)
C3—C2—C1	125.1 (4)	C22—C21—Sb1	120.5 (4)
C7—C2—C1	116.2 (4)	C26—C21—Sb1	120.2 (4)
C2—C3—C4	120.3 (5)	C21—C22—C23	120.1 (5)
C2—C3—C11	120.6 (4)	C21—C22—H22	119.9
C4—C3—C11	119.0 (4)	C23—C22—H22	119.9
C5—C4—C3	119.9 (5)	C24—C23—C22	120.1 (6)
C5—C4—H4	120.0	C24—C23—H23	120.0
C3—C4—H4	120.0	C22—C23—H23	120.0
C4—C5—C6	120.9 (5)	C23—C24—C25	120.6 (6)
C4—C5—H5	119.6	C23—C24—H24	119.7
C6—C5—H5	119.6	C25—C24—H24	119.7
C7—C6—C5	118.6 (5)	C24—C25—C26	119.8 (7)
C7—C6—N1	118.9 (5)	C24—C25—H25	120.1
C5—C6—N1	122.3 (5)	C26—C25—H25	120.1
C6—C7—C2	121.7 (5)	C21—C26—C25	120.0 (6)
C6—C7—H7	119.1	C21—C26—H26	120.0
C2—C7—H7	119.1	C25—C26—H26	120.0

O4—C8—O3	123.2 (5)	C32—C27—C28	120.6 (5)
O4—C8—C9	121.8 (4)	C32—C27—Sb1	124.4 (4)
O3—C8—C9	114.9 (4)	C28—C27—Sb1	114.8 (4)
C10—C9—C14	118.6 (5)	C27—C28—C29	119.5 (5)
C10—C9—C8	124.7 (4)	C27—C28—H28	120.2
C14—C9—C8	116.6 (4)	C29—C28—H28	120.2
C9—C10—C11	120.1 (5)	C30—C29—C28	120.3 (6)
C9—C10—C12	121.1 (4)	C30—C29—H29	119.9
C11—C10—C12	118.6 (4)	C28—C29—H29	119.9
C12—C11—C10	120.5 (5)	C29—C30—C31	119.8 (5)
C12—C11—H11	119.8	C29—C30—H30	120.1
C10—C11—H11	119.8	C31—C30—H30	120.1
C11—C12—C13	120.9 (5)	C30—C31—C32	121.0 (6)
C11—C12—H12	119.6	C30—C31—H31	119.5
C13—C12—H12	119.6	C32—C31—H31	119.5
C12—C13—N2	121.3 (5)	C27—C32—C31	118.7 (5)
C12—C13—C14	117.8 (5)	C27—C32—H32	120.6
N2—C13—C14	120.6 (5)	C31—C32—H32	120.6

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C32—H32 \cdots O2	0.93	2.39	3.105 (6)	134
C20—H20 \cdots O4	0.93	2.47	3.120 (6)	127
N1—H1B \cdots N2 ⁱ	0.865 (19)	2.43 (3)	3.211 (8)	150 (5)
N1—H1A \cdots O2 ⁱⁱ	0.86 (2)	2.43 (5)	3.114 (6)	137 (5)
N2—H2A \cdots N1 ⁱⁱⁱ	0.847 (19)	2.44 (2)	3.272 (7)	169 (5)
N2—H2B \cdots O4 ^{iv}	0.847 (19)	2.36 (3)	3.151 (6)	156 (5)
C14—H14 \cdots O4 ^{iv}	0.93	2.58	3.369 (6)	143
C12—H12 \cdots O2 ^v	0.93	2.56	3.433 (6)	157
C23—H23 \cdots Cg1 ^{vi}	0.93	2.72	3.567 (7)	151

Symmetry codes: (i) $x, -y+2, z+1/2$; (ii) $-x, -y+2, -z+1$; (iii) $-x, y-1, -z+1/2$; (iv) $-x, y, -z+1/2$; (v) $x, -y+1, z-1/2$; (vi) $-x+1/2, -y+3/2, -z+1$.

Fig. 1




Fig. 2

