

Tetraaquabis[2-(thiosemicarbazono-methyl)benzenesulfonato]manganese(II)

Xi-Shi Tai^{a*} and Zeng-Bing Zhao^b^aCollege of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China, and ^bDepartment of Chemistry, Qinghai Normal University, Xining 810008, People's Republic of China

Correspondence e-mail: taixishi@lzu.edu.cn


Received 27 October 2009; accepted 27 October 2009

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.028; wR factor = 0.077; data-to-parameter ratio = 13.0.

In the title compound, $[\text{Mn}(\text{C}_8\text{H}_8\text{N}_3\text{O}_3\text{S}_2)_2(\text{H}_2\text{O})_4]$, the Mn^{II} atom (site symmetry $\bar{1}$) adopts a slightly distorted octahedral MnO_6 geometry. The molecular conformation is supported by $\text{N}-\text{H}\cdots\text{N}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds. In the crystal, molecules interact by $\text{O}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{S}$, $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds, thereby forming (011) sheets.

Related literature

For background to coordination networks, see: Ranford *et al.* (1998).



Experimental

Crystal data

$[\text{Mn}(\text{C}_8\text{H}_8\text{N}_3\text{O}_3\text{S}_2)_2(\text{H}_2\text{O})_4]$
 $M_r = 643.59$

Triclinic, $P\bar{1}$
 $a = 6.8096$ (6) Å

$b = 9.5498$ (8) Å
 $c = 10.7898$ (9) Å
 $\alpha = 64.386$ (1)°
 $\beta = 88.495$ (1)°
 $\gamma = 83.791$ (1)°
 $V = 628.83$ (9) Å³

$Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.92$ mm⁻¹
 $T = 273$ K
 $0.19 \times 0.14 \times 0.12$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2000)
 $T_{\text{min}} = 0.845$, $T_{\text{max}} = 0.898$

3329 measured reflections
 2207 independent reflections
 2005 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.077$
 $S = 1.07$
 2207 reflections

170 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.36$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Selected bond lengths (Å).

Mn1—O4	2.1369 (17)	Mn1—O1	2.2166 (14)
Mn1—O5	2.1495 (16)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A ⁱ ⋯N3	0.86	2.29	2.638 (3)	104
O4—H10 ⁱ ⋯O3	0.85	2.02	2.761 (3)	146
N1—H1B ⁱ ⋯O3 ⁱ	0.86	2.33	2.986 (3)	134
N2—H2 ⁱⁱ ⋯S2 ⁱⁱⁱ	0.86	2.57	3.4231 (19)	170
O4—H9 ⁱⁱ ⋯S2 ⁱⁱⁱ	0.85	2.36	3.182 (2)	162
O5—H11 ⁱⁱ ⋯S2 ^{iv}	0.85	2.46	3.2603 (19)	156
O5—H12 ^v ⋯O2 ^v	0.85	1.87	2.712 (3)	172

Symmetry codes: (i) $x, y - 1, z + 1$; (ii) $-x + 2, -y, -z + 1$; (iii) $x - 1, y + 1, z - 1$; (iv) $x, y + 1, z - 1$; (v) $-x + 2, -y + 1, -z$.

Data collection: SMART (Bruker, 2000); cell refinement: SAINT (Bruker, 2000); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

This project was supported by the National Natural Science Foundation of China (20671073), the Natural Science Foundation of Shandong (Y2007B60) and the Science Foundation of Weifang University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5192).

References

- Bruker (2000). SMART, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Ranford, J. D., Vittal, J. J. & Wang, Y. M. (1998). *Inorg. Chem.* **37**, 1226–1231.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m1491 [doi:10.1107/S160053680904495X]

Tetraaquabis[2-(thiosemicarbazonomethyl)benzenesulfonato]manganese(II)

X.-S. Tai and Z.-B. Zhao

Experimental

A solution of 1.0 mmol 2-formyl-benzenesulfonate-thiosemicarbazide was added to a solution of 0.5 mmol $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ in 5 ml ethanol at room temperature. The mixture was refluxed for 4 h with stirring, then the resulting precipitate was filtered, washed, and dried *in vacuo* over P_4O_{10} for 48 h. Pink blocks of (I) were obtained by slowly evaporating from methanol at room temperature.

Refinement

The H atoms were positioned geometrically (C—H = 0.93, N—H = 0.86, O—H = 0.85 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$ or $1.5 U_{\text{eq}}(\text{O})$.

Figures




Fig. 1. The molecular structure of (I) showing 30% displacement ellipsoids. Atoms with the suffix A are generated by the symmetry operation (1-x, 1-y, -z).

Tetraaquabis[2-(thiosemicarbazonomethyl)benzenesulfonato]manganese(II)

Crystal data

$[\text{Mn}(\text{C}_8\text{H}_8\text{N}_3\text{O}_3\text{S}_2)_2(\text{H}_2\text{O})_4]$

$M_r = 643.59$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.8096$ (6) Å

$b = 9.5498$ (8) Å

$c = 10.7898$ (9) Å

$\alpha = 64.386$ (1)°

$\beta = 88.495$ (1)°

$\gamma = 83.791$ (1)°

$V = 628.83$ (9) Å³

$Z = 1$

$F_{000} = 331$

$D_x = 1.700$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2123 reflections

$\theta = 2.4\text{--}28.2^\circ$

$\mu = 0.92$ mm⁻¹

$T = 273$ K

Block, pink

$0.19 \times 0.14 \times 0.12$ mm

Data collection

Bruker SMART CCD diffractometer	2207 independent reflections
Radiation source: fine-focus sealed tube	2005 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.015$
$T = 273$ K	$\theta_{\text{max}} = 25.1^\circ$
ω scans	$\theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2000)	$h = -8 \rightarrow 8$
$T_{\text{min}} = 0.845$, $T_{\text{max}} = 0.898$	$k = -11 \rightarrow 11$
3329 measured reflections	$l = -9 \rightarrow 12$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.028$	$w = 1/[\sigma^2(F_o^2) + (0.0384P)^2 + 0.3206P]$
$wR(F^2) = 0.077$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\text{max}} = 0.001$
2207 reflections	$\Delta\rho_{\text{max}} = 0.36 \text{ e } \text{\AA}^{-3}$
170 parameters	$\Delta\rho_{\text{min}} = -0.34 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.049 (3)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.5000	0.5000	0.0000	0.02389 (16)
S1	0.73935 (8)	0.67535 (6)	0.15413 (5)	0.02518 (16)
S2	1.00275 (9)	-0.13908 (6)	0.71686 (6)	0.03340 (18)

O1	0.6334 (2)	0.54631 (17)	0.16197 (15)	0.0333 (4)
O2	0.9517 (2)	0.6407 (2)	0.15616 (17)	0.0417 (4)
O3	0.6673 (3)	0.82183 (18)	0.04065 (15)	0.0386 (4)
O4	0.3965 (3)	0.7431 (2)	-0.0981 (2)	0.0652 (6)
H9	0.2884	0.7905	-0.1404	0.098*
H10	0.4415	0.7985	-0.0637	0.098*
O5	0.7698 (2)	0.5559 (2)	-0.11007 (18)	0.0433 (4)
H12	0.8605	0.4911	-0.1165	0.065*
H11	0.8157	0.6433	-0.1363	0.065*
N1	0.8828 (3)	0.0620 (2)	0.81969 (19)	0.0400 (5)
H1A	0.8452	0.1545	0.8118	0.048*
H1B	0.8919	-0.0160	0.8996	0.048*
N2	0.9104 (3)	0.16402 (19)	0.58726 (18)	0.0271 (4)
H2	0.9474	0.1543	0.5143	0.033*
N3	0.8342 (2)	0.30737 (19)	0.57900 (17)	0.0257 (4)
C1	0.9263 (3)	0.0400 (2)	0.7097 (2)	0.0269 (5)
C2	0.8091 (3)	0.4167 (2)	0.4577 (2)	0.0242 (4)
H2A	0.8437	0.3975	0.3821	0.029*
C3	0.7252 (3)	0.5734 (2)	0.4382 (2)	0.0216 (4)
C4	0.6849 (3)	0.6967 (2)	0.3074 (2)	0.0212 (4)
C5	0.6053 (3)	0.8429 (2)	0.2938 (2)	0.0284 (5)
H5A	0.5780	0.9236	0.2066	0.034*
C6	0.5664 (3)	0.8694 (3)	0.4078 (2)	0.0345 (5)
H6	0.5140	0.9679	0.3979	0.041*
C7	0.6056 (3)	0.7491 (3)	0.5373 (2)	0.0329 (5)
H7	0.5791	0.7667	0.6147	0.040*
C8	0.6835 (3)	0.6034 (3)	0.5523 (2)	0.0283 (5)
H8	0.7090	0.5235	0.6401	0.034*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0270 (3)	0.0197 (2)	0.0237 (3)	-0.00156 (17)	-0.00312 (17)	-0.00820 (19)
S1	0.0331 (3)	0.0220 (3)	0.0199 (3)	-0.0055 (2)	0.0001 (2)	-0.0080 (2)
S2	0.0437 (3)	0.0198 (3)	0.0306 (3)	0.0006 (2)	-0.0044 (2)	-0.0057 (2)
O1	0.0521 (10)	0.0266 (8)	0.0230 (8)	-0.0120 (7)	-0.0042 (7)	-0.0104 (6)
O2	0.0352 (9)	0.0540 (11)	0.0448 (10)	-0.0042 (8)	0.0067 (7)	-0.0301 (9)
O3	0.0603 (11)	0.0259 (8)	0.0221 (8)	-0.0100 (7)	-0.0032 (7)	-0.0020 (7)
O4	0.0675 (13)	0.0263 (9)	0.0932 (16)	0.0101 (9)	-0.0507 (12)	-0.0182 (10)
O5	0.0405 (10)	0.0403 (10)	0.0559 (11)	-0.0125 (8)	0.0177 (8)	-0.0262 (9)
N1	0.0639 (14)	0.0245 (10)	0.0246 (10)	-0.0001 (9)	0.0020 (9)	-0.0051 (8)
N2	0.0313 (9)	0.0206 (9)	0.0239 (9)	0.0028 (7)	-0.0010 (7)	-0.0056 (7)
N3	0.0269 (9)	0.0203 (9)	0.0267 (10)	-0.0011 (7)	-0.0023 (7)	-0.0074 (8)
C1	0.0250 (11)	0.0235 (11)	0.0261 (11)	-0.0033 (8)	-0.0033 (8)	-0.0047 (9)
C2	0.0263 (10)	0.0228 (10)	0.0225 (10)	-0.0017 (8)	-0.0020 (8)	-0.0091 (9)
C3	0.0206 (10)	0.0211 (10)	0.0238 (10)	-0.0031 (8)	-0.0027 (8)	-0.0100 (8)
C4	0.0217 (10)	0.0212 (10)	0.0216 (10)	-0.0043 (8)	-0.0016 (8)	-0.0095 (8)
C5	0.0320 (11)	0.0194 (10)	0.0308 (12)	-0.0016 (9)	-0.0024 (9)	-0.0080 (9)

supplementary materials

C6	0.0375 (13)	0.0260 (12)	0.0450 (14)	0.0009 (10)	-0.0004 (10)	-0.0210 (11)
C7	0.0362 (12)	0.0367 (13)	0.0361 (12)	-0.0053 (10)	0.0022 (10)	-0.0250 (11)
C8	0.0306 (11)	0.0297 (11)	0.0240 (11)	-0.0039 (9)	-0.0013 (9)	-0.0108 (9)

Geometric parameters (\AA , $^\circ$)

Mn1—O4 ⁱ	2.1369 (17)	N1—H1B	0.8600
Mn1—O4	2.1369 (17)	N2—C1	1.337 (3)
Mn1—O5 ⁱ	2.1495 (16)	N2—N3	1.376 (2)
Mn1—O5	2.1495 (16)	N2—H2	0.8600
Mn1—O1	2.2166 (14)	N3—C2	1.274 (3)
Mn1—O1 ⁱ	2.2166 (14)	C2—C3	1.469 (3)
S1—O2	1.4468 (17)	C2—H2A	0.9300
S1—O3	1.4489 (16)	C3—C8	1.396 (3)
S1—O1	1.4647 (15)	C3—C4	1.403 (3)
S1—C4	1.7767 (19)	C4—C5	1.388 (3)
S2—C1	1.702 (2)	C5—C6	1.374 (3)
O4—H9	0.8500	C5—H5A	0.9300
O4—H10	0.8499	C6—C7	1.381 (3)
O5—H12	0.8499	C6—H6	0.9300
O5—H11	0.8499	C7—C8	1.376 (3)
N1—C1	1.313 (3)	C7—H7	0.9300
N1—H1A	0.8600	C8—H8	0.9300
O4 ⁱ —Mn1—O4	180.0	H1A—N1—H1B	120.0
O4 ⁱ —Mn1—O5 ⁱ	88.04 (8)	C1—N2—N3	119.42 (18)
O4—Mn1—O5 ⁱ	91.96 (8)	C1—N2—H2	120.3
O4 ⁱ —Mn1—O5	91.96 (8)	N3—N2—H2	120.3
O4—Mn1—O5	88.04 (8)	C2—N3—N2	115.48 (18)
O5 ⁱ —Mn1—O5	180.0	N1—C1—N2	118.31 (19)
O4 ⁱ —Mn1—O1	92.72 (7)	N1—C1—S2	122.79 (16)
O4—Mn1—O1	87.28 (7)	N2—C1—S2	118.89 (16)
O5 ⁱ —Mn1—O1	92.43 (6)	N3—C2—C3	119.54 (19)
O5—Mn1—O1	87.57 (6)	N3—C2—H2A	120.2
O4 ⁱ —Mn1—O1 ⁱ	87.28 (7)	C3—C2—H2A	120.2
O4—Mn1—O1 ⁱ	92.72 (7)	C8—C3—C4	117.81 (18)
O5 ⁱ —Mn1—O1 ⁱ	87.57 (6)	C8—C3—C2	119.86 (18)
O5—Mn1—O1 ⁱ	92.43 (6)	C4—C3—C2	122.33 (18)
O1—Mn1—O1 ⁱ	180.0	C5—C4—C3	120.37 (18)
O2—S1—O3	113.09 (10)	C5—C4—S1	117.47 (15)
O2—S1—O1	112.81 (10)	C3—C4—S1	122.14 (15)
O3—S1—O1	111.89 (9)	C6—C5—C4	120.6 (2)
O2—S1—C4	105.67 (9)	C6—C5—H5A	119.7
O3—S1—C4	106.74 (9)	C4—C5—H5A	119.7
O1—S1—C4	105.98 (9)	C5—C6—C7	119.6 (2)
S1—O1—Mn1	131.56 (9)	C5—C6—H6	120.2
Mn1—O4—H9	131.3	C7—C6—H6	120.2

Mn1—O4—H10	115.1	C8—C7—C6	120.3 (2)
H9—O4—H10	108.1	C8—C7—H7	119.8
Mn1—O5—H12	126.2	C6—C7—H7	119.8
Mn1—O5—H11	124.0	C7—C8—C3	121.2 (2)
H12—O5—H11	108.1	C7—C8—H8	119.4
C1—N1—H1A	120.0	C3—C8—H8	119.4
C1—N1—H1B	120.0		
O2—S1—O1—Mn1	-96.18 (14)	C8—C3—C4—S1	177.75 (15)
O3—S1—O1—Mn1	32.68 (16)	C2—C3—C4—S1	-2.0 (3)
C4—S1—O1—Mn1	148.66 (12)	O2—S1—C4—C5	115.31 (17)
O4 ⁱ —Mn1—O1—S1	138.91 (14)	O3—S1—C4—C5	-5.33 (19)
O4—Mn1—O1—S1	-41.09 (14)	O1—S1—C4—C5	-124.75 (17)
O5 ⁱ —Mn1—O1—S1	-132.94 (13)	O2—S1—C4—C3	-62.83 (18)
O5—Mn1—O1—S1	47.06 (13)	O3—S1—C4—C3	176.53 (15)
O1 ⁱ —Mn1—O1—S1	-25 (100)	O1—S1—C4—C3	57.11 (18)
C1—N2—N3—C2	-174.55 (18)	C3—C4—C5—C6	0.6 (3)
N3—N2—C1—N1	-5.3 (3)	S1—C4—C5—C6	-177.58 (17)
N3—N2—C1—S2	174.98 (14)	C4—C5—C6—C7	-0.5 (3)
N2—N3—C2—C3	178.94 (17)	C5—C6—C7—C8	0.2 (3)
N3—C2—C3—C8	3.4 (3)	C6—C7—C8—C3	0.1 (3)
N3—C2—C3—C4	-176.81 (18)	C4—C3—C8—C7	0.0 (3)
C8—C3—C4—C5	-0.3 (3)	C2—C3—C8—C7	179.78 (19)
C2—C3—C4—C5	179.89 (19)		

Symmetry codes: (i) $-x+1, -y+1, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots N3	0.86	2.29	2.638 (3)	104
O4—H10 \cdots O3	0.85	2.02	2.761 (3)	146
N1—H1B \cdots O3 ⁱⁱ	0.86	2.33	2.986 (3)	134
N2—H2 \cdots S2 ⁱⁱⁱ	0.86	2.57	3.4231 (19)	170
O4—H9 \cdots S2 ^{iv}	0.85	2.36	3.182 (2)	162
O5—H11 \cdots S2 ^v	0.85	2.46	3.2603 (19)	156
O5—H12 \cdots O2 ^{vi}	0.85	1.87	2.712 (3)	172

Symmetry codes: (ii) $x, y-1, z+1$; (iii) $-x+2, -y, -z+1$; (iv) $x-1, y+1, z-1$; (v) $x, y+1, z-1$; (vi) $-x+2, -y+1, -z$.

Fig. 1

