

Diaquabis(4-oxo-1,4-dihydropyridine-3-sulfonato- $\kappa^2 O^3, O^4$)zinc(II)

Zhi-Biao Zhu,^a Shan Gao^a and Seik Weng Ng^{b*}

^aCollege of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 27 October 2009; accepted 27 October 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.025; wR factor = 0.074; data-to-parameter ratio = 13.7.

In the crystal structure of the title compound, $[\text{Zn}(\text{C}_5\text{H}_4\text{NO}_4\text{S})_2(\text{H}_2\text{O})_2]$, the 4-oxo-1,4-dihydropyridine-3-sulfonate anion chelates to water-coordinated zinc centres through the carbonyl O atom and through one O atom of the sulfonate group. The Zn^{II} atom lies on a center of inversion, and adjacent molecules are linked by $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming a three-dimensional network.

Related literature

For the structure of the 4-oxo-1,4-dihydropyridine-3-sulfonate anion, see: Zhu *et al.* (2009).



Experimental

Crystal data

$[\text{Zn}(\text{C}_5\text{H}_4\text{NO}_4\text{S})_2(\text{H}_2\text{O})_2]$	$c = 7.4437 (2)\text{ \AA}$
$M_r = 449.71$	$\beta = 98.9371 (9)^\circ$
Monoclinic, $P2_1/c$	$V = 759.01 (3)\text{ \AA}^3$
$a = 4.9263 (1)\text{ \AA}$	$Z = 2$
$b = 20.9529 (6)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 1.95\text{ mm}^{-1}$
 $T = 293\text{ K}$

$0.23 \times 0.17 \times 0.14\text{ mm}$

Data collection

Rigaku R-AXIS RAPID IP
diffractometer
Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.662$, $T_{\max} = 0.772$

7334 measured reflections
1738 independent reflections
1629 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.021$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.025$
 $wR(F^2) = 0.074$
 $S = 1.10$
1738 reflections
127 parameters
3 restraints

H atoms treated by a mixture of
independent and constrained
refinement
 $\Delta\rho_{\max} = 0.45\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.46\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1w-H1 \cdots O2 ⁱ	0.84 (1)	2.05 (2)	2.797 (2)	147 (2)
O1w-H2 \cdots O4 ⁱⁱ	0.84 (1)	1.92 (1)	2.744 (2)	171 (3)
N1-H3 \cdots O3 ⁱⁱⁱ	0.85 (1)	1.91 (1)	2.754 (2)	175 (3)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x + 1, y, z$; (iii) $x - 1, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Natural Science Foundation of Heilongjiang Province (No. B200501), Heilongjiang University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5120).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.
- Zhu, Z.-B., Gao, S. & Ng, S. W. (2009). *Acta Cryst. E* **65**, o2687.

supplementary materials

Acta Cryst. (2009). E65, m1493 [doi:10.1107/S1600536809044948]

Diaquabis(4-oxo-1,4-dihydropyridine-3-sulfonato- $\kappa^2 O^3, O^4$)zinc(II)

Z.-B. Zhu, S. Gao and S. W. Ng

Experimental

Zinc carbonate (0.25 g, 2 mmol) was added to a hot aqueous solution of 4-hydroxypyridine-3-sulfonic acid (0.35 g, 2 mmol); the pH value was adjusted to 6 with 0.1 M sodium hydroxide. The solution was allowed to evaporate slowly at room temperature, and colorless prismatic crystals were isolated after about five days. CH&N elemental analysis. Calc. for $C_{10}H_{12}N_2O_{10}S_2Zn$: C 26.71, H 2.69, N 6.23%; found: C 26.73, H 2.73, N 6.21%.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. The amino and water H-atoms were located in a difference Fourier map, and were refined isotropically with a distance restraint of N—H = O—H = 0.85 ± 0.01 Å.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of $Zn(H_2O)_2(C_5H_4NO_4S)_2$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

Diaquabis(4-oxo-1,4-dihydropyridine-3-sulfonato- $\kappa^2 O^3, O^4$)zinc(II)

Crystal data

$[Zn(C_5H_4NO_4S)_2(H_2O)_2]$	$F_{000} = 456$
$M_r = 449.71$	$D_x = 1.968 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Hall symbol: -P 2ybc	Cell parameters from 6865 reflections
$a = 4.9263$ (1) Å	$\theta = 3.4\text{--}27.5^\circ$
$b = 20.9529$ (6) Å	$\mu = 1.95 \text{ mm}^{-1}$
$c = 7.4437$ (2) Å	$T = 293$ K
$\beta = 98.9371$ (9)°	Prism, colorless
$V = 759.01$ (3) Å ³	$0.23 \times 0.17 \times 0.14$ mm
$Z = 2$	

Data collection

Rigaku R-AXIS RAPID IP diffractometer	1738 independent reflections
Radiation source: fine-focus sealed tube	1629 reflections with $I > 2\sigma(I)$

supplementary materials

Monochromator: graphite	$R_{\text{int}} = 0.021$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 3.4^\circ$
Absorption correction: Multi-scan (ABSCOR; Higashi, 1995)	$h = -6 \rightarrow 5$
$T_{\text{min}} = 0.662, T_{\text{max}} = 0.772$	$k = -27 \rightarrow 27$
7334 measured reflections	$l = -9 \rightarrow 9$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.025$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.074$	$w = 1/[\sigma^2(F_o^2) + (0.0399P)^2 + 0.5207P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\text{max}} = 0.001$
1738 reflections	$\Delta\rho_{\text{max}} = 0.45 \text{ e \AA}^{-3}$
127 parameters	$\Delta\rho_{\text{min}} = -0.45 \text{ e \AA}^{-3}$
3 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.5000	0.5000	0.01991 (11)
S1	0.54548 (9)	0.35166 (2)	0.43742 (6)	0.02251 (13)
O1	0.6943 (3)	0.41242 (6)	0.46588 (19)	0.0246 (3)
O2	0.4026 (3)	0.33653 (7)	0.5890 (2)	0.0327 (3)
O3	0.7182 (3)	0.30080 (7)	0.3867 (2)	0.0380 (4)
O4	0.1877 (3)	0.47222 (6)	0.29786 (18)	0.0238 (3)
O1W	0.7120 (3)	0.54012 (7)	0.3004 (2)	0.0279 (3)
N1	-0.0061 (4)	0.30854 (10)	0.0136 (2)	0.0361 (5)
C1	0.2078 (4)	0.30795 (10)	0.1490 (3)	0.0303 (4)
H1A	0.3053	0.2703	0.1764	0.036*
C2	0.2849 (4)	0.36189 (8)	0.2474 (3)	0.0221 (4)
C3	0.1359 (4)	0.42039 (8)	0.2098 (2)	0.0209 (4)
C4	-0.0876 (4)	0.41740 (10)	0.0613 (3)	0.0294 (4)
H4	-0.1897	0.4540	0.0274	0.035*
C5	-0.1522 (5)	0.36232 (11)	-0.0300 (3)	0.0353 (5)
H5A	-0.2998	0.3615	-0.1245	0.042*
H1	0.738 (5)	0.5794 (5)	0.318 (4)	0.035 (7)*
H2	0.866 (3)	0.5233 (13)	0.301 (4)	0.040 (7)*
H3	-0.081 (6)	0.2742 (9)	-0.028 (4)	0.051 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.01975 (17)	0.01417 (16)	0.02420 (18)	-0.00106 (10)	-0.00165 (12)	-0.00169 (10)
S1	0.0258 (2)	0.0128 (2)	0.0270 (2)	0.00172 (16)	-0.00212 (18)	0.00056 (16)
O1	0.0221 (6)	0.0168 (6)	0.0333 (7)	0.0005 (5)	-0.0006 (5)	-0.0015 (5)
O2	0.0433 (8)	0.0247 (7)	0.0286 (7)	-0.0061 (6)	0.0009 (6)	0.0042 (6)
O3	0.0398 (8)	0.0203 (7)	0.0503 (10)	0.0124 (6)	-0.0037 (7)	-0.0046 (6)
O4	0.0226 (6)	0.0168 (6)	0.0295 (7)	0.0010 (5)	-0.0042 (5)	-0.0031 (5)
O1W	0.0263 (7)	0.0244 (7)	0.0333 (8)	0.0020 (6)	0.0059 (6)	0.0008 (6)
N1	0.0443 (11)	0.0297 (10)	0.0324 (10)	-0.0131 (8)	0.0006 (8)	-0.0141 (7)
C1	0.0366 (11)	0.0203 (9)	0.0337 (10)	-0.0038 (8)	0.0046 (8)	-0.0065 (8)
C2	0.0253 (9)	0.0178 (8)	0.0224 (9)	-0.0022 (7)	0.0008 (7)	-0.0024 (6)
C3	0.0215 (8)	0.0189 (8)	0.0212 (8)	-0.0032 (7)	-0.0005 (7)	0.0008 (6)
C4	0.0300 (10)	0.0295 (10)	0.0256 (9)	-0.0034 (8)	-0.0061 (8)	0.0025 (8)
C5	0.0364 (11)	0.0414 (12)	0.0248 (10)	-0.0108 (9)	-0.0056 (8)	-0.0042 (8)

Geometric parameters (\AA , $^\circ$)

Zn1—O4	2.0618 (12)	O1W—H2	0.835 (10)
Zn1—O4 ⁱ	2.0618 (12)	N1—C1	1.340 (3)
Zn1—O1 ⁱ	2.1031 (13)	N1—C5	1.349 (3)
Zn1—O1	2.1031 (13)	N1—H3	0.846 (10)
Zn1—O1W ⁱ	2.1183 (15)	C1—C2	1.368 (3)
Zn1—O1W	2.1183 (15)	C1—H1A	0.9300
S1—O3	1.4496 (15)	C2—C3	1.434 (2)
S1—O2	1.4547 (17)	C3—C4	1.435 (2)
S1—O1	1.4681 (13)	C4—C5	1.352 (3)
S1—C2	1.7694 (19)	C4—H4	0.9300
O4—C3	1.274 (2)	C5—H5A	0.9300
O1W—H1	0.841 (10)		
O4—Zn1—O4 ⁱ	180.00 (7)	C3—O4—Zn1	133.12 (12)
O4—Zn1—O1 ⁱ	91.86 (5)	Zn1—O1W—H1	111.3 (18)
O4 ⁱ —Zn1—O1 ⁱ	88.14 (5)	Zn1—O1W—H2	112 (2)
O4—Zn1—O1	88.14 (5)	H1—O1W—H2	107 (3)
O4 ⁱ —Zn1—O1	91.86 (5)	C1—N1—C5	121.09 (18)
O1 ⁱ —Zn1—O1	180.0	C1—N1—H3	121 (2)
O4—Zn1—O1W ⁱ	90.38 (6)	C5—N1—H3	116 (2)
O4 ⁱ —Zn1—O1W ⁱ	89.62 (6)	N1—C1—C2	121.0 (2)
O1 ⁱ —Zn1—O1W ⁱ	88.76 (5)	N1—C1—H1A	119.5
O1—Zn1—O1W ⁱ	91.24 (5)	C2—C1—H1A	119.5
O4—Zn1—O1W	89.62 (6)	C1—C2—C3	120.70 (17)
O4 ⁱ —Zn1—O1W	90.38 (6)	C1—C2—S1	115.75 (15)
O1 ⁱ —Zn1—O1W	91.24 (5)	C3—C2—S1	123.06 (13)
O1—Zn1—O1W	88.76 (5)	O4—C3—C2	124.95 (16)

supplementary materials

O1W ⁱ —Zn1—O1W	180.00 (7)	O4—C3—C4	120.14 (17)
O3—S1—O2	114.58 (10)	C2—C3—C4	114.91 (16)
O3—S1—O1	112.02 (9)	C5—C4—C3	121.1 (2)
O2—S1—O1	111.59 (9)	C5—C4—H4	119.4
O3—S1—C2	105.27 (9)	C3—C4—H4	119.4
O2—S1—C2	105.55 (9)	C4—C5—N1	121.12 (18)
O1—S1—C2	107.12 (8)	C4—C5—H5A	119.4
S1—O1—Zn1	123.21 (8)	N1—C5—H5A	119.4
O3—S1—O1—Zn1	170.03 (10)	O2—S1—C2—C1	-89.44 (17)
O2—S1—O1—Zn1	-59.98 (12)	O1—S1—C2—C1	151.53 (16)
C2—S1—O1—Zn1	55.08 (12)	O3—S1—C2—C3	-155.77 (16)
O4—Zn1—O1—S1	-40.93 (10)	O2—S1—C2—C3	82.66 (17)
O4 ⁱ —Zn1—O1—S1	139.07 (10)	O1—S1—C2—C3	-36.38 (18)
O1W ⁱ —Zn1—O1—S1	49.40 (10)	Zn1—O4—C3—C2	6.9 (3)
O1W—Zn1—O1—S1	-130.60 (10)	Zn1—O4—C3—C4	-173.69 (14)
O1 ⁱ —Zn1—O4—C3	-173.83 (17)	C1—C2—C3—O4	177.73 (19)
O1—Zn1—O4—C3	6.17 (17)	S1—C2—C3—O4	6.0 (3)
O1W ⁱ —Zn1—O4—C3	-85.06 (18)	C1—C2—C3—C4	-1.7 (3)
O1W—Zn1—O4—C3	94.94 (18)	S1—C2—C3—C4	-173.44 (15)
C5—N1—C1—C2	0.0 (3)	O4—C3—C4—C5	-177.7 (2)
N1—C1—C2—C3	0.9 (3)	C2—C3—C4—C5	1.8 (3)
N1—C1—C2—S1	173.18 (17)	C3—C4—C5—N1	-1.0 (4)
O3—S1—C2—C1	32.14 (19)	C1—N1—C5—C4	0.0 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O1w—H1 ⁱ —O2 ⁱ	0.84 (1)	2.05 (2)	2.797 (2)	147 (2)
O1w—H2 ⁱⁱ —O4 ⁱⁱ	0.84 (1)	1.92 (1)	2.744 (2)	171 (3)
N1—H3 ⁱⁱⁱ —O3 ⁱⁱⁱ	0.85 (1)	1.91 (1)	2.754 (2)	175 (3)

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $x+1, y, z$; (iii) $x-1, -y+1/2, z-1/2$.

Fig. 1

