

$\alpha = 97.895(3)^\circ$
 $\beta = 98.520(3)^\circ$
 $\gamma = 91.661(3)^\circ$
 $V = 1080.41(10)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.28\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.21 \times 0.09 \times 0.06\text{ mm}$

2-(Phenylsulfanyl)pyridine-3-carboxylic acid

Muhammad Naeem Khan,^a‡ Misbahul Ain Khan,^a
Muhammad Nadeem Arshad,^b Islam Ullah Khan^{b*} and
Salma Rehman^c

^aDepartment of Chemistry, Islamia University, Bahawalpur, Pakistan, ^bMaterials Chemistry laboratory, Department of Chemistry, GC University, Lahore, Pakistan, and ^cApplied Chemistry Research Center, PCSIR Laboratories Complex, Lahore 54600, Pakistan

Correspondence e-mail: iukhan.gcu@gmail.com


Received 19 September 2009; accepted 23 September 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.006\text{ \AA}$; R factor = 0.073; wR factor = 0.224; data-to-parameter ratio = 18.5.

The title compound, $C_{12}H_9NO_2S$, belongs to the nitrogen-containing group of heterocyclic organic compounds and crystallized with two molecules per asymmetric unit. In the crystal, both molecules form inversion dimers linked by pairs of O—H—O hydrogen bonds. Weak symmetry-related C—H—O interactions link the carboxyl dimers along b axis. The dihedral angle between the two aromatic rings in the two molecules are 55.75 (14) and 58.33 (13)°.

Related literature

For the pharmacological effects of heteroaromatic antitumor compounds: Denny *et al.* (1982); Fujiwara (1997); Antonini & Martelli (1992); Cholody *et al.* (1992). For the title compound as an intermediate for heterocycles, see: Khan *et al.* (2008a,b). For the synthesis, see: Mann & Reid (1952).



Experimental

Crystal data

$C_{12}H_9NO_2S$
 $M_r = 231.26$
Triclinic, $P\bar{1}$
 $a = 7.2201(4)\text{ \AA}$
 $b = 7.6653(4)\text{ \AA}$
 $c = 19.9537(11)\text{ \AA}$

Data collection

Bruker Kappa APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.911$, $T_{\max} = 0.983$

23200 measured reflections
5397 independent reflections
2766 reflections with $I > 2s(I)$
 $R_{\text{int}} = 0.063$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.073$
 $wR(F^2) = 0.224$
 $S = 1.09$
5397 reflections

292 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.59\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.53\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2···O3 ⁱ	0.82	1.82	2.624 (2)	167 (1)
O4—H4o···O1 ⁱ	0.82	1.83	2.642 (2)	170 (1)
C3—H3···O4 ⁱⁱ	0.93	2.50	3.264 (5)	139
C4—H4···O1 ⁱⁱⁱ	0.93	2.55	3.458 (5)	164
C15—H15···O2 ⁱⁱ	0.93	2.54	3.294 (5)	138
C16—H16···O3 ⁱⁱⁱ	0.93	2.58	3.467 (5)	160

Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + 1, -y + 1, -z$; (iii) $x, y + 1, z$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON*.

The authors acknowledge the Higher Education Commission of Pakistan for providing a grant under the project strengthening the Materials Chemistry Laboratory at GC University, Lahore.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HG2570).

References

- Antonini, I. & Martelli, S. (1992). *J. Heterocycl. Chem.* **29**, 471–473.
Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cholody, W. M., Martelli, S. & Konopa, J. (1992). *J. Med. Chem.* **35**, 276–315.
Denny, W. A., Cain, B. F., Atwell, G. J., Hansch, C., Panthanickal, A. & Leo, A. (1982). *J. Med. Chem.* **25**, 276–315.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Fujiwara, H. (1997). *Heterocycles*, **45**, 119–127.
Khan, M. N., Tahir, M. N., Khan, M. A., Khan, I. U. & Arshad, M. N. (2008a). *Acta Cryst. E64*, o730.
Khan, M. N., Tahir, M. N., Khan, M. A., Khan, I. U. & Arshad, M. N. (2008b). *Acta Cryst. E64*, o1704.
Mann, F. G. & Reid, J. A. (1952). *J. Chem. Soc.* pp. 2057–2062.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

‡ Applied Chemistry Research Center, PCSIR Laboratories Complex, Lahore 54600, Pakistan.

supplementary materials

Acta Cryst. (2009). E65, o2662 [doi:10.1107/S1600536809038586]

2-(Phenylsulfanyl)pyridine-3-carboxylic acid

M. N. Khan, M. A. Khan, M. N. Arshad, I. U. Khan and S. Rehman

Comment

In continuation of our studies on pyridine-containing heterocyclic compounds, the title compound was synthesized. It is an intermediate for our previously reported crystal structures of 7-nitro-5*H*-thiochromeno[2,3-*b*]pyridin-5-one (Khan *et al.*, 2008a) and 5*H*-thiochromeno[2,3-*b*]pyridin-5-one (Khan *et al.*, 2008b). Pyridine containing compounds are widely distributed in nature. Heteroaromatic antitumor compounds have been prepared in recent years with the hope of increasing pharmacological effects (Denny *et al.*, 1982), (Fujiwara, 1997), (Antonini & Martelli, 1992) (Cholody *et al.*, 1992).

The title compound was crystallized with two independent molecules in the asymmetric unit (Fig 1). The dihedral angles between the two aromatic rings in molecule A and molecule B are 55.75 (14) $^{\circ}$ and 58.33 (13) $^{\circ}$ respectively. The carboxylic group present in each molecule forms dimers which are linked through weak C—H—O type interaction along the *b* axis to stabilize the structure Table. 1 & Fig. 2.

Experimental

A mixture of 2-chloronicotinic acid (1.57 g, 10 mmol) and thiophenol (2 ml) was heated under reflux for two hours to produce 2-(Phenylsulfanyl)pyridine-3-carboxylic acid (Mann & Reid, 1952). Suitable crystals for X-ray diffractions were obtained on cooling the saturated solution of (I) in ethanol.

Refinement

The H-atoms for aromatic carbons and carboxylic O atoms were refined geometrically and treated as riding atoms: C—H = 0.93 Å with $U_{\text{iso}}(\text{H})$ = 1.2 and O—H = 0.82 with $U_{\text{iso}}(\text{H})$ = 1.5.

Figures




Fig. 1. The molecular structure diagram of the title compound showing the atom labels. Thermal ellipsoids are drawn at the 50% probability level.

supplementary materials




Fig. 2. Unit cell packing diagram showing the intermolecular hydrogen bonding using dashed lines. The hydrogen atoms not involved in hydrogen bonding have been omitted.

2-(Phenylsulfanyl)pyridine-3-carboxylic acid

Crystal data

C ₁₂ H ₉ NO ₂ S	Z = 4
M _r = 231.26	F ₀₀₀ = 480
Triclinic, P <bar{1}< td=""><td>D_x = 1.422 Mg m⁻³</td></bar{1}<>	D _x = 1.422 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 7.2201 (4) Å	Cell parameters from 3133 reflections
b = 7.6653 (4) Å	θ = 2.9–23.8°
c = 19.9537 (11) Å	μ = 0.28 mm ⁻¹
α = 97.895 (3)°	T = 296 K
β = 98.520 (3)°	Needle, white
γ = 91.661 (3)°	0.21 × 0.09 × 0.06 mm
V = 1080.41 (10) Å ³	

Data collection

Bruker Kappa APEXII CCD diffractometer	5397 independent reflections
Radiation source: fine-focus sealed tube	2766 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.063$
T = 296 K	$\theta_{\text{max}} = 28.4^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.0^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -9 \rightarrow 9$
$T_{\text{min}} = 0.911$, $T_{\text{max}} = 0.983$	$k = -10 \rightarrow 10$
23200 measured reflections	$l = -26 \rightarrow 26$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.073$	$w = 1/[\sigma^2(F_{\text{o}}^2) + (0.1023P)^2 + 0.1581P]$

$wR(F^2) = 0.224$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.09$	$(\Delta/\sigma)_{\max} < 0.001$
5397 reflections	$\Delta\rho_{\max} = 0.59 \text{ e \AA}^{-3}$
292 parameters	$\Delta\rho_{\min} = -0.52 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$
Secondary atom site location: difference Fourier map	Extinction coefficient: 0.045 (5)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.15013 (14)	-0.12613 (12)	0.21464 (5)	0.0407 (3)
S2	0.70162 (15)	0.36894 (12)	0.22267 (5)	0.0447 (3)
O1	0.1649 (4)	-0.2094 (3)	0.07364 (13)	0.0508 (7)
O2	0.2377 (6)	-0.0151 (4)	0.00790 (14)	0.0709 (10)
H2	0.2653	-0.1043	-0.0155	0.106*
O3	0.7077 (5)	0.2844 (3)	0.08218 (13)	0.0568 (8)
O4	0.7667 (5)	0.4795 (4)	0.01516 (15)	0.0689 (10)
H4O	0.7958	0.3911	-0.0083	0.103*
N1	0.1813 (5)	0.2239 (4)	0.23401 (15)	0.0422 (8)
N2	0.6430 (4)	0.7119 (4)	0.23511 (15)	0.0414 (8)
C1	0.1727 (5)	0.0818 (4)	0.18659 (17)	0.0341 (8)
C2	0.1832 (5)	0.0971 (5)	0.11768 (18)	0.0382 (8)
C3	0.1939 (6)	0.2639 (5)	0.1002 (2)	0.0496 (10)
H3	0.2005	0.2783	0.0551	0.059*
C4	0.1951 (6)	0.4099 (5)	0.1488 (2)	0.0521 (11)
H4	0.1978	0.5234	0.1372	0.062*
C5	0.1921 (6)	0.3809 (5)	0.2150 (2)	0.0485 (10)
H5	0.1981	0.4787	0.2486	0.058*
C6	0.1937 (6)	-0.0559 (5)	0.06520 (18)	0.0413 (9)
C7	0.1766 (5)	-0.0719 (5)	0.30494 (18)	0.0380 (8)
C8	0.3176 (6)	-0.1484 (5)	0.34292 (19)	0.0490 (10)
H8	0.4002	-0.2190	0.3214	0.059*
C9	0.3357 (7)	-0.1189 (7)	0.4143 (2)	0.0656 (13)

supplementary materials

H9	0.4306	-0.1709	0.4403	0.079*
C10	0.2159 (8)	-0.0149 (7)	0.4461 (2)	0.0738 (15)
H10	0.2290	0.0048	0.4936	0.089*
C11	0.0762 (7)	0.0605 (6)	0.4077 (2)	0.0665 (14)
H11	-0.0053	0.1320	0.4295	0.080*
C12	0.0542 (6)	0.0326 (5)	0.3379 (2)	0.0509 (10)
H12	-0.0426	0.0836	0.3125	0.061*
C13	0.6743 (5)	0.5728 (5)	0.19104 (18)	0.0357 (8)
C14	0.6895 (5)	0.5891 (5)	0.12249 (18)	0.0373 (8)
C15	0.6724 (6)	0.7539 (5)	0.1026 (2)	0.0457 (10)
H15	0.6818	0.7686	0.0577	0.055*
C16	0.6416 (6)	0.8959 (5)	0.1483 (2)	0.0463 (10)
H16	0.6317	1.0081	0.1358	0.056*
C17	0.6261 (5)	0.8659 (5)	0.2134 (2)	0.0442 (9)
H17	0.6017	0.9612	0.2444	0.053*
C18	0.7230 (6)	0.4371 (5)	0.07237 (18)	0.0416 (9)
C19	0.7107 (5)	0.4283 (5)	0.31237 (18)	0.0380 (8)
C20	0.5879 (6)	0.3405 (6)	0.3442 (2)	0.0492 (10)
H20	0.4968	0.2604	0.3184	0.059*
C21	0.5999 (7)	0.3716 (7)	0.4149 (2)	0.0658 (13)
H21	0.5161	0.3128	0.4363	0.079*
C22	0.7329 (8)	0.4871 (7)	0.4529 (2)	0.0717 (14)
H22	0.7415	0.5066	0.5004	0.086*
C23	0.8545 (7)	0.5748 (6)	0.4213 (2)	0.0683 (14)
H23	0.9445	0.6552	0.4476	0.082*
C24	0.8463 (6)	0.5467 (6)	0.3516 (2)	0.0528 (11)
H24	0.9308	0.6064	0.3308	0.063*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0630 (7)	0.0289 (5)	0.0303 (5)	0.0032 (4)	0.0088 (4)	0.0027 (4)
S2	0.0728 (7)	0.0313 (5)	0.0326 (5)	0.0075 (5)	0.0147 (5)	0.0060 (4)
O1	0.092 (2)	0.0293 (15)	0.0329 (15)	0.0027 (13)	0.0166 (14)	0.0041 (12)
O2	0.148 (3)	0.0337 (16)	0.0387 (17)	0.0088 (18)	0.0413 (19)	0.0042 (13)
O3	0.111 (2)	0.0291 (15)	0.0366 (16)	0.0060 (14)	0.0284 (15)	0.0063 (12)
O4	0.137 (3)	0.0358 (16)	0.0428 (18)	0.0118 (18)	0.0420 (19)	0.0057 (13)
N1	0.062 (2)	0.0298 (17)	0.0348 (18)	0.0000 (14)	0.0090 (15)	0.0022 (14)
N2	0.056 (2)	0.0325 (17)	0.0378 (18)	0.0111 (14)	0.0137 (15)	0.0047 (14)
C1	0.041 (2)	0.0320 (19)	0.0291 (19)	0.0071 (15)	0.0034 (15)	0.0037 (15)
C2	0.054 (2)	0.0300 (19)	0.0309 (19)	0.0066 (16)	0.0075 (16)	0.0034 (15)
C3	0.078 (3)	0.034 (2)	0.040 (2)	0.0069 (19)	0.016 (2)	0.0065 (18)
C4	0.084 (3)	0.028 (2)	0.048 (3)	0.0085 (19)	0.016 (2)	0.0099 (18)
C5	0.068 (3)	0.029 (2)	0.046 (2)	0.0008 (18)	0.013 (2)	-0.0059 (18)
C6	0.065 (3)	0.032 (2)	0.029 (2)	0.0089 (17)	0.0106 (17)	0.0056 (16)
C7	0.053 (2)	0.0293 (19)	0.031 (2)	-0.0026 (16)	0.0100 (17)	0.0019 (15)
C8	0.063 (3)	0.048 (2)	0.038 (2)	0.005 (2)	0.0113 (19)	0.0104 (19)
C9	0.077 (3)	0.080 (3)	0.041 (3)	-0.003 (3)	0.003 (2)	0.021 (2)

C10	0.101 (4)	0.087 (4)	0.032 (2)	-0.020 (3)	0.020 (3)	0.000 (3)
C11	0.087 (4)	0.060 (3)	0.054 (3)	-0.001 (3)	0.035 (3)	-0.010 (2)
C12	0.063 (3)	0.048 (2)	0.045 (2)	0.004 (2)	0.020 (2)	0.0043 (19)
C13	0.044 (2)	0.0323 (19)	0.0295 (19)	0.0029 (15)	0.0065 (15)	0.0008 (15)
C14	0.052 (2)	0.0310 (19)	0.0305 (19)	0.0016 (16)	0.0122 (16)	0.0042 (15)
C15	0.067 (3)	0.032 (2)	0.041 (2)	0.0050 (18)	0.0151 (19)	0.0080 (17)
C16	0.064 (3)	0.032 (2)	0.048 (2)	0.0099 (18)	0.016 (2)	0.0126 (18)
C17	0.060 (2)	0.030 (2)	0.043 (2)	0.0088 (17)	0.0143 (18)	-0.0001 (17)
C18	0.063 (3)	0.032 (2)	0.032 (2)	0.0039 (17)	0.0154 (18)	0.0047 (17)
C19	0.052 (2)	0.033 (2)	0.0301 (19)	0.0079 (16)	0.0102 (16)	0.0056 (16)
C20	0.056 (3)	0.053 (3)	0.039 (2)	-0.001 (2)	0.0074 (19)	0.0115 (19)
C21	0.074 (3)	0.087 (4)	0.044 (3)	0.006 (3)	0.023 (2)	0.024 (3)
C22	0.093 (4)	0.090 (4)	0.033 (2)	0.012 (3)	0.014 (3)	0.005 (3)
C23	0.079 (3)	0.072 (3)	0.046 (3)	0.004 (3)	-0.008 (2)	0.001 (2)
C24	0.064 (3)	0.053 (3)	0.042 (2)	-0.004 (2)	0.009 (2)	0.009 (2)

Geometric parameters (Å, °)

S1—C1	1.771 (4)	C9—C10	1.360 (7)
S1—C7	1.773 (4)	C9—H9	0.9300
S2—C13	1.769 (4)	C10—C11	1.365 (7)
S2—C19	1.776 (4)	C10—H10	0.9300
O1—C6	1.228 (4)	C11—C12	1.365 (6)
O2—C6	1.306 (4)	C11—H11	0.9300
O2—H2	0.8200	C12—H12	0.9300
O3—C18	1.218 (4)	C13—C14	1.411 (5)
O4—C18	1.308 (4)	C14—C15	1.379 (5)
O4—H4O	0.8200	C14—C18	1.478 (5)
N1—C5	1.315 (5)	C15—C16	1.367 (5)
N1—C1	1.334 (4)	C15—H15	0.9300
N2—C17	1.315 (5)	C16—C17	1.369 (5)
N2—C13	1.333 (4)	C16—H16	0.9300
C1—C2	1.409 (5)	C17—H17	0.9300
C2—C3	1.374 (5)	C19—C20	1.376 (5)
C2—C6	1.472 (5)	C19—C24	1.386 (5)
C3—C4	1.374 (5)	C20—C21	1.386 (5)
C3—H3	0.9300	C20—H20	0.9300
C4—C5	1.372 (5)	C21—C22	1.354 (7)
C4—H4	0.9300	C21—H21	0.9300
C5—H5	0.9300	C22—C23	1.366 (7)
C7—C8	1.372 (5)	C22—H22	0.9300
C7—C12	1.387 (5)	C23—C24	1.371 (6)
C8—C9	1.397 (5)	C23—H23	0.9300
C8—H8	0.9300	C24—H24	0.9300
C1—S1—C7	103.18 (16)	C11—C12—C7	119.9 (4)
C13—S2—C19	103.20 (16)	C11—C12—H12	120.1
C6—O2—H2	109.5	C7—C12—H12	120.1
C18—O4—H4O	109.5	N2—C13—C14	121.1 (3)
C5—N1—C1	118.8 (3)	N2—C13—S2	117.3 (3)

supplementary materials

C17—N2—C13	118.6 (3)	C14—C13—S2	121.6 (3)
N1—C1—C2	121.4 (3)	C15—C14—C13	117.9 (3)
N1—C1—S1	116.8 (3)	C15—C14—C18	119.8 (3)
C2—C1—S1	121.8 (3)	C13—C14—C18	122.3 (3)
C3—C2—C1	117.6 (3)	C16—C15—C14	120.5 (4)
C3—C2—C6	119.1 (3)	C16—C15—H15	119.8
C1—C2—C6	123.1 (3)	C14—C15—H15	119.8
C2—C3—C4	120.7 (4)	C15—C16—C17	117.2 (3)
C2—C3—H3	119.6	C15—C16—H16	121.4
C4—C3—H3	119.6	C17—C16—H16	121.4
C5—C4—C3	117.1 (4)	N2—C17—C16	124.7 (3)
C5—C4—H4	121.5	N2—C17—H17	117.6
C3—C4—H4	121.5	C16—C17—H17	117.6
N1—C5—C4	124.3 (3)	O3—C18—O4	122.1 (3)
N1—C5—H5	117.8	O3—C18—C14	123.5 (3)
C4—C5—H5	117.8	O4—C18—C14	114.3 (3)
O1—C6—O2	121.9 (3)	C20—C19—C24	119.4 (4)
O1—C6—C2	123.8 (3)	C20—C19—S2	117.8 (3)
O2—C6—C2	114.2 (3)	C24—C19—S2	122.5 (3)
C8—C7—C12	119.6 (4)	C19—C20—C21	120.0 (4)
C8—C7—S1	117.3 (3)	C19—C20—H20	120.0
C12—C7—S1	122.9 (3)	C21—C20—H20	120.0
C7—C8—C9	119.3 (4)	C22—C21—C20	120.4 (4)
C7—C8—H8	120.3	C22—C21—H21	119.8
C9—C8—H8	120.3	C20—C21—H21	119.8
C10—C9—C8	120.6 (5)	C21—C22—C23	119.7 (4)
C10—C9—H9	119.7	C21—C22—H22	120.2
C8—C9—H9	119.7	C23—C22—H22	120.2
C9—C10—C11	119.6 (4)	C22—C23—C24	121.3 (4)
C9—C10—H10	120.2	C22—C23—H23	119.3
C11—C10—H10	120.2	C24—C23—H23	119.3
C10—C11—C12	121.1 (4)	C23—C24—C19	119.3 (4)
C10—C11—H11	119.5	C23—C24—H24	120.4
C12—C11—H11	119.5	C19—C24—H24	120.4
C5—N1—C1—C2	2.8 (5)	C17—N2—C13—C14	0.1 (5)
C5—N1—C1—S1	-177.9 (3)	C17—N2—C13—S2	178.6 (3)
C7—S1—C1—N1	-7.2 (3)	C19—S2—C13—N2	-8.3 (3)
C7—S1—C1—C2	172.1 (3)	C19—S2—C13—C14	170.2 (3)
N1—C1—C2—C3	-2.9 (6)	N2—C13—C14—C15	0.4 (6)
S1—C1—C2—C3	177.8 (3)	S2—C13—C14—C15	-177.9 (3)
N1—C1—C2—C6	173.5 (3)	N2—C13—C14—C18	-179.6 (3)
S1—C1—C2—C6	-5.8 (5)	S2—C13—C14—C18	2.0 (5)
C1—C2—C3—C4	0.2 (6)	C13—C14—C15—C16	0.0 (6)
C6—C2—C3—C4	-176.3 (4)	C18—C14—C15—C16	-179.9 (4)
C2—C3—C4—C5	2.4 (6)	C14—C15—C16—C17	-1.0 (6)
C1—N1—C5—C4	0.1 (6)	C13—N2—C17—C16	-1.3 (6)
C3—C4—C5—N1	-2.6 (7)	C15—C16—C17—N2	1.7 (6)
C3—C2—C6—O1	-171.7 (4)	C15—C14—C18—O3	-166.9 (4)
C1—C2—C6—O1	12.0 (6)	C13—C14—C18—O3	13.2 (6)

C3—C2—C6—O2	8.6 (6)	C15—C14—C18—O4	11.6 (6)
C1—C2—C6—O2	-167.8 (4)	C13—C14—C18—O4	-168.3 (4)
C1—S1—C7—C8	-122.4 (3)	C13—S2—C19—C20	126.3 (3)
C1—S1—C7—C12	62.3 (4)	C13—S2—C19—C24	-59.6 (4)
C12—C7—C8—C9	-0.2 (6)	C24—C19—C20—C21	0.3 (6)
S1—C7—C8—C9	-175.7 (3)	S2—C19—C20—C21	174.6 (3)
C7—C8—C9—C10	-0.3 (7)	C19—C20—C21—C22	-0.5 (7)
C8—C9—C10—C11	0.3 (8)	C20—C21—C22—C23	0.7 (8)
C9—C10—C11—C12	0.3 (8)	C21—C22—C23—C24	-0.8 (8)
C10—C11—C12—C7	-0.8 (7)	C22—C23—C24—C19	0.7 (7)
C8—C7—C12—C11	0.8 (6)	C20—C19—C24—C23	-0.4 (6)
S1—C7—C12—C11	176.0 (3)	S2—C19—C24—C23	-174.4 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H2···O3 ⁱ	0.82	1.82	2.624 (2)	168 (1)
O4—H4 ₀ ···O1 ⁱ	0.820	1.83000	2.642 (2)	169.62 (19)
C3—H3···O4 ⁱⁱ	0.93	2.50	3.264 (5)	139
C4—H4···O1 ⁱⁱⁱ	0.93	2.55	3.458 (5)	164
C15—H15···O2 ⁱⁱ	0.93	2.54	3.294 (5)	138
C16—H16···O3 ⁱⁱⁱ	0.93	2.58	3.467 (5)	160

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+1, -y+1, -z$; (iii) $x, y+1, z$.

supplementary materials

Fig. 1




Fig. 2

