

1-(6-Chloro-2-methyl-4-phenyl-3-quinolyl)ethanone

Hoong-Kun Fun,^{a,*‡} Wan-Sin Loh,^a S. Sarveswari,^b V. Vijayakumar^b and B. Palakshi Reddy^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bOrganic Chemistry Division, School of Science, VIT University, Vellore 632 014, India
Correspondence e-mail: hkfun@usm.my


Received 30 September 2009; accepted 2 October 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.036; wR factor = 0.107; data-to-parameter ratio = 39.7.

In the title compound, $\text{C}_{18}\text{H}_{14}\text{ClNO}$, the quinoline ring system is approximately planar with a maximum deviation of $0.022(1)\text{ \AA}$ and forms a dihedral angle of $62.70(3)^\circ$ with the phenyl ring. In the crystal, pairs of $\text{C}-\text{H} \cdots \text{O}$ intermolecular hydrogen bonds link neighbouring molecules into inversion dimers, forming $R_2^2(14)$ ring motifs. These inversion dimers are stacked along the b axis. The structure is further stabilized by $\text{C}-\text{H} \cdots \pi$ interactions.

Related literature

For reference bond-length data, see: Allen *et al.* (1987). For background to quinolines, see: Morimoto *et al.* (1991); Michael (1997); Markees *et al.* (1970); Campbell *et al.* (1988); Maguire *et al.* (1994); Kalluraya & Sreenivasa (1998); Roma *et al.* (2000); Chen *et al.* (2001); Skraup (1880); Katritzky & Arend (1998); Jiang & Si (2002). For the biological activity of chalcones, see: Dimmock *et al.* (1999); Yamazaki *et al.* (2002). For a related structure, see: Fun *et al.* (2009). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



‡ Thomson Reuters ResearcherID: A-3561-2009.

Experimental

Crystal data

$\text{C}_{18}\text{H}_{14}\text{ClNO}$
 $M_r = 295.75$
Monoclinic, $P2_1/n$
 $a = 10.4633(2)\text{ \AA}$
 $b = 7.7959(1)\text{ \AA}$
 $c = 17.5925(3)\text{ \AA}$
 $\beta = 90.887(1)^\circ$

$V = 1434.86(4)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 0.26\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.57 \times 0.34 \times 0.27\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.865$, $T_{\max} = 0.932$

32340 measured reflections
7613 independent reflections
6588 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.107$
 $S = 1.07$
7613 reflections

192 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.58\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
C15—H15A…O1 ⁱ	0.93	2.55	3.2047 (10)	128
C11—H11A…Cg1 ⁱⁱ	0.93	2.78	3.6416 (7)	155
C13—H13A…Cg2 ⁱⁱⁱ	0.93	2.92	3.6255 (8)	133

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$. Cg1 and Cg2 are the centroids of the Cl1-C9/N1 and C10-C15 ring systems, respectively.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL* software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and WSL thank USM for the Research University Golden Goose Grant (No. 1001/PFIZIK/811012). WSL thanks the Malaysian Government and USM for the award of the post of Assistant Research Officer under the Research University Golden Goose Grant (No. 1001/PFIZIK/811012). VV is grateful to DST-India for funding through the Young Scientist Scheme (Fast Track Proposal).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: WN2352).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Campbell, S. F., Hardstone, J. D. & Palmer, M. J. (1988). *J. Med. Chem.* **31**, 1031–1035.
Chen, Y.-L., Fang, K.-C., Sheu, J.-Y., Hsu, S.-L. & Tzeng, C.-C. (2001). *J. Med. Chem.* **44**, 2374–2377.

- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Dimmock, J. R., Elias, D. W., Beazely, M. A. & Kandepu, N. M. (1999). *Curr. Med. Chem.* **6**, 1125–1149.
- Fun, H.-K., Yeap, C. S., Sarveswari, S., Vijayakumar, V. & Prasath, R. (2009). *Acta Cryst. E* **65**, o2665–o2666.
- Jiang, B. & Si, Y.-G. (2002). *J. Org. Chem.* **67**, 9449–9451.
- Kalluraya, B. & Sreenivasa, S. (1998). *Farmaco*, **53**, 399–404.
- Katritzky, A. R. & Arend, M. I. (1998). *J. Org. Chem.* **63**, 9989–9991.
- Maguire, M. P., Sheets, K. R., McVety, K., Spada, A. P. & Zilberstein, A. (1994). *J. Med. Chem.* **37**, 2129–2137.
- Markees, D. G., Dewey, V. C. & Kidder, G. W. (1970). *J. Med. Chem.* **13**, 324–326.
- Michael, J. P. (1997). *Nat. Prod. Rep.* **14**, 605–608.
- Morimoto, Y., Matsuda, F. & Shirahama, H. (1991). *Synlett*, **3**, 202–203.
- Roma, G., Braccio, M. D., Grossi, G., Mattioli, F. & Ghia, M. (2000). *Eur. J. Med. Chem.* **35**, 1021–1026.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Skraup, H. (1880). *Ber. Dtsch. Chem. Ges.* **13**, 2086–2088.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Yamazaki, S., Morita, T. & Endo, H. (2002). *Cancer Lett.* **183**, 23–30.

supplementary materials

Acta Cryst. (2009). E65, o2688–o2689 [doi:10.1107/S1600536809040306]

1-(6-Chloro-2-methyl-4-phenyl-3-quinolyl)ethanone

H.-K. Fun, W.-S. Loh, S. Sarveswari, V. Vijayakumar and B. P. Reddy

Comment

Quinolines and their derivatives are very important compounds because of their wide occurrence in natural products (Morimoto *et al.*, 1991; Michael, 1997) and biologically active compounds (Markees *et al.*, 1970; Campbell *et al.*, 1988). A large variety of quinolines have interesting physiological activities and have found attractive applications as pharmaceuticals, agrochemicals and as synthetic building blocks (Maguire *et al.*, 1994; Kalluraya & Sreenivasa, 1998; Roma *et al.*, 2000; Chen *et al.*, 2001; Skraup, 1880). Many synthetic methods such as the Skraup, Doeblner-Von Miller, Friedländer and Combes reactions have been developed for the preparation of quinolines, but due to their great importance, the synthesis of new derivatives of quinoline remains an active research area (Katritzky & Arend, 1998; Jiang & Si, 2002). Chalcones are open-chain flavonoids, possessing a variety of biological activities, including antioxidant, anti-inflammatory, antimicrobial, antiprotozoal, antiulcer, as well as other activities (Dimmock *et al.*, 1999). More importantly, chalcones have shown several anticancer activities, such as inhibitors of cancer cell proliferation, carcinogenesis, and metastasis (Yamazaki *et al.*, 2002).

In the crystal structure (Fig. 1), bond lengths (Allen *et al.*, 1987) and angles are within normal ranges and comparable to those in a closely related structure (Fun *et al.*, 2009). The quinoline ring system (C1–C9/N1) is approximately planar, with a maximum deviation of 0.022 (1) Å at atom C1. The phenyl ring (C10–C15) forms a dihedral angle of 62.70 (3)° with the mean plane of the quinoline ring system. In the crystal packing (Fig. 2), pairs of C15—H15A…O1 hydrogen bonds link neighbouring molecules into dimers, forming $R_2^2(14)$ ring motifs (Bernstein *et al.*, 1995). These inversion dimers are stacked along the *b* axis. The crystal structure is further stabilized by C—H…π interactions (Table 1), involving the C1–C9/N1 (centroid *Cg1*) and C10–C15 (centroid *Cg2*) ring systems.

Experimental

A mixture of 2-amino-5-chlorobenzophenone (2.3 g, 0.01 mol) and acetylacetone (1.0 g, 0.01 mol) with 0.15 ml concentrated HCl in a beaker was subjected to microwave irradiation for about 6 min. After completion of the reaction (monitored by TLC), the reaction mixture was washed with saturated solvent NaHCO₃ (10 ml) and then it was dried. After that it was washed with petroleum ether and recrystallized with chloroform (*M. p.* 224–226°C). IR (cm⁻¹): 1704, 1480, 1385, 840, 711.

Refinement

All H atoms were positioned geometrically [C—H = 0.93 or 0.96 Å] and were refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{Csp}^2)$ or $1.5U_{\text{eq}}(\text{methyl C})$. A rotating-group model was applied for the methyl groups.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. The crystal packing of the title compound, viewed along the b axis, showing the $R_2^2(14)$ ring motifs. C—H···O intermolecular interactions are shown as dashed lines.

1-(6-Chloro-2-methyl-4-phenyl-3-quinolyl)ethanone

Crystal data

$C_{18}H_{14}ClNO$	$F_{000} = 616$
$M_r = 295.75$	$D_x = 1.369 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 9994 reflections
$a = 10.4633 (2) \text{ \AA}$	$\theta = 2.3\text{--}37.6^\circ$
$b = 7.7959 (1) \text{ \AA}$	$\mu = 0.26 \text{ mm}^{-1}$
$c = 17.5925 (3) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 90.887 (1)^\circ$	Block, yellow
$V = 1434.86 (4) \text{ \AA}^3$	$0.57 \times 0.34 \times 0.27 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	7613 independent reflections
Radiation source: fine-focus sealed tube	6588 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.023$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 37.6^\circ$
φ and ω scans	$\theta_{\text{min}} = 2.3^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -17 \rightarrow 16$
$T_{\text{min}} = 0.865$, $T_{\text{max}} = 0.932$	$k = -13 \rightarrow 12$
32340 measured reflections	$l = -30 \rightarrow 30$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.036$	H-atom parameters constrained
$wR(F^2) = 0.107$	$w = 1/[\sigma^2(F_o^2) + (0.0574P)^2 + 0.2858P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.07$	$(\Delta/\sigma)_{\max} < 0.001$
7613 reflections	$\Delta\rho_{\max} = 0.58 \text{ e \AA}^{-3}$
192 parameters	$\Delta\rho_{\min} = -0.24 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.179322 (16)	0.65521 (3)	0.247848 (10)	0.02152 (5)
O1	0.56817 (6)	0.27543 (9)	0.51440 (4)	0.02585 (12)
N1	0.98196 (6)	0.23575 (8)	0.49910 (3)	0.01532 (10)
C1	0.98802 (6)	0.50843 (9)	0.32517 (4)	0.01414 (10)
H1A	0.9336	0.5589	0.2894	0.017*
C2	1.11760 (6)	0.53229 (9)	0.32124 (4)	0.01548 (11)
C3	1.20321 (6)	0.46250 (9)	0.37595 (4)	0.01719 (11)
H3A	1.2907	0.4809	0.3723	0.021*
C4	1.15518 (6)	0.36706 (9)	0.43465 (4)	0.01614 (11)
H4A	1.2107	0.3226	0.4714	0.019*
C5	1.02224 (6)	0.33535 (8)	0.43995 (4)	0.01347 (10)
C6	0.93765 (6)	0.40614 (8)	0.38437 (3)	0.01243 (10)
C7	0.80451 (6)	0.36943 (8)	0.39077 (3)	0.01238 (10)
C8	0.76613 (6)	0.26872 (8)	0.45085 (3)	0.01334 (10)
C9	0.85868 (6)	0.20413 (9)	0.50445 (4)	0.01465 (10)
C10	0.70864 (6)	0.44156 (8)	0.33610 (3)	0.01260 (10)

supplementary materials

C11	0.70861 (6)	0.39520 (9)	0.25902 (4)	0.01491 (10)
H11A	0.7714	0.3219	0.2410	0.018*
C12	0.61476 (6)	0.45857 (9)	0.20938 (4)	0.01612 (11)
H12A	0.6143	0.4263	0.1585	0.019*
C13	0.52150 (6)	0.57038 (9)	0.23602 (4)	0.01573 (11)
H13A	0.4589	0.6126	0.2029	0.019*
C14	0.52217 (6)	0.61880 (9)	0.31229 (4)	0.01545 (11)
H14A	0.4607	0.6947	0.3298	0.019*
C15	0.61476 (6)	0.55367 (9)	0.36233 (4)	0.01429 (10)
H15A	0.6141	0.5849	0.4133	0.017*
C16	0.81736 (8)	0.09262 (10)	0.56922 (4)	0.02031 (13)
H16B	0.8914	0.0471	0.5951	0.030*
H16C	0.7660	-0.0001	0.5499	0.030*
H16A	0.7683	0.1595	0.6041	0.030*
C17	0.62761 (6)	0.22144 (9)	0.46068 (4)	0.01645 (11)
C18	0.56993 (9)	0.09744 (14)	0.40461 (5)	0.02852 (17)
H18A	0.4785	0.1000	0.4083	0.043*
H18B	0.6004	-0.0162	0.4156	0.043*
H18C	0.5940	0.1292	0.3541	0.043*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.01532 (7)	0.02794 (10)	0.02141 (8)	-0.00307 (5)	0.00332 (5)	0.00537 (6)
O1	0.0226 (3)	0.0312 (3)	0.0242 (3)	0.0004 (2)	0.0103 (2)	-0.0010 (2)
N1	0.0162 (2)	0.0145 (2)	0.0153 (2)	0.00089 (17)	-0.00179 (17)	0.00089 (17)
C1	0.0122 (2)	0.0158 (2)	0.0145 (2)	0.00001 (19)	0.00039 (18)	0.00054 (19)
C2	0.0132 (2)	0.0166 (3)	0.0167 (2)	-0.00137 (19)	0.00138 (19)	0.0000 (2)
C3	0.0123 (2)	0.0174 (3)	0.0219 (3)	-0.0004 (2)	-0.0010 (2)	-0.0008 (2)
C4	0.0133 (2)	0.0153 (3)	0.0197 (3)	0.00069 (19)	-0.0032 (2)	-0.0008 (2)
C5	0.0136 (2)	0.0125 (2)	0.0143 (2)	0.00098 (18)	-0.00177 (18)	-0.00102 (18)
C6	0.0118 (2)	0.0130 (2)	0.0125 (2)	0.00046 (18)	-0.00022 (17)	-0.00075 (18)
C7	0.0121 (2)	0.0135 (2)	0.0115 (2)	0.00065 (17)	0.00041 (17)	-0.00052 (18)
C8	0.0136 (2)	0.0143 (2)	0.0122 (2)	0.00014 (18)	0.00090 (18)	-0.00009 (18)
C9	0.0168 (2)	0.0135 (2)	0.0136 (2)	0.00065 (19)	-0.00055 (19)	0.00067 (19)
C10	0.0108 (2)	0.0151 (2)	0.0119 (2)	-0.00022 (18)	0.00032 (17)	0.00094 (18)
C11	0.0140 (2)	0.0181 (3)	0.0127 (2)	0.0014 (2)	0.00037 (18)	-0.0009 (2)
C12	0.0151 (2)	0.0200 (3)	0.0132 (2)	-0.0002 (2)	-0.00113 (19)	0.0002 (2)
C13	0.0127 (2)	0.0181 (3)	0.0164 (2)	-0.00090 (19)	-0.00199 (19)	0.0024 (2)
C14	0.0116 (2)	0.0171 (3)	0.0176 (3)	0.00086 (19)	0.00059 (19)	0.0011 (2)
C15	0.0122 (2)	0.0171 (3)	0.0136 (2)	0.00082 (19)	0.00122 (18)	-0.00023 (19)
C16	0.0232 (3)	0.0199 (3)	0.0177 (3)	0.0000 (2)	0.0000 (2)	0.0065 (2)
C17	0.0150 (2)	0.0194 (3)	0.0150 (2)	-0.0010 (2)	0.00195 (19)	0.0031 (2)
C18	0.0256 (4)	0.0377 (5)	0.0222 (3)	-0.0151 (3)	0.0014 (3)	-0.0032 (3)

Geometric parameters (\AA , $^\circ$)

Cl1—C2	1.7401 (7)	C10—C15	1.3985 (9)
O1—C17	1.2146 (9)	C10—C11	1.4032 (9)

N1—C9	1.3181 (9)	C11—C12	1.3944 (9)
N1—C5	1.3701 (9)	C11—H11A	0.9300
C1—C2	1.3714 (9)	C12—C13	1.3949 (10)
C1—C6	1.4196 (9)	C12—H12A	0.9300
C1—H1A	0.9300	C13—C14	1.3937 (10)
C2—C3	1.4137 (10)	C13—H13A	0.9300
C3—C4	1.3742 (10)	C14—C15	1.3947 (9)
C3—H3A	0.9300	C14—H14A	0.9300
C4—C5	1.4173 (9)	C15—H15A	0.9300
C4—H4A	0.9300	C16—H16B	0.9600
C5—C6	1.4203 (9)	C16—H16C	0.9600
C6—C7	1.4284 (9)	C16—H16A	0.9600
C7—C8	1.3813 (9)	C17—C18	1.5012 (11)
C7—C10	1.4891 (9)	C18—H18A	0.9600
C8—C9	1.4325 (9)	C18—H18B	0.9600
C8—C17	1.5081 (9)	C18—H18C	0.9600
C9—C16	1.5021 (10)		
C9—N1—C5	118.23 (6)	C12—C11—C10	120.24 (6)
C2—C1—C6	119.46 (6)	C12—C11—H11A	119.9
C2—C1—H1A	120.3	C10—C11—H11A	119.9
C6—C1—H1A	120.3	C11—C12—C13	120.03 (6)
C1—C2—C3	122.01 (6)	C11—C12—H12A	120.0
C1—C2—Cl1	119.39 (5)	C13—C12—H12A	120.0
C3—C2—Cl1	118.59 (5)	C14—C13—C12	119.97 (6)
C4—C3—C2	119.00 (6)	C14—C13—H13A	120.0
C4—C3—H3A	120.5	C12—C13—H13A	120.0
C2—C3—H3A	120.5	C13—C14—C15	120.14 (6)
C3—C4—C5	120.97 (6)	C13—C14—H14A	119.9
C3—C4—H4A	119.5	C15—C14—H14A	119.9
C5—C4—H4A	119.5	C14—C15—C10	120.25 (6)
N1—C5—C4	117.56 (6)	C14—C15—H15A	119.9
N1—C5—C6	123.19 (6)	C10—C15—H15A	119.9
C4—C5—C6	119.25 (6)	C9—C16—H16B	109.5
C1—C6—C5	119.27 (6)	C9—C16—H16C	109.5
C1—C6—C7	122.95 (6)	H16B—C16—H16C	109.5
C5—C6—C7	117.78 (6)	C9—C16—H16A	109.5
C8—C7—C6	118.01 (6)	H16B—C16—H16A	109.5
C8—C7—C10	120.53 (5)	H16C—C16—H16A	109.5
C6—C7—C10	121.43 (5)	O1—C17—C18	121.94 (7)
C7—C8—C9	120.13 (6)	O1—C17—C8	120.60 (7)
C7—C8—C17	121.19 (6)	C18—C17—C8	117.34 (6)
C9—C8—C17	118.67 (6)	C17—C18—H18A	109.5
N1—C9—C8	122.66 (6)	C17—C18—H18B	109.5
N1—C9—C16	117.12 (6)	H18A—C18—H18B	109.5
C8—C9—C16	120.21 (6)	C17—C18—H18C	109.5
C15—C10—C11	119.35 (6)	H18A—C18—H18C	109.5
C15—C10—C7	119.49 (5)	H18B—C18—H18C	109.5
C11—C10—C7	121.13 (6)		

supplementary materials


C6—C1—C2—C3	−2.09 (10)	C5—N1—C9—C8	0.38 (10)
C6—C1—C2—Cl1	179.09 (5)	C5—N1—C9—C16	179.18 (6)
C1—C2—C3—C4	0.36 (11)	C7—C8—C9—N1	−0.40 (10)
Cl1—C2—C3—C4	179.19 (5)	C17—C8—C9—N1	178.37 (6)
C2—C3—C4—C5	1.27 (10)	C7—C8—C9—C16	−179.17 (6)
C9—N1—C5—C4	−179.52 (6)	C17—C8—C9—C16	−0.40 (9)
C9—N1—C5—C6	0.11 (10)	C8—C7—C10—C15	61.18 (9)
C3—C4—C5—N1	178.51 (6)	C6—C7—C10—C15	−116.81 (7)
C3—C4—C5—C6	−1.13 (10)	C8—C7—C10—C11	−116.99 (7)
C2—C1—C6—C5	2.18 (10)	C6—C7—C10—C11	65.01 (9)
C2—C1—C6—C7	−177.46 (6)	C15—C10—C11—C12	−0.88 (10)
N1—C5—C6—C1	179.78 (6)	C7—C10—C11—C12	177.30 (6)
C4—C5—C6—C1	−0.60 (9)	C10—C11—C12—C13	0.91 (10)
N1—C5—C6—C7	−0.57 (9)	C11—C12—C13—C14	0.02 (10)
C4—C5—C6—C7	179.06 (6)	C12—C13—C14—C15	−0.97 (10)
C1—C6—C7—C8	−179.83 (6)	C13—C14—C15—C10	1.00 (10)
C5—C6—C7—C8	0.52 (9)	C11—C10—C15—C14	−0.07 (10)
C1—C6—C7—C10	−1.79 (10)	C7—C10—C15—C14	−178.28 (6)
C5—C6—C7—C10	178.56 (6)	C7—C8—C17—O1	−113.46 (8)
C6—C7—C8—C9	−0.08 (9)	C9—C8—C17—O1	67.78 (9)
C10—C7—C8—C9	−178.14 (6)	C7—C8—C17—C18	70.31 (9)
C6—C7—C8—C17	−178.81 (6)	C9—C8—C17—C18	−108.45 (8)
C10—C7—C8—C17	3.13 (9)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C15—H15A···O1 ⁱ	0.93	2.55	3.2047 (10)	128
C11—H11A···Cg1 ⁱⁱ	0.93	2.78	3.6416 (7)	155
C13—H13A···Cg2 ⁱⁱⁱ	0.93	2.92	3.6255 (8)	133

Symmetry codes: (i) $-x+1, -y+1, -z+1$; (ii) $-x+3/2, y-1/2, -z+1/2$; (iii) $-x+3/2, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

