

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Chloro-7,8-dimethylquinoline-3-carbaldehyde

F. Nawaz Khan,^a R. Subashini,^a Atul Kumar Kushwaha,^a Venkatesha R. Hathwar^b and Seik Weng Ng^{c*}

^aChemistry Division, School of Science and Humanities, VIT University, Vellore 632 014, Tamil Nadu, India, ^bSolid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560 012, Karnataka, India, and ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
Correspondence e-mail: seikweng@um.edu.my


Received 6 October 2009; accepted 6 October 2009

Key indicators: single-crystal X-ray study; $T = 290$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.034; wR factor = 0.115; data-to-parameter ratio = 13.8.

All the non-H atoms of the title compound, $\text{C}_{12}\text{H}_{10}\text{ClNO}$, lie on a crystallographic mirror plane orientated perpendicular to the crystallographic b axis.

Related literature

For a review of the synthesis of quinolines by the Vilsmeier–Haack reaction, see: Meth-Cohn (1993).



Experimental

Crystal data

 $\text{C}_{12}\text{H}_{10}\text{ClNO}$ $M_r = 219.66$ Orthorhombic, $Pnma$ $a = 20.4542$ (13) Å $b = 6.7393$ (4) Å $c = 7.5675$ (4) Å $V = 1043.16$ (11) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.34$ mm⁻¹ $T = 290$ K $0.28 \times 0.21 \times 0.17$ mm

Data collection

Bruker SMART area-detector diffractometer

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.912$, $T_{\max} = 0.945$

7145 measured reflections

1299 independent reflections

1078 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.022$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.034$ $wR(F^2) = 0.115$ $S = 1.07$

1299 reflections

94 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.29$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.30$ e Å⁻³

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINTE* (Bruker, 2004); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Department of Science and Technology, India, for use of the diffraction facility at IISc under the IRHPA–DST program; FNK thanks the DST for Fast Track Proposal funding. We thank VIT University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5086).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Bruker (2004). *SAINTE* and *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Meth-Cohn, O. (1993). *Heterocycles*, **35**, 539–557.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, o2709 [doi:10.1107/S1600536809040860]

2-Chloro-7,8-dimethylquinoline-3-carbaldehyde

F. N. Khan, R. Subashini, A. K. Kushwaha, V. R. Hathwar and S. W. Ng

Experimental

A Vilsmeier-Haack adduct prepared from phosphorus oxytrichloride (6.5 ml, 70 mmol) and *N,N*-dimethylformamide (2.3 ml, 30 mmol) at 273 K was added to *N*-(2,3-dimethylphenyl)acetamide (1.63 g, 10 mmol). The mixture was heated at 353 K for 15 h. The mixture was poured onto ice; the white product was collected and dried. The compound was purified by recrystallization from a petroleum ether/ethyl acetate mixture.

Refinement

H-atoms were placed in calculated positions (C–H 0.93–0.96 Å) and were included in the refinement in the riding model approximation, with $U(\text{H})$ set to 1.2–1.5 $U(\text{C})$.

Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of $\text{C}_{12}\text{H}_{10}\text{ClNO}$ at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

2-Chloro-7,8-dimethylquinoline-3-carbaldehyde

Crystal data

$\text{C}_{12}\text{H}_{10}\text{ClNO}$

$M_r = 219.66$

Orthorhombic, *Pnma*

Hall symbol: -P 2ac 2n

$a = 20.4542$ (13) Å

$b = 6.7393$ (4) Å

$c = 7.5675$ (4) Å

$V = 1043.16$ (11) Å³

$Z = 4$

$F_{000} = 456$

$D_x = 1.399$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 753 reflections

$\theta = 2.0$ – 24.4°

$\mu = 0.34$ mm⁻¹

$T = 290$ K

Block, colorless

$0.28 \times 0.21 \times 0.17$ mm

Data collection

Bruker SMART area-detector
diffractometer

1299 independent reflections

Radiation source: fine-focus sealed tube

1078 reflections with $I > 2\sigma(I)$

supplementary materials

Monochromator: graphite
 $T = 290$ K
 φ and ω scans
 Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.912$, $T_{\max} = 0.945$
 7145 measured reflections

$R_{\text{int}} = 0.022$
 $\theta_{\max} = 27.5^\circ$
 $\theta_{\min} = 2.0^\circ$
 $h = -25 \rightarrow 26$
 $k = -8 \rightarrow 4$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.115$
 $S = 1.07$
 1299 reflections
 94 parameters
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0706P)^2 + 0.1712P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{\AA}^{-3}$
 Extinction correction: SHELXL97 (Sheldrick, 2008),
 $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.010 (2)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C11	0.59400 (3)	0.2500	0.07517 (7)	0.0568 (2)	
N1	0.48555 (8)	0.2500	0.24995 (19)	0.0363 (4)	
O1	0.68317 (9)	0.2500	0.5872 (3)	0.0672 (5)	
C1	0.54815 (9)	0.2500	0.2711 (2)	0.0369 (4)	
C2	0.58228 (10)	0.2500	0.4343 (3)	0.0385 (4)	
C3	0.54382 (10)	0.2500	0.5832 (3)	0.0382 (4)	
H3	0.5635	0.2500	0.6939	0.046*	
C4	0.47569 (10)	0.2500	0.5712 (2)	0.0347 (4)	
C5	0.43407 (11)	0.2500	0.7208 (3)	0.0428 (5)	
H5	0.4517	0.2500	0.8341	0.051*	
C6	0.36779 (11)	0.2500	0.6971 (3)	0.0461 (5)	
H6	0.3408	0.2500	0.7960	0.055*	
C7	0.33908 (10)	0.2500	0.5279 (3)	0.0411 (5)	
C8	0.37829 (9)	0.2500	0.3777 (3)	0.0361 (4)	
C9	0.44701 (9)	0.2500	0.3993 (2)	0.0322 (4)	
C10	0.65428 (11)	0.2500	0.4503 (3)	0.0531 (6)	
H10	0.6787	0.2500	0.3467	0.064*	
C11	0.26571 (11)	0.2500	0.5126 (4)	0.0583 (6)	
H11A	0.2468	0.2766	0.6262	0.087*	0.50
H11B	0.2512	0.1227	0.4713	0.087*	0.50
H11C	0.2524	0.3507	0.4305	0.087*	0.50

C12	0.34939 (11)	0.2500	0.1950 (3)	0.0532 (6)	
H12A	0.3794	0.1882	0.1144	0.080*	0.50
H12B	0.3414	0.3842	0.1583	0.080*	0.50
H12C	0.3090	0.1776	0.1957	0.080*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0424 (3)	0.0881 (5)	0.0400 (3)	0.000	0.0087 (2)	0.000
N1	0.0355 (8)	0.0437 (8)	0.0298 (7)	0.000	-0.0008 (6)	0.000
O1	0.0480 (10)	0.0820 (12)	0.0716 (12)	0.000	-0.0252 (9)	0.000
C1	0.0356 (10)	0.0415 (9)	0.0336 (9)	0.000	0.0018 (7)	0.000
C2	0.0344 (10)	0.0387 (9)	0.0423 (11)	0.000	-0.0058 (7)	0.000
C3	0.0417 (11)	0.0389 (9)	0.0341 (9)	0.000	-0.0097 (7)	0.000
C4	0.0411 (10)	0.0344 (9)	0.0286 (9)	0.000	-0.0022 (7)	0.000
C5	0.0497 (12)	0.0513 (11)	0.0275 (9)	0.000	0.0003 (8)	0.000
C6	0.0483 (12)	0.0556 (11)	0.0343 (10)	0.000	0.0078 (9)	0.000
C7	0.0357 (10)	0.0453 (10)	0.0423 (10)	0.000	0.0028 (8)	0.000
C8	0.0351 (10)	0.0386 (9)	0.0346 (9)	0.000	-0.0021 (7)	0.000
C9	0.0361 (9)	0.0324 (8)	0.0281 (8)	0.000	-0.0014 (7)	0.000
C10	0.0375 (11)	0.0623 (13)	0.0595 (14)	0.000	-0.0064 (10)	0.000
C11	0.0367 (11)	0.0783 (16)	0.0600 (15)	0.000	0.0085 (10)	0.000
C12	0.0395 (11)	0.0814 (15)	0.0387 (11)	0.000	-0.0054 (9)	0.000

Geometric parameters (\AA , $^\circ$)

C11—C1	1.7545 (19)	C6—C7	1.409 (3)
N1—C1	1.290 (2)	C6—H6	0.9300
N1—C9	1.378 (2)	C7—C8	1.391 (3)
O1—C10	1.192 (3)	C7—C11	1.505 (3)
C1—C2	1.419 (3)	C8—C9	1.415 (3)
C2—C3	1.374 (3)	C8—C12	1.504 (3)
C2—C10	1.478 (3)	C10—H10	0.9300
C3—C4	1.396 (3)	C11—H11A	0.9600
C3—H3	0.9300	C11—H11B	0.9600
C4—C5	1.416 (3)	C11—H11C	0.9600
C4—C9	1.427 (2)	C12—H12A	0.9600
C5—C6	1.367 (3)	C12—H12B	0.9600
C5—H5	0.9300	C12—H12C	0.9600
C1—N1—C9	117.76 (16)	C7—C8—C9	118.58 (17)
N1—C1—C2	126.61 (17)	C7—C8—C12	121.64 (18)
N1—C1—C11	115.19 (14)	C9—C8—C12	119.79 (17)
C2—C1—C11	118.20 (15)	N1—C9—C8	118.25 (15)
C3—C2—C1	115.60 (18)	N1—C9—C4	120.82 (16)
C3—C2—C10	120.21 (18)	C8—C9—C4	120.92 (16)
C1—C2—C10	124.20 (19)	O1—C10—C2	124.4 (2)
C2—C3—C4	121.19 (17)	O1—C10—H10	117.8
C2—C3—H3	119.4	C2—C10—H10	117.8

supplementary materials

C4—C3—H3	119.4	C7—C11—H11A	109.5
C3—C4—C5	123.21 (17)	C7—C11—H11B	109.5
C3—C4—C9	118.02 (16)	H11A—C11—H11B	109.5
C5—C4—C9	118.77 (18)	C7—C11—H11C	109.5
C6—C5—C4	119.43 (18)	H11A—C11—H11C	109.5
C6—C5—H5	120.3	H11B—C11—H11C	109.5
C4—C5—H5	120.3	C8—C12—H12A	109.5
C5—C6—C7	122.15 (18)	C8—C12—H12B	109.5
C5—C6—H6	118.9	H12A—C12—H12B	109.5
C7—C6—H6	118.9	C8—C12—H12C	109.5
C8—C7—C6	120.15 (19)	H12A—C12—H12C	109.5
C8—C7—C11	120.8 (2)	H12B—C12—H12C	109.5
C6—C7—C11	119.0 (2)		
C9—N1—C1—C2	0.0	C11—C7—C8—C9	180.0
C9—N1—C1—C11	180.0	C6—C7—C8—C12	180.0
N1—C1—C2—C3	0.0	C11—C7—C8—C12	0.0
C11—C1—C2—C3	180.0	C1—N1—C9—C8	180.0
N1—C1—C2—C10	180.0	C1—N1—C9—C4	0.0
C11—C1—C2—C10	0.0	C7—C8—C9—N1	180.0
C1—C2—C3—C4	0.0	C12—C8—C9—N1	0.0
C10—C2—C3—C4	180.0	C7—C8—C9—C4	0.0
C2—C3—C4—C5	180.0	C12—C8—C9—C4	180.0
C2—C3—C4—C9	0.0	C3—C4—C9—N1	0.0
C3—C4—C5—C6	180.0	C5—C4—C9—N1	180.0
C9—C4—C5—C6	0.0	C3—C4—C9—C8	180.0
C4—C5—C6—C7	0.0	C5—C4—C9—C8	0.0
C5—C6—C7—C8	0.0	C3—C2—C10—O1	0.0
C5—C6—C7—C11	180.0	C1—C2—C10—O1	180.0
C6—C7—C8—C9	0.0		

Fig. 1

