

1,3-Bis(hydroxymethyl)benzimidazolin-2-one

H. C. Devarajegowda,^{a*} V. Madhura,^b

B. S. Palakshamurthy,^a S. Jeyaseelan^a and Manohar V. Kulkarni^b

^aDepartment of Physics, Yuvaraja's College (Constituent College), University of Mysore, Mysore 570 005, Karnataka, India, and ^bDepartment of Chemistry, Karnataka University, Dharwad 580 003, Karnataka, India
Correspondence e-mail: devarajegowda@yahoo.com


Received 24 September 2009; accepted 7 October 2009

Key indicators: single-crystal X-ray study; $T = 273\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.040; wR factor = 0.110; data-to-parameter ratio = 14.1.

The title compound, $\text{C}_9\text{H}_{10}\text{N}_2\text{O}_3$, crystallizes with one and a half molecules in the asymmetric unit, one lying on a general position and the other on a twofold rotation axis. The dihedral angle between the two independent benzimidazole ring systems is $18.96(5)^\circ$. In the crystal, molecules are linked into a three-dimensional network by $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonding involving *N*-hydroxymethyl and carbonyl groups, and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For general background to 2-benzimidazolones, see: Raghu *et al.* (2005); Porret & Hebermeier (1974); Habermeier (1976); Trask-Morrel *et al.* (1988); Hammach *et al.* (2006); Bansal *et al.* (1981). For related structures, see: Anklekar & Kulkarni (1995); Schwiebert *et al.* (1996). For the synthesis, see: Zinner & Spangenberg (1958).



Experimental

Crystal data

$\text{C}_9\text{H}_{10}\text{N}_2\text{O}_3$
 $M_r = 194.19$
Monoclinic, $C2/c$
 $a = 13.5515(14)\text{ \AA}$
 $b = 11.0848(12)\text{ \AA}$

$c = 17.6253(19)\text{ \AA}$
 $\beta = 94.216(2)^\circ$
 $V = 2640.4(5)\text{ \AA}^3$
 $Z = 12$
Mo $K\alpha$ radiation

$\mu = 0.11\text{ mm}^{-1}$
 $T = 273\text{ K}$

$0.22 \times 0.20 \times 0.10\text{ mm}$

Data collection

Bruker SMART CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2004)
 $T_{\min} = 0.976$, $T_{\max} = 0.986$

13521 measured reflections
2684 independent reflections
2395 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.110$
 $S = 1.10$
2684 reflections

191 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.16\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.24\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1A \cdots O4	0.82	1.99	2.8003 (16)	169
O2—H2A \cdots O5 ⁱ	0.82	1.93	2.7503 (18)	176
O5—H5A \cdots O3 ⁱⁱ	0.82	1.84	2.6551 (17)	175
C3—H3 \cdots O2 ⁱⁱⁱ	0.93	2.58	3.489 (2)	164
C14—H14B \cdots O1 ⁱⁱ	0.97	2.54	3.364 (2)	143

Symmetry codes: (i) $x, -y + 1, z - \frac{1}{2}$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + \frac{3}{2}, -y + \frac{3}{2}, -z$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors thank Professor T. N. Guru Row and Miss Brinda Selvaraj, Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, for their help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2925).

References

- Anklekar, K. Y. & Kulkarni, M. V. (1995). *Indian J. Chem. Sect. B*, **34**, 677–678.
- Bansal, P. C., Ian, H., Pitman, I. H. & Higuchi, T. (1981). *J. Pharm. Sci.* **70**, 855–857.
- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Habermeier, J. (1976). *Ger. Offen.* 2 453 450.
- Hammach, A., Barbosa, A., Gaenzler, F. C., Fadra, T., Goldberg, D., Hao, M.-H., Kroe, R. R., Liu, P., Qian, K. C., Ralph, M., Sarko, C., Soleymanzadeh, F. & Moss, N. (2006). *Bioorg. Med. Chem. Lett.* **16**, 6316–6320.
- Porret, D. & Hebermeier, J. (1974). *Ger. Offen.* 2 342 432.
- Raghu, A. V., Gadaginamath, G. S. & Amminabavi, T. M. (2005). *J. Appl. Polym. Sci.* **98**, 2236–2244.
- Schwiebert, K. E., Chin, D. N., MacDonald, J. C. & Whitesides, G. M. (1996). *J. Am. Chem. Soc.* **118**, 4018–4029.
- Sheldrick, G. M. (2004). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Trask-Morrel, B. J., Franklin, W. E. & Liu, R. H. (1988). *Text. Chem. Color*, pp. 20–21.
- Zinner, H. & Spangenberg, B. (1958). *Chem. Ber.* **91**, 1432–1437.

supplementary materials

Acta Cryst. (2009). E65, o2735 [doi:10.1107/S1600536809040963]

1,3-Bis(hydroxymethyl)benzimidazolin-2-one

H. C. Devarajegowda, V. Madhura, B. S. Palakshamurthy, S. Jeyaseelan and M. V. Kulkarni

Comment

1,3-Bishydroxyalkylated benzimidazolones are an important class of functionalized benzimidazoles which have been found to be useful as polymer intermediates (Raghu *et al.*, 2005; Porret *et al.*, 1974), fire retardants (Habermeier, 1976), and in curing process in textile industry (Trask-Morrel *et al.*, 1988). Solid state chemistry of hydroxy methylated benzimidazole derivatives leading to thermal extrusion of formaldehyde has been reported (Anklekar *et al.*, 1995) by our group. Design and synthesis of benzimidazolone p38 MAP kinase inhibitors (Hammach *et al.*, 2006) is based on the analysis of their crystal structure data. Benzimidazolones have been reported to crystallize as hydrogen bonded molecular tapes (Schwiebert *et al.*, 1996) which has been used to engineer structures of organic solids. In view of the therapeutic importance of aromatic and hetero aromatic compounds (Bansal *et al.*, 1981) containing *N*-hydroxymethyl group, we report here the crystal structure of title compound.

The asymmetric unit of the title compound contains one and a half molecules, one lying on a general position and the other on a twofold rotation axis (Fig.1). Atoms O4 and C13 lie on the twofold rotation axis. The two independent benzimidazole ring systems form a dihedral angle of 18.96 (5) $^{\circ}$. There are no intramolecular hydrogen bonding between *N*-hydroxymethyl and carbonyl groups.

In the crystal, O—H \cdots O hydrogen bonding involving *N*-hydroxymethyl and carbonyl groups results in the formation of three-dimensional network (Fig.2). In addition, C—H \cdots O hydrogen bonds (Table 1) are observed. This type of intermolecular association is similar to that observed in the structure of benzimidazolone (Schwiebert *et al.*, 1996).

Experimental

The title compound was prepared by following a literature method (Zinner *et al.*, 1958). A mixture of 2-hydroxy benzimidazole (13.4 g, 0.01 *M*) and 37% formalin (30 ml, 1*M*) was refluxed for 30 minutes in presence of 100 ml water. The solid product formed was filtered and single crystals were grown by slow evaporation in water (yield 92%, m.p. 433 K). Spectral data IR_{VCO} = 1700 cm $^{-1}$, ν_{OH} = 3300 cm $^{-1}$. ¹H NMR -(CDCl₃+DMSO-d6)δ p.p.m. - 5.3(4*H*, d, CH₂) appeared as singlet on D₂O exchange, 6.2(2*H*, t, OH) vanished on D₂O exchange, 7.4–7.8 (4*H*, m, Ar-H). Mass m/z = 134 (100%).

Refinement

H atoms were positioned at calculated positions [O-H = 0.82 Å and C-H = 0.93–0.97 Å] and refined using a riding model with *U*_{iso}(H) = 1.2*U*_{eq}(C) and 1.5*U*_{eq}(O).

supplementary materials

Figures




Fig. 1. The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. Packing of the molecules viewed down the *b* axis. Hydrogen bonds are shown as dashed lines.

1,3-Bis(hydroxymethyl)benzimidazolin-2-one

Crystal data

C ₉ H ₁₀ N ₂ O ₃	<i>F</i> ₀₀₀ = 1224
<i>M_r</i> = 194.19	<i>D_x</i> = 1.465 Mg m ⁻³
Monoclinic, <i>C</i> 2/c	Melting point: 433 K
Hall symbol: -C 2yc	Mo <i>K</i> α radiation, λ = 0.71073 Å
<i>a</i> = 13.5515 (14) Å	Cell parameters from 2684 reflections
<i>b</i> = 11.0848 (12) Å	θ = 2.3–26.4°
<i>c</i> = 17.6253 (19) Å	μ = 0.11 mm ⁻¹
β = 94.216 (2)°	<i>T</i> = 273 K
<i>V</i> = 2640.4 (5) Å ³	Plate, white
<i>Z</i> = 12	0.22 × 0.20 × 0.10 mm

Data collection

Bruker SMART CCD area-detector diffractometer	2684 independent reflections
Radiation source: fine-focus sealed tube	2395 reflections with $I > 2\sigma(I)$
Monochromator: graphite	R_{int} = 0.025
<i>T</i> = 273 K	θ_{max} = 26.4°
ω and φ scans	θ_{min} = 2.3°
Absorption correction: multi-scan (SADABS; Sheldrick, 2004)	h = -16→16
T_{min} = 0.976, T_{max} = 0.986	k = -13→13
13521 measured reflections	l = -22→22

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
---------------------	--

Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0486P)^2 + 1.5821P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\max} = 0.001$
2684 reflections	$\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$
191 parameters	$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.36019 (9)	0.44213 (11)	0.19000 (7)	0.0555 (3)
H1A	0.4009	0.3984	0.2131	0.083*
O2	0.51268 (9)	0.85515 (11)	-0.04107 (7)	0.0560 (3)
H2A	0.4597	0.8565	-0.0667	0.084*
O3	0.32846 (8)	0.69395 (11)	0.07135 (7)	0.0525 (3)
N1	0.44553 (9)	0.54147 (11)	0.09507 (7)	0.0390 (3)
N2	0.49663 (9)	0.72131 (11)	0.06132 (7)	0.0420 (3)
C1	0.61458 (12)	0.44272 (15)	0.11034 (9)	0.0469 (4)
H1	0.5932	0.3665	0.1238	0.056*
C2	0.71420 (13)	0.46756 (18)	0.10526 (10)	0.0568 (5)
H2	0.7604	0.4063	0.1146	0.068*
C3	0.74639 (13)	0.5819 (2)	0.08647 (10)	0.0582 (5)
H3	0.8138	0.5963	0.0847	0.070*
C4	0.67978 (12)	0.67501 (17)	0.07028 (9)	0.0499 (4)
H4	0.7012	0.7516	0.0576	0.060*
C5	0.58057 (11)	0.64950 (13)	0.07380 (8)	0.0388 (3)
C6	0.54853 (11)	0.53545 (13)	0.09463 (8)	0.0372 (3)
C7	0.41401 (11)	0.65604 (14)	0.07545 (8)	0.0398 (3)
C8	0.49544 (14)	0.84582 (14)	0.03582 (10)	0.0511 (4)
H8A	0.4317	0.8813	0.0440	0.061*
H8B	0.5458	0.8909	0.0658	0.061*
C9	0.37880 (12)	0.44495 (14)	0.11304 (9)	0.0456 (4)

supplementary materials

H9A	0.3166	0.4549	0.0827	0.055*
H9B	0.4072	0.3684	0.0993	0.055*
O4	0.5000	0.27510 (13)	0.2500	0.0467 (4)
O5	0.33811 (9)	0.12882 (13)	0.36950 (7)	0.0628 (4)
H5A	0.2848	0.1462	0.3858	0.094*
N3	0.42148 (9)	0.09011 (11)	0.26254 (7)	0.0396 (3)
C10	0.45046 (15)	-0.24348 (15)	0.25741 (9)	0.0553 (5)
H10	0.4179	-0.3167	0.2617	0.066*
C11	0.39872 (13)	-0.13699 (15)	0.26601 (9)	0.0487 (4)
H11	0.3325	-0.1370	0.2765	0.058*
C12	0.45049 (11)	-0.03066 (13)	0.25817 (8)	0.0385 (3)
C13	0.5000	0.16353 (19)	0.2500	0.0379 (4)
C14	0.32941 (11)	0.13248 (16)	0.28957 (9)	0.0468 (4)
H14A	0.3162	0.2143	0.2722	0.056*
H14B	0.2752	0.0814	0.2701	0.056*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0523 (7)	0.0567 (7)	0.0602 (7)	0.0107 (6)	0.0216 (5)	0.0184 (6)
O2	0.0488 (7)	0.0629 (8)	0.0571 (7)	0.0001 (6)	0.0090 (5)	0.0226 (6)
O3	0.0415 (6)	0.0568 (7)	0.0603 (7)	0.0122 (5)	0.0109 (5)	0.0124 (5)
N1	0.0370 (6)	0.0371 (6)	0.0436 (7)	0.0016 (5)	0.0077 (5)	0.0042 (5)
N2	0.0454 (7)	0.0366 (7)	0.0451 (7)	0.0036 (5)	0.0122 (5)	0.0077 (5)
C1	0.0514 (9)	0.0423 (8)	0.0477 (9)	0.0097 (7)	0.0092 (7)	0.0043 (7)
C2	0.0463 (10)	0.0675 (12)	0.0574 (10)	0.0190 (8)	0.0094 (8)	0.0066 (9)
C3	0.0382 (9)	0.0814 (13)	0.0561 (10)	0.0022 (8)	0.0110 (7)	0.0020 (9)
C4	0.0479 (9)	0.0547 (10)	0.0486 (9)	-0.0070 (7)	0.0143 (7)	0.0017 (7)
C5	0.0424 (8)	0.0402 (8)	0.0349 (7)	0.0031 (6)	0.0095 (6)	0.0012 (6)
C6	0.0392 (8)	0.0397 (8)	0.0335 (7)	0.0020 (6)	0.0079 (5)	-0.0001 (6)
C7	0.0419 (8)	0.0417 (8)	0.0366 (7)	0.0048 (6)	0.0084 (6)	0.0045 (6)
C8	0.0629 (11)	0.0366 (8)	0.0547 (10)	0.0025 (7)	0.0103 (8)	0.0081 (7)
C9	0.0437 (8)	0.0404 (8)	0.0528 (9)	-0.0040 (7)	0.0052 (7)	0.0027 (7)
O4	0.0426 (8)	0.0344 (8)	0.0635 (10)	0.000	0.0068 (7)	0.000
O5	0.0400 (6)	0.0918 (10)	0.0575 (7)	0.0004 (6)	0.0105 (5)	-0.0143 (7)
N3	0.0328 (6)	0.0377 (7)	0.0490 (7)	-0.0017 (5)	0.0069 (5)	-0.0008 (5)
C10	0.0860 (13)	0.0370 (8)	0.0426 (9)	-0.0116 (8)	0.0021 (8)	0.0014 (7)
C11	0.0562 (10)	0.0459 (9)	0.0439 (8)	-0.0124 (7)	0.0032 (7)	0.0019 (7)
C12	0.0416 (8)	0.0376 (8)	0.0363 (7)	-0.0008 (6)	0.0020 (6)	0.0000 (6)
C13	0.0346 (10)	0.0381 (11)	0.0410 (11)	0.000	0.0025 (8)	0.000
C14	0.0329 (8)	0.0504 (9)	0.0575 (10)	0.0001 (7)	0.0052 (6)	-0.0001 (7)

Geometric parameters (\AA , $^\circ$)

O1—C9	1.3981 (19)	C5—C6	1.395 (2)
O1—H1A	0.82	C8—H8A	0.97
O2—C8	1.396 (2)	C8—H8B	0.97
O2—H2A	0.82	C9—H9A	0.97
O3—C7	1.2304 (18)	C9—H9B	0.97

N1—C7	1.3759 (19)	O4—C13	1.237 (3)
N1—C6	1.3980 (19)	O5—C14	1.406 (2)
N1—C9	1.4509 (19)	O5—H5A	0.82
N2—C7	1.371 (2)	N3—C13	1.3704 (17)
N2—C5	1.3925 (19)	N3—C12	1.3990 (19)
N2—C8	1.451 (2)	N3—C14	1.4462 (19)
C1—C6	1.378 (2)	C10—C10 ⁱ	1.387 (4)
C1—C2	1.387 (2)	C10—C11	1.387 (3)
C1—H1	0.93	C11—C12	1.384 (2)
C2—C3	1.388 (3)	C11—H11	0.93
C2—H2	0.93	C12—C12 ⁱ	1.393 (3)
C3—C4	1.387 (3)	C13—N3 ⁱ	1.3704 (17)
C3—H3	0.93	C14—H14A	0.97
C4—C5	1.380 (2)	C14—H14B	0.97
C4—H4	0.93		
C9—O1—H1A	109.4	O2—C8—H8B	109.2
C8—O2—H2A	109.5	N2—C8—H8B	109.2
C7—N1—C6	109.53 (12)	H8A—C8—H8B	107.9
C7—N1—C9	123.27 (13)	O1—C9—N1	112.84 (13)
C6—N1—C9	127.20 (12)	O1—C9—H9A	109.0
C7—N2—C5	109.76 (12)	N1—C9—H9A	109.0
C7—N2—C8	124.58 (14)	O1—C9—H9B	109.0
C5—N2—C8	125.66 (13)	N1—C9—H9B	109.0
C6—C1—C2	117.39 (16)	H9A—C9—H9B	107.8
C6—C1—H1	121.3	C14—O5—H5A	109.5
C2—C1—H1	121.3	C13—N3—C12	109.55 (12)
C1—C2—C3	121.40 (16)	C13—N3—C14	124.00 (13)
C1—C2—H2	119.3	C12—N3—C14	125.57 (13)
C3—C2—H2	119.3	C10 ⁱ —C10—C11	121.66 (10)
C4—C3—C2	121.16 (16)	C10 ⁱ —C10—H10	119.2
C4—C3—H3	119.4	C11—C10—H10	119.2
C2—C3—H3	119.4	C12—C11—C10	116.74 (16)
C5—C4—C3	117.36 (16)	C12—C11—H11	121.6
C5—C4—H4	121.3	C10—C11—H11	121.6
C3—C4—H4	121.3	C11—C12—C12 ⁱ	121.59 (10)
C4—C5—N2	131.54 (15)	C11—C12—N3	131.53 (14)
C4—C5—C6	121.39 (14)	C12 ⁱ —C12—N3	106.88 (8)
N2—C5—C6	107.05 (13)	O4—C13—N3	126.43 (9)
C1—C6—C5	121.26 (14)	O4—C13—N3 ⁱ	126.43 (9)
C1—C6—N1	131.97 (14)	N3—C13—N3 ⁱ	107.14 (18)
C5—C6—N1	106.76 (12)	O5—C14—N3	108.05 (13)
O3—C7—N2	125.99 (14)	O5—C14—H14A	110.1
O3—C7—N1	127.14 (14)	N3—C14—H14A	110.1
N2—C7—N1	106.87 (12)	O5—C14—H14B	110.1
O2—C8—N2	111.86 (14)	N3—C14—H14B	110.1
O2—C8—H8A	109.2	H14A—C14—H14B	108.4
N2—C8—H8A	109.2		

supplementary materials

C6—C1—C2—C3	1.2 (3)	C8—N2—C7—N1	177.40 (14)
C1—C2—C3—C4	-1.5 (3)	C6—N1—C7—O3	-178.71 (15)
C2—C3—C4—C5	0.1 (3)	C9—N1—C7—O3	1.2 (2)
C3—C4—C5—N2	179.76 (16)	C6—N1—C7—N2	1.10 (16)
C3—C4—C5—C6	1.6 (2)	C9—N1—C7—N2	-178.97 (13)
C7—N2—C5—C4	-177.04 (16)	C7—N2—C8—O2	-105.28 (17)
C8—N2—C5—C4	4.1 (3)	C5—N2—C8—O2	73.5 (2)
C7—N2—C5—C6	1.36 (16)	C7—N1—C9—O1	-88.27 (17)
C8—N2—C5—C6	-177.55 (14)	C6—N1—C9—O1	91.64 (17)
C2—C1—C6—C5	0.5 (2)	C10 ⁱ —C10—C11—C12	0.8 (3)
C2—C1—C6—N1	-179.32 (15)	C10—C11—C12—C12 ⁱ	0.5 (3)
C4—C5—C6—C1	-1.9 (2)	C10—C11—C12—N3	179.38 (15)
N2—C5—C6—C1	179.46 (14)	C13—N3—C12—C11	-179.51 (14)
C4—C5—C6—N1	177.94 (13)	C14—N3—C12—C11	10.9 (3)
N2—C5—C6—N1	-0.65 (16)	C13—N3—C12—C12 ⁱ	-0.50 (18)
C7—N1—C6—C1	179.60 (16)	C14—N3—C12—C12 ⁱ	-170.07 (15)
C9—N1—C6—C1	-0.3 (3)	C12—N3—C13—O4	-179.81 (7)
C7—N1—C6—C5	-0.27 (16)	C14—N3—C13—O4	-10.04 (16)
C9—N1—C6—C5	179.81 (14)	C12—N3—C13—N3 ⁱ	0.19 (7)
C5—N2—C7—O3	178.30 (15)	C14—N3—C13—N3 ⁱ	169.96 (16)
C8—N2—C7—O3	-2.8 (2)	C13—N3—C14—O5	-89.75 (16)
C5—N2—C7—N1	-1.52 (16)	C12—N3—C14—O5	78.38 (18)


Symmetry codes: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O1—H1A \cdots O4	0.82	1.99	2.8003 (16)	169
O2—H2A \cdots O5 ⁱⁱ	0.82	1.93	2.7503 (18)	176
O5—H5A \cdots O3 ⁱⁱⁱ	0.82	1.84	2.6551 (17)	175
C3—H3 \cdots O2 ^{iv}	0.93	2.58	3.489 (2)	164
C14—H14B \cdots O1 ⁱⁱⁱ	0.97	2.54	3.364 (2)	143

Symmetry codes: (ii) $x, -y+1, z-1/2$; (iii) $-x+1/2, y-1/2, -z+1/2$; (iv) $-x+3/2, -y+3/2, -z$.

Fig. 1



supplementary materials

Fig. 2

