

tert-Butyl 3-[2,2-bis(ethoxycarbonyl)-vinyl]-2-bromomethyl-1*H*-indole-1-carboxylate

M. Thenmozhi,^a T. Kavitha,^a V. Dhayalan,^b
A. K. Mohanakrishnan^b and M. N. Ponnuswamy^{a*}

^aCentre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and ^bDepartment of Organic Chemistry, University of Madras, Guindy Campus, Chennai 600 025, India
Correspondence e-mail: mnpsy2004@yahoo.com


Received 12 September 2009; accepted 12 October 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.040; wR factor = 0.114; data-to-parameter ratio = 32.0.

In the title compound, $\text{C}_{22}\text{H}_{26}\text{BrNO}_6$, the indole ring system is planar [maximum deviation 0.029 (2) Å]. The *tert*-butyl bound carboxylate group forms a dihedral angle of 17.54 (8)° with the indole ring system. In the crystal, molecules are linked into centrosymmetric $R_2^2(10)$ dimers by paired C—H...O hydrogen bonds.

Related literature

For general background to indoles, see: Gribble (1996); Jing-Ru *et al.* (2007); Ximenes *et al.* (2005). For hybridization, see: Beddoes *et al.* (1986). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{22}\text{H}_{26}\text{BrNO}_6$

$M_r = 480.35$

Triclinic, $P\bar{1}$
 $a = 10.8682$ (3) Å
 $b = 11.1094$ (4) Å
 $c = 11.5699$ (6) Å
 $\alpha = 111.984$ (3)°
 $\beta = 105.841$ (2)°
 $\gamma = 106.926$ (2)°

$V = 1118.51$ (9) Å³
 $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 1.88$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker Kappa APEXII area-detector diffractometer
Absorption correction: multi-scan (SADABS, Sheldrick, 2001)
 $T_{\min} = 0.603$, $T_{\max} = 0.706$

32165 measured reflections
8669 independent reflections
5490 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.028$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.114$
 $S = 1.01$
8669 reflections
271 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.29$ e Å⁻³
 $\Delta\rho_{\min} = -0.60$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C18}-\text{H18A}\cdots\text{O4}^i$	0.97	2.56	3.392 (3)	144

Symmetry code: (i) $-x, -y + 2, -z + 2$.

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT (Bruker, 2004); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97 and PLATON (Spek, 2009).

MT thanks Dr Babu Varghese, SAIF, IIT-Madras, Chennai, India, for his help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2911).

References

- Beddoes, R. L., Dalton, L., Joule, T. A., Mills, O. S., Street, J. D. & Watt, C. I. F. (1986). *J. Chem. Soc. Perkin Trans. 2*, pp. 787–797.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N. L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2004). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Gribble, G. W. (1996). *Comprehensive Heterocyclic Chemistry*, 2nd ed., Vol. 2, pp. 207–257. New York: Pergamon Press.
Jing-Ru, W., Chen-Hsun, T., Samuel, K. K., Dasheng, W., Chia-Hui, L., Hsiao-Ching, Y., Yihui, M., Aaron, S., Chang-Fang, C., Ming-Hsui, T. & Ching-Shih, C. (2007). *Cancer Res.* **67**, 7815–7824.
Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Ximenes, V. F., Paino, I. M. M., Faria-Oliveira, O. M. M., Fonseca, L. M. & Brunetti, I. L. (2005). *Braz. J. Med. Biol. Res.* **38**, 1575–1583.

supplementary materials

Acta Cryst. (2009). E65, o2796 [doi:10.1107/S1600536809041567]

***tert*-Butyl 3-[2,2-bis(ethoxycarbonyl)vinyl]-2-bromomethyl-1*H*-indole-1-carboxylate**

M. Thenmozhi, T. Kavitha, V. Dhayalan, A. K. Mohanakrishnan and M. N. Ponnuswamy

Comment

Indole is a common motif for a drug target and, as such, the development of new diversity-tolerant routes to this privileged biological scaffold continues to be of significant benefit (Gribble *et al.*, 1996) and forms the basis of a wide variety of drugs, including the anti-inflammatory agent indomethacin, reserpine (exploited as hypotensive agent) and sumatriptan (used for the treatment of migraine). The indole derivatives are the effective inhibitors of myeloperoxidase(MPO)-chlorinating activity (Ximenes *et al.*, 2005). Indole-3-carbinol has emerged as a promising chemopreventive agent due to its *in vivo* efficacy in prostate cancer cells of various animal models (Jing-Ru *et al.*, 2007).

The indole ring system of the title molecule (Fig.1) is planar and the bromomethyl group is oriented at an angle of 74.98 (8)°. The *tert* butyl carboxylate group substituted at N1 of the indole ring is in an extended conformation [N1–C10–O1–C11 = 176.24 (16)°]. Both ethoxycarbonyl groups adopt extended conformations as can be seen from torsion angles C16–C17–O3–C18 [-179.61 (16)°], C17–O3–C18–C19 [-156.5 (2)°], C16–C20–O5–C21 [179.18 (14)°] and C20–O5–C21–C22 [-177.9 (2)°]. The sum of bond angles around N1 [360.0 (4)°] indicates that atom N1 exhibits sp^2 hybridization (Beddoes *et al.*, 1986).

The crystal structure is stabilized by C–H...O hydrogen bonds. The molecules form centrosymmetric $R_2^2(10)$ dimers through paired C18–H18A...O4 hydrogen bonds (Fig. 2) (Bernstein *et al.*, 1995).

Experimental

A solution of *tert*-butyl 3-(2,2-di(ethoxycarbonyl)vinyl)-2-methyl-1*H*-indole-1-carboxylate (2 g, 4.98 mmol) in dry carbon tetrachloride (80 ml), azobis(isobutyronitrile)(AIBN) (0.07 g) and finely powdered *N*-bromosuccinimide(NBS) (0.93 g, 5.23 mmol) were added and refluxed for 2 h. Then, the reaction mixture was cooled to room temperature. The floated succinimide was filtered off and washed with carbon tetrachloride (10 ml). The combined filtrate was concentrated *in vacuo* to afford the title compound (1.91 g, 80%) as colourless crystals.

Refinement

H atoms were positioned geometrically (C–H = 0.93–0.97 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H and $1.2U_{\text{eq}}(\text{C})$ for other H atoms. The C18–C19 bond distance was restrained to 1.50 (5) Å.

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. H atoms have been omitted for clarity.




Fig. 2. A view of the crystal packing of molecules, showing C–H...O interactions (dashed lines), leading to dimer formation.

***tert*-Butyl 3-[2,2-bis(ethoxycarbonyl)vinyl]-2-bromomethyl-1*H*-indole-1-carboxylate**

Crystal data

$C_{22}H_{26}BrNO_6$

$M_r = 480.35$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 10.8682$ (3) Å

$b = 11.1094$ (4) Å

$c = 11.5699$ (6) Å

$\alpha = 111.984$ (3)°

$\beta = 105.841$ (2)°

$\gamma = 106.926$ (2)°

$V = 1118.51$ (9) Å³

$Z = 2$

$F_{000} = 496$

$D_x = 1.426$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8669 reflections

$\theta = 2.1$ – 33.8 °

$\mu = 1.88$ mm⁻¹

$T = 293$ K

Block, colourless

$0.30 \times 0.25 \times 0.20$ mm

Data collection

Bruker Kappa APEXII area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293$ K

ω and φ scans

Absorption correction: multi-scan (SADABS, Sheldrick, 2001)

8669 independent reflections

5490 reflections with $I > 2\sigma(I)$

$R_{int} = 0.028$

$\theta_{max} = 33.8$ °

$\theta_{min} = 2.1$ °

$h = -17 \rightarrow 16$

$T_{\min} = 0.603$, $T_{\max} = 0.706$
32165 measured reflections

$k = -16 \rightarrow 17$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.114$	$w = 1/[\sigma^2(F_o^2) + (0.0566P)^2 + 0.1525P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
8669 reflections	$(\Delta/\sigma)_{\max} = 0.001$
271 parameters	$\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\min} = -0.60 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C2	0.09184 (16)	0.74325 (17)	1.16827 (16)	0.0342 (3)
C3	0.07086 (19)	0.8044 (2)	1.28578 (19)	0.0441 (4)
H3	0.1358	0.8307	1.3725	0.053*
C4	-0.0521 (2)	0.8242 (2)	1.2671 (2)	0.0537 (5)
H4	-0.0694	0.8659	1.3435	0.064*
C5	-0.14938 (19)	0.7836 (2)	1.1379 (2)	0.0528 (5)
H5	-0.2299	0.7997	1.1293	0.063*
C6	-0.12946 (17)	0.7201 (2)	1.0217 (2)	0.0427 (4)
H6	-0.1966	0.6909	0.9349	0.051*
C7	-0.00584 (15)	0.70013 (16)	1.03705 (17)	0.0341 (3)
C8	0.04611 (15)	0.63561 (16)	0.94006 (16)	0.0328 (3)
C9	0.17083 (15)	0.63964 (16)	1.01285 (15)	0.0323 (3)
C10	0.31912 (16)	0.73467 (18)	1.26701 (16)	0.0365 (3)
C11	0.51726 (16)	0.67167 (19)	1.32235 (17)	0.0391 (3)
C12	0.5566 (2)	0.5718 (3)	1.2261 (2)	0.0578 (5)

supplementary materials

H12A	0.4829	0.4747	1.1791	0.087*
H12B	0.6448	0.5760	1.2782	0.087*
H12C	0.5671	0.6010	1.1598	0.087*
C13	0.4815 (2)	0.6177 (3)	1.4162 (2)	0.0565 (5)
H13A	0.4495	0.6784	1.4716	0.085*
H13B	0.5647	0.6199	1.4752	0.085*
H13C	0.4074	0.5202	1.3616	0.085*
C14	0.6292 (2)	0.8251 (2)	1.3980 (2)	0.0603 (5)
H14A	0.6402	0.8560	1.3328	0.090*
H14B	0.7180	0.8316	1.4517	0.090*
H14C	0.6009	0.8861	1.4582	0.090*
C15	-0.02203 (15)	0.56833 (17)	0.78999 (16)	0.0361 (3)
H15	-0.0246	0.4788	0.7407	0.043*
C16	-0.08096 (15)	0.62017 (18)	0.71535 (16)	0.0367 (3)
C17	-0.06921 (18)	0.76980 (19)	0.77787 (18)	0.0425 (4)
C18	-0.1952 (2)	0.9111 (2)	0.8069 (3)	0.0658 (6)
H18A	-0.1147	0.9770	0.8966	0.079*
H18B	-0.1864	0.9505	0.7460	0.079*
C19	-0.3273 (3)	0.8959 (3)	0.8200 (4)	0.1028 (11)
H19A	-0.4062	0.8172	0.7355	0.154*
H19B	-0.3367	0.9837	0.8387	0.154*
H19C	-0.3262	0.8766	0.8944	0.154*
C20	-0.14814 (17)	0.5379 (2)	0.56162 (18)	0.0428 (4)
C21	-0.2390 (2)	0.3118 (2)	0.36364 (19)	0.0571 (5)
H21A	-0.1787	0.3480	0.3245	0.069*
H21B	-0.3299	0.3122	0.3247	0.069*
C22	-0.2597 (4)	0.1643 (3)	0.3329 (3)	0.0913 (9)
H22A	-0.1688	0.1647	0.3688	0.137*
H22B	-0.3073	0.1009	0.2347	0.137*
H22C	-0.3165	0.1311	0.3750	0.137*
C23	0.26606 (17)	0.59735 (18)	0.95525 (16)	0.0373 (3)
H23A	0.2140	0.5362	0.8558	0.045*
H23B	0.2997	0.5423	0.9929	0.045*
N1	0.20188 (13)	0.70606 (14)	1.15357 (13)	0.0331 (3)
O1	0.38397 (12)	0.65566 (13)	1.22367 (11)	0.0397 (2)
O2	0.34760 (15)	0.81621 (16)	1.38221 (13)	0.0570 (4)
O3	-0.19532 (13)	0.77004 (13)	0.75104 (14)	0.0476 (3)
O4	0.04207 (15)	0.87474 (16)	0.84512 (19)	0.0707 (4)
O5	-0.17187 (14)	0.40133 (14)	0.51266 (12)	0.0483 (3)
O6	-0.17445 (18)	0.59307 (18)	0.49278 (15)	0.0668 (4)
Br1	0.429587 (19)	0.76941 (2)	0.99976 (2)	0.05385 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C2	0.0370 (7)	0.0337 (8)	0.0387 (8)	0.0197 (6)	0.0174 (6)	0.0204 (7)
C3	0.0481 (9)	0.0488 (10)	0.0408 (9)	0.0256 (8)	0.0221 (8)	0.0219 (8)
C4	0.0559 (10)	0.0622 (12)	0.0581 (12)	0.0346 (10)	0.0359 (10)	0.0291 (10)

C5	0.0445 (9)	0.0635 (12)	0.0712 (13)	0.0335 (9)	0.0335 (9)	0.0392 (11)
C6	0.0344 (7)	0.0478 (10)	0.0531 (10)	0.0204 (7)	0.0174 (7)	0.0311 (8)
C7	0.0342 (7)	0.0320 (7)	0.0414 (8)	0.0164 (6)	0.0152 (6)	0.0224 (7)
C8	0.0346 (7)	0.0316 (7)	0.0334 (7)	0.0160 (6)	0.0114 (6)	0.0184 (6)
C9	0.0361 (7)	0.0323 (7)	0.0299 (7)	0.0182 (6)	0.0112 (6)	0.0163 (6)
C10	0.0384 (7)	0.0385 (8)	0.0335 (8)	0.0202 (7)	0.0123 (6)	0.0185 (7)
C11	0.0334 (7)	0.0500 (10)	0.0342 (8)	0.0223 (7)	0.0086 (6)	0.0221 (7)
C12	0.0541 (10)	0.0722 (14)	0.0528 (11)	0.0435 (10)	0.0189 (9)	0.0271 (10)
C13	0.0495 (10)	0.0813 (15)	0.0576 (12)	0.0335 (10)	0.0208 (9)	0.0497 (11)
C14	0.0430 (9)	0.0559 (12)	0.0664 (14)	0.0151 (9)	0.0151 (9)	0.0268 (11)
C15	0.0340 (7)	0.0369 (8)	0.0340 (8)	0.0155 (6)	0.0096 (6)	0.0181 (7)
C16	0.0311 (7)	0.0401 (8)	0.0356 (8)	0.0142 (6)	0.0084 (6)	0.0209 (7)
C17	0.0402 (8)	0.0410 (9)	0.0419 (9)	0.0148 (7)	0.0079 (7)	0.0256 (8)
C18	0.0744 (14)	0.0448 (11)	0.0804 (16)	0.0348 (11)	0.0287 (12)	0.0297 (11)
C19	0.0846 (18)	0.0762 (19)	0.129 (3)	0.0495 (16)	0.0475 (19)	0.0222 (18)
C20	0.0338 (7)	0.0545 (11)	0.0392 (9)	0.0184 (7)	0.0104 (7)	0.0268 (8)
C21	0.0551 (10)	0.0651 (13)	0.0308 (9)	0.0161 (10)	0.0135 (8)	0.0163 (9)
C22	0.129 (3)	0.0657 (16)	0.0488 (14)	0.0332 (17)	0.0320 (15)	0.0117 (12)
C23	0.0413 (7)	0.0407 (8)	0.0314 (8)	0.0238 (7)	0.0135 (6)	0.0164 (7)
N1	0.0359 (6)	0.0358 (7)	0.0297 (6)	0.0204 (5)	0.0115 (5)	0.0165 (5)
O1	0.0410 (5)	0.0488 (7)	0.0298 (5)	0.0282 (5)	0.0088 (5)	0.0179 (5)
O2	0.0569 (7)	0.0709 (9)	0.0311 (6)	0.0389 (7)	0.0105 (6)	0.0118 (6)
O3	0.0427 (6)	0.0366 (6)	0.0566 (8)	0.0189 (5)	0.0121 (6)	0.0220 (6)
O4	0.0441 (7)	0.0440 (8)	0.0922 (12)	0.0081 (6)	0.0071 (8)	0.0281 (8)
O5	0.0547 (7)	0.0486 (7)	0.0314 (6)	0.0190 (6)	0.0119 (5)	0.0178 (6)
O6	0.0823 (10)	0.0742 (10)	0.0452 (8)	0.0385 (9)	0.0118 (7)	0.0387 (8)
Br1	0.05092 (11)	0.05899 (14)	0.05578 (13)	0.02350 (9)	0.02807 (10)	0.02914 (10)

Geometric parameters (Å, °)

C2—C3	1.389 (2)	C14—H14A	0.96
C2—C7	1.393 (2)	C14—H14B	0.96
C2—N1	1.4057 (17)	C14—H14C	0.96
C3—C4	1.390 (2)	C15—C16	1.331 (2)
C3—H3	0.93	C15—H15	0.93
C4—C5	1.383 (3)	C16—C20	1.489 (2)
C4—H4	0.93	C16—C17	1.493 (2)
C5—C6	1.374 (3)	C17—O4	1.189 (2)
C5—H5	0.93	C17—O3	1.321 (2)
C6—C7	1.399 (2)	C18—C19	1.452 (3)
C6—H6	0.93	C18—O3	1.454 (2)
C7—C8	1.439 (2)	C18—H18A	0.97
C8—C9	1.365 (2)	C18—H18B	0.97
C8—C15	1.458 (2)	C19—H19A	0.96
C9—N1	1.4017 (19)	C19—H19B	0.96
C9—C23	1.4701 (19)	C19—H19C	0.96
C10—O2	1.183 (2)	C20—O6	1.198 (2)
C10—O1	1.3199 (18)	C20—O5	1.318 (2)
C10—N1	1.409 (2)	C21—O5	1.450 (2)

supplementary materials

C11—O1	1.4929 (17)	C21—C22	1.471 (4)
C11—C14	1.498 (3)	C21—H21A	0.97
C11—C12	1.504 (2)	C21—H21B	0.97
C11—C13	1.507 (2)	C22—H22A	0.96
C12—H12A	0.96	C22—H22B	0.96
C12—H12B	0.96	C22—H22C	0.96
C12—H12C	0.96	C23—Br1	1.9654 (17)
C13—H13A	0.96	C23—H23A	0.97
C13—H13B	0.96	C23—H23B	0.97
C13—H13C	0.96		
C3—C2—C7	122.51 (14)	H14B—C14—H14C	109.5
C3—C2—N1	129.77 (15)	C16—C15—C8	127.53 (15)
C7—C2—N1	107.63 (12)	C16—C15—H15	116.2
C2—C3—C4	116.62 (17)	C8—C15—H15	116.2
C2—C3—H3	121.7	C15—C16—C20	121.44 (15)
C4—C3—H3	121.7	C15—C16—C17	122.82 (15)
C5—C4—C3	121.64 (17)	C20—C16—C17	115.34 (14)
C5—C4—H4	119.2	O4—C17—O3	125.11 (17)
C3—C4—H4	119.2	O4—C17—C16	122.78 (16)
C6—C5—C4	121.34 (15)	O3—C17—C16	112.10 (14)
C6—C5—H5	119.3	C19—C18—O3	109.1 (2)
C4—C5—H5	119.3	C19—C18—H18A	109.9
C5—C6—C7	118.48 (17)	O3—C18—H18A	109.9
C5—C6—H6	120.8	C19—C18—H18B	109.9
C7—C6—H6	120.8	O3—C18—H18B	109.9
C2—C7—C6	119.38 (14)	H18A—C18—H18B	108.3
C2—C7—C8	107.57 (12)	C18—C19—H19A	109.5
C6—C7—C8	133.01 (15)	C18—C19—H19B	109.5
C9—C8—C7	107.60 (13)	H19A—C19—H19B	109.5
C9—C8—C15	124.17 (13)	C18—C19—H19C	109.5
C7—C8—C15	128.12 (13)	H19A—C19—H19C	109.5
C8—C9—N1	109.10 (12)	H19B—C19—H19C	109.5
C8—C9—C23	125.35 (14)	O6—C20—O5	125.09 (17)
N1—C9—C23	125.16 (13)	O6—C20—C16	122.46 (18)
O2—C10—O1	127.84 (15)	O5—C20—C16	112.44 (14)
O2—C10—N1	122.20 (14)	O5—C21—C22	107.12 (18)
O1—C10—N1	109.93 (13)	O5—C21—H21A	110.3
O1—C11—C14	110.16 (14)	C22—C21—H21A	110.3
O1—C11—C12	101.60 (13)	O5—C21—H21B	110.3
C14—C11—C12	111.45 (16)	C22—C21—H21B	110.3
O1—C11—C13	108.63 (13)	H21A—C21—H21B	108.5
C14—C11—C13	113.56 (17)	C21—C22—H22A	109.5
C12—C11—C13	110.76 (17)	C21—C22—H22B	109.5
C11—C12—H12A	109.5	H22A—C22—H22B	109.5
C11—C12—H12B	109.5	C21—C22—H22C	109.5
H12A—C12—H12B	109.5	H22A—C22—H22C	109.5
C11—C12—H12C	109.5	H22B—C22—H22C	109.5
H12A—C12—H12C	109.5	C9—C23—Br1	110.47 (11)
H12B—C12—H12C	109.5	C9—C23—H23A	109.6

C11—C13—H13A	109.5	Br1—C23—H23A	109.6
C11—C13—H13B	109.5	C9—C23—H23B	109.6
H13A—C13—H13B	109.5	Br1—C23—H23B	109.6
C11—C13—H13C	109.5	H23A—C23—H23B	108.1
H13A—C13—H13C	109.5	C9—N1—C2	108.08 (12)
H13B—C13—H13C	109.5	C9—N1—C10	129.37 (12)
C11—C14—H14A	109.5	C2—N1—C10	122.55 (13)
C11—C14—H14B	109.5	C10—O1—C11	120.94 (12)
H14A—C14—H14B	109.5	C17—O3—C18	116.27 (15)
C11—C14—H14C	109.5	C20—O5—C21	116.52 (15)
H14A—C14—H14C	109.5		
C7—C2—C3—C4	-1.5 (3)	C17—C16—C20—O6	9.1 (2)
N1—C2—C3—C4	-177.75 (17)	C15—C16—C20—O5	15.1 (2)
C2—C3—C4—C5	0.7 (3)	C17—C16—C20—O5	-171.97 (14)
C3—C4—C5—C6	0.8 (3)	C8—C9—C23—Br1	101.19 (16)
C4—C5—C6—C7	-1.6 (3)	N1—C9—C23—Br1	-70.93 (17)
C3—C2—C7—C6	0.8 (2)	C8—C9—N1—C2	0.55 (17)
N1—C2—C7—C6	177.74 (14)	C23—C9—N1—C2	173.76 (14)
C3—C2—C7—C8	-177.20 (15)	C8—C9—N1—C10	-179.59 (15)
N1—C2—C7—C8	-0.23 (17)	C23—C9—N1—C10	-6.4 (3)
C5—C6—C7—C2	0.8 (2)	C3—C2—N1—C9	176.49 (17)
C5—C6—C7—C8	178.15 (17)	C7—C2—N1—C9	-0.18 (17)
C2—C7—C8—C9	0.57 (17)	C3—C2—N1—C10	-3.4 (3)
C6—C7—C8—C9	-177.01 (17)	C7—C2—N1—C10	179.95 (14)
C2—C7—C8—C15	176.90 (15)	O2—C10—N1—C9	164.58 (17)
C6—C7—C8—C15	-0.7 (3)	O1—C10—N1—C9	-17.1 (2)
C7—C8—C9—N1	-0.68 (17)	O2—C10—N1—C2	-15.6 (3)
C15—C8—C9—N1	-177.20 (14)	O1—C10—N1—C2	162.71 (14)
C7—C8—C9—C23	-173.88 (15)	O2—C10—O1—C11	-5.6 (3)
C15—C8—C9—C23	9.6 (2)	N1—C10—O1—C11	176.24 (13)
C9—C8—C15—C16	-136.69 (17)	C14—C11—O1—C10	-57.2 (2)
C7—C8—C15—C16	47.5 (3)	C12—C11—O1—C10	-175.45 (16)
C8—C15—C16—C20	-179.46 (14)	C13—C11—O1—C10	67.7 (2)
C8—C15—C16—C17	8.1 (3)	O4—C17—O3—C18	1.3 (3)
C15—C16—C17—O4	57.9 (3)	C16—C17—O3—C18	-179.61 (16)
C20—C16—C17—O4	-115.0 (2)	C19—C18—O3—C17	-156.5 (2)
C15—C16—C17—O3	-121.20 (17)	O6—C20—O5—C21	-1.9 (3)
C20—C16—C17—O3	65.95 (19)	C16—C20—O5—C21	179.18 (14)
C15—C16—C20—O6	-163.84 (17)	C22—C21—O5—C20	-177.9 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C18—H18A \cdots O4 ⁱ	0.97	2.56	3.392 (3)	144

Symmetry codes: (i) $-x, -y+2, -z+2$.

Fig. 1




Fig. 2

