

$b = 14.5407(3)$ Å
 $c = 12.7050(4)$ Å
 $\beta = 110.977(1)^\circ$
 $V = 1913.91(10)$ Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

1-Formyl-*r*-2,6-bis(4-methoxyphenyl)-*t*-3,*t*-5-dimethylpiperidin-4-one

T. Kavitha,^a P. Sakthivel,^b S. Ponnuswamy,^b S. S. Ilango^b and M. N. Ponnuswamy^{a*}

^aCentre of Advanced Study in Crystallography and Biophysics, University of Madras, Guindy Campus, Chennai 600 025, India, and ^bDepartment of Chemistry, Government Arts College (Autonomous), Coimbatore 641 018, Tamil Nadu, India
 Correspondence e-mail: mnpsony2004@yahoo.com


Received 12 September 2009; accepted 12 October 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(C-C) = 0.002$ Å; disorder in main residue; R factor = 0.048; wR factor = 0.140; data-to-parameter ratio = 21.6.

In the title compound, C₂₂H₂₅NO₄, the piperidine ring adopts a distorted boat conformation. The methyl groups at the 3 and 5 positions of the piperidine ring are in axial and equatorial orientations, respectively. Both H and O atoms in the aldehyde group are disordered over two positions with occupancies of 0.534 (5) and 0.466 (5). In the crystal, the molecules are linked into a three-dimensional network by C–H···O hydrogen bonds.

Related literature

For general background to piperidine derivatives, see: Escalano & Amat (2006); Wang & Wuorola (1992); Grishina *et al.* (1994). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

C₂₂H₂₅NO₄
 $M_r = 367.43$

Monoclinic, $P2_1/n$
 $a = 11.0954(4)$ Å

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (*SADABS*; Sheldrick, 2001)
 $R_{\text{min}} = 0.974$, $T_{\text{max}} = 0.974$

24720 measured reflections
 5524 independent reflections
 3703 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.029$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.140$
 $S = 1.05$
 5524 reflections

256 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.23$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C3—H3···O1 ⁱ	0.98	2.48	3.429 (2)	163
C15—H15C···O2 ⁱⁱ	0.96	2.53	3.240 (4)	131
C20—H20···O1 ⁱⁱⁱ	0.93	2.51	3.432 (2)	171
Symmetry codes: (i) $-x + 2, -y, -z$; (ii) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$				

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

TK thanks Dr Babu Varghese, SAIF, IIT-Madras, Chennai, India, for his help with the data collection. SP thanks the UGC, India, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2909).

References

- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Escalano, C. & Amat, M. (2006). *Chem. Eur. J.* **12**, 8198–8207.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Grishina, G. V., Gaidarov, E. L. & Zefirov, N. S. (1994). *Chem. Heterocycl. Compd.* **30**, 1401–1426.
- Sheldrick, G. M. (2001). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wang, C.-L. & Wuorola, M. A. (1992). *Org. Prep. Proc. Int.* **24**, 585–621.

supplementary materials

Acta Cryst. (2009). E65, o2818 [doi:10.1107/S1600536809041609]

1-Formyl-r-2,c-6-bis(4-methoxyphenyl)-t-3,t-5-dimethylpiperidin-4-one

T. Kavitha, P. Sakthivel, S. Ponnuswamy, S. S. Ilango and M. N. Ponnuswamy

Comment

The piperidine ring is a common feature occurring in many biologically active natural products and therapeutic agents. Piperidine containing entities constitute important targets for pharmaceutical research (Escolano & Amat, 2006). Piperidine derivatives, namely 4-piperidones are synthetic intermediates in the preparation of various alkaloids and pharmaceutical products (Wang *et al.*, 1992; Grishina *et al.*, 1994).

In the title molecule (Fig.1), the piperidine ring adopts a distorted boat conformation. The methyl groups at 3 and 5 positions of the piperidine ring are in axial and equatorial orientations [$\text{N}1—\text{C}2—\text{C}3—\text{C}14 = -63.02$ (15) $^\circ$ and $\text{N}1—\text{C}6—\text{C}5—\text{C}15 = 176.60$ (11) $^\circ$]. The phenyl rings at 2 and 6 positions of the piperidine ring are axially [$\text{C}4—\text{C}3—\text{C}2—\text{C}8 = -68.32$ (14) $^\circ$] and equatorially [$\text{C}4—\text{C}5—\text{C}6—\text{C}16 = 176.69$ (10) $^\circ$] oriented. The dihedral angle between the two phenyl rings is 41.97 (8) $^\circ$.

Centrosymmetrically related molecules form $R_2^2(8)$ (Bernstein *et al.*, 1995) dimers through C—H \cdots O hydrogen bonds involving atoms C3 and O1. The dimers are linked into a zigzag C(8) chain running along the *b* axis by intermolecular C—H \cdots O hydrogen bonds involving atoms C20 and O1 (Table 1). Further, C15—H15C \cdots O2 interactions link the chains along the *c* axis to form a three-dimensional network.

Experimental

An ice-cold solution of acetic-formic anhydride was prepared from acetic anhydride (10 ml) and 85% formic acid (5 ml) and was added slowly to a cold solution of r-2,c-6-bis(4-methoxyphenyl)-t-3,t-5-dimethylpiperidin-4-one (1.69 g) in benzene (30 ml). The reaction mixture was stirred at room temperature for 5 h. The organic layer was separated, dried over anhydrous Na_2SO_4 and concentrated. The resulting mass was purified by crystallization from benzene-petroleum ether (333–353 K) in the ratio 1:1.

Refinement

The O and H atoms of the formyl group is disordered over two positions with occupancies of 0.534 (5) and 0.466 (5). H atoms were positioned geometrically ($\text{C}-\text{H} = 0.93$ –0.98 Å) and allowed to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$ and $1.2U_{\text{eq}}(\text{C})$. A rotating group model was used for the methoxy methyl groups.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound. Displacement ellipsoids are drawn at the 30% probability level. Both disorder components are shown. H atoms have been omitted for clarity.




Fig. 2. The crystal packing of the title compound. H atoms not involved in hydrogen bonding (dashed lines) have been omitted.

1-Formyl-*t*-2,6-bis(4-methoxyphenyl)-*t*-3,5-dimethylpiperidin-4-one

Crystal data

C ₂₂ H ₂₅ NO ₄	$F_{000} = 784$
$M_r = 367.43$	$D_x = 1.275 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 5524 reflections
$a = 11.0954 (4) \text{ \AA}$	$\theta = 2.1\text{--}29.9^\circ$
$b = 14.5407 (3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 12.7050 (4) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 110.977 (1)^\circ$	Block, colourless
$V = 1913.91 (10) \text{ \AA}^3$	$0.30 \times 0.20 \times 0.20 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer	5524 independent reflections
Radiation source: fine-focus sealed tube	3703 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.029$
$T = 293 \text{ K}$	$\theta_{\max} = 29.9^\circ$
ω and φ scans	$\theta_{\min} = 2.1^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 2001)	$h = -15 \rightarrow 12$
$T_{\min} = 0.974$, $T_{\max} = 0.974$	$k = -20 \rightarrow 19$
24720 measured reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.048$	H-atom parameters constrained
$wR(F^2) = 0.140$	$w = 1/[\sigma^2(F_o^2) + (0.0598P)^2 + 0.3189P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\max} = 0.001$
5524 reflections	$\Delta\rho_{\max} = 0.23 \text{ e \AA}^{-3}$
256 parameters	$\Delta\rho_{\min} = -0.23 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.87856 (11)	0.03534 (7)	0.05971 (10)	0.0574 (3)	
O2	1.0929 (3)	0.3733 (2)	0.3086 (2)	0.1143 (15)	0.534 (5)
O2'	1.2050 (3)	0.3881 (2)	0.2198 (3)	0.1036 (16)	0.466 (5)
O3	0.92438 (12)	0.33025 (8)	-0.37580 (9)	0.0647 (3)	
O4	0.60958 (11)	0.58982 (7)	0.13202 (9)	0.0600 (3)	
N1	1.03399 (11)	0.28962 (8)	0.15129 (9)	0.0446 (3)	
C2	1.08936 (13)	0.22990 (9)	0.08641 (11)	0.0433 (3)	
H2	1.1831	0.2389	0.1177	0.052*	
C3	1.06451 (13)	0.13017 (10)	0.10905 (12)	0.0455 (3)	
H3	1.0976	0.0906	0.0631	0.055*	
C4	0.92172 (13)	0.11238 (9)	0.07622 (11)	0.0408 (3)	
C5	0.83662 (12)	0.19522 (9)	0.06811 (11)	0.0397 (3)	
H5	0.8182	0.2227	-0.0064	0.048*	
C6	0.90568 (13)	0.26797 (9)	0.15605 (10)	0.0402 (3)	
H6	0.9198	0.2415	0.2305	0.048*	
C7	1.1102 (2)	0.34871 (18)	0.22651 (19)	0.0930 (7)	
H7	1.1815	0.3720	0.2133	0.112*	0.534 (5)
H7'	1.0887	0.3619	0.2918	0.112*	0.466 (5)

supplementary materials

C8	1.04508 (12)	0.25691 (9)	-0.03708 (11)	0.0405 (3)
C9	1.01446 (16)	0.19282 (10)	-0.12353 (13)	0.0536 (4)
H9	1.0200	0.1305	-0.1062	0.064*
C10	0.97608 (17)	0.21971 (11)	-0.23422 (13)	0.0582 (4)
H10	0.9554	0.1753	-0.2906	0.070*
C11	0.96764 (14)	0.31160 (10)	-0.26313 (12)	0.0472 (3)
C12	1.00016 (15)	0.37652 (10)	-0.17907 (13)	0.0517 (4)
H12	0.9968	0.4387	-0.1968	0.062*
C13	1.03795 (14)	0.34869 (10)	-0.06774 (13)	0.0496 (3)
H13	1.0593	0.3932	-0.0115	0.060*
C14	1.13194 (17)	0.10344 (14)	0.23312 (14)	0.0675 (5)
H14A	1.2219	0.1187	0.2568	0.101*
H14B	1.1227	0.0385	0.2417	0.101*
H14C	1.0934	0.1364	0.2786	0.101*
C15	0.70806 (15)	0.16822 (11)	0.07688 (17)	0.0619 (4)
H15A	0.6531	0.2213	0.0637	0.093*
H15B	0.7224	0.1444	0.1509	0.093*
H15C	0.6675	0.1219	0.0216	0.093*
C16	0.82499 (13)	0.35393 (9)	0.14503 (11)	0.0406 (3)
C17	0.79081 (14)	0.40931 (9)	0.05079 (11)	0.0435 (3)
H17	0.8174	0.3932	-0.0084	0.052*
C18	0.71800 (14)	0.48820 (9)	0.04184 (11)	0.0444 (3)
H18	0.6960	0.5246	-0.0225	0.053*
C19	0.67825 (13)	0.51235 (9)	0.12957 (11)	0.0433 (3)
C20	0.70878 (16)	0.45640 (11)	0.22319 (12)	0.0533 (4)
H20	0.6803	0.4715	0.2815	0.064*
C21	0.78103 (16)	0.37843 (10)	0.23047 (12)	0.0518 (4)
H21	0.8009	0.3413	0.2940	0.062*
C22	0.9119 (2)	0.42401 (13)	-0.40951 (16)	0.0748 (5)
H22A	0.8799	0.4276	-0.4903	0.112*
H22B	0.9946	0.4535	-0.3797	0.112*
H22C	0.8527	0.4543	-0.3813	0.112*
C23	0.58744 (18)	0.65403 (12)	0.04332 (15)	0.0655 (5)
H23A	0.5442	0.7070	0.0578	0.098*
H23B	0.5346	0.6263	-0.0265	0.098*
H23C	0.6685	0.6723	0.0386	0.098*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0733 (7)	0.0333 (5)	0.0769 (7)	-0.0017 (5)	0.0407 (6)	-0.0034 (5)
O2	0.096 (2)	0.140 (3)	0.085 (2)	0.0135 (19)	0.0052 (15)	-0.072 (2)
O2'	0.0510 (18)	0.085 (2)	0.156 (3)	-0.0229 (15)	0.0140 (18)	-0.046 (2)
O3	0.0798 (8)	0.0614 (7)	0.0550 (6)	0.0062 (6)	0.0265 (6)	0.0104 (5)
O4	0.0746 (7)	0.0486 (6)	0.0589 (6)	0.0254 (5)	0.0266 (5)	0.0045 (5)
N1	0.0411 (6)	0.0452 (6)	0.0437 (6)	-0.0008 (5)	0.0105 (5)	-0.0090 (5)
C2	0.0353 (6)	0.0446 (8)	0.0507 (7)	0.0034 (5)	0.0162 (6)	-0.0003 (6)
C3	0.0479 (8)	0.0426 (8)	0.0514 (8)	0.0141 (6)	0.0243 (6)	0.0085 (6)

C4	0.0526 (8)	0.0337 (7)	0.0423 (7)	0.0041 (6)	0.0244 (6)	0.0024 (5)
C5	0.0399 (6)	0.0335 (6)	0.0467 (7)	0.0032 (5)	0.0166 (5)	0.0013 (5)
C6	0.0479 (7)	0.0379 (7)	0.0366 (6)	0.0073 (6)	0.0173 (5)	0.0014 (5)
C7	0.0728 (13)	0.1154 (18)	0.0846 (14)	-0.0275 (13)	0.0205 (11)	-0.0533 (13)
C8	0.0373 (6)	0.0380 (7)	0.0518 (7)	0.0001 (5)	0.0227 (6)	0.0001 (6)
C9	0.0768 (11)	0.0344 (7)	0.0575 (9)	0.0039 (7)	0.0337 (8)	0.0006 (6)
C10	0.0841 (12)	0.0440 (8)	0.0535 (9)	0.0016 (8)	0.0330 (8)	-0.0051 (7)
C11	0.0452 (7)	0.0494 (8)	0.0525 (8)	0.0033 (6)	0.0242 (6)	0.0044 (6)
C12	0.0569 (9)	0.0368 (7)	0.0652 (9)	-0.0021 (6)	0.0264 (7)	0.0065 (7)
C13	0.0550 (8)	0.0379 (7)	0.0588 (8)	-0.0072 (6)	0.0237 (7)	-0.0043 (6)
C14	0.0601 (10)	0.0778 (12)	0.0615 (10)	0.0226 (9)	0.0179 (8)	0.0232 (9)
C15	0.0469 (8)	0.0496 (9)	0.0950 (13)	0.0020 (7)	0.0323 (8)	-0.0004 (8)
C16	0.0500 (7)	0.0346 (7)	0.0385 (6)	0.0058 (5)	0.0175 (5)	-0.0017 (5)
C17	0.0535 (8)	0.0428 (7)	0.0378 (6)	0.0063 (6)	0.0208 (6)	-0.0009 (5)
C18	0.0526 (8)	0.0398 (7)	0.0402 (7)	0.0050 (6)	0.0159 (6)	0.0042 (6)
C19	0.0473 (7)	0.0361 (7)	0.0460 (7)	0.0059 (6)	0.0160 (6)	-0.0032 (5)
C20	0.0732 (10)	0.0496 (9)	0.0462 (8)	0.0149 (7)	0.0324 (7)	0.0006 (6)
C21	0.0746 (10)	0.0454 (8)	0.0412 (7)	0.0165 (7)	0.0280 (7)	0.0066 (6)
C22	0.0856 (13)	0.0710 (12)	0.0707 (11)	0.0147 (10)	0.0316 (10)	0.0259 (9)
C23	0.0765 (11)	0.0466 (9)	0.0673 (10)	0.0210 (8)	0.0183 (9)	0.0075 (8)

Geometric parameters (Å, °)

O1—C4	1.2067 (16)	C10—C11	1.380 (2)
O2—C7	1.182 (3)	C10—H10	0.93
O2'—C7	1.227 (4)	C11—C12	1.373 (2)
O3—C11	1.3642 (18)	C12—C13	1.384 (2)
O3—C22	1.421 (2)	C12—H12	0.93
O4—C19	1.3665 (16)	C13—H13	0.93
O4—C23	1.416 (2)	C14—H14A	0.96
N1—C7	1.337 (2)	C14—H14B	0.96
N1—C2	1.4743 (17)	C14—H14C	0.96
N1—C6	1.4803 (17)	C15—H15A	0.96
C2—C8	1.5183 (19)	C15—H15B	0.96
C2—C3	1.523 (2)	C15—H15C	0.96
C2—H2	0.98	C16—C17	1.3787 (18)
C3—C4	1.509 (2)	C16—C21	1.3865 (18)
C3—C14	1.533 (2)	C17—C18	1.3840 (19)
C3—H3	0.98	C17—H17	0.93
C4—C5	1.5111 (18)	C18—C19	1.3829 (19)
C5—C15	1.521 (2)	C18—H18	0.93
C5—C6	1.5302 (18)	C19—C20	1.379 (2)
C5—H5	0.98	C20—C21	1.372 (2)
C6—C16	1.5145 (18)	C20—H20	0.93
C6—H6	0.98	C21—H21	0.93
C7—H7	0.93	C22—H22A	0.96
C7—H7'	0.96	C22—H22B	0.96
C8—C13	1.385 (2)	C22—H22C	0.96
C8—C9	1.387 (2)	C23—H23A	0.96

supplementary materials

C9—C10	1.372 (2)	C23—H23B	0.96
C9—H9	0.93	C23—H23C	0.96
C11—O3—C22	117.82 (13)	C12—C11—C10	118.99 (14)
C19—O4—C23	117.70 (12)	C11—C12—C13	119.53 (14)
C7—N1—C2	119.66 (14)	C11—C12—H12	120.2
C7—N1—C6	118.59 (14)	C13—C12—H12	120.2
C2—N1—C6	119.80 (10)	C12—C13—C8	122.36 (14)
N1—C2—C8	112.33 (11)	C12—C13—H13	118.8
N1—C2—C3	108.39 (11)	C8—C13—H13	118.8
C8—C2—C3	115.34 (12)	C3—C14—H14A	109.5
N1—C2—H2	106.8	C3—C14—H14B	109.5
C8—C2—H2	106.8	H14A—C14—H14B	109.5
C3—C2—H2	106.8	C3—C14—H14C	109.5
C4—C3—C2	110.75 (11)	H14A—C14—H14C	109.5
C4—C3—C14	108.56 (12)	H14B—C14—H14C	109.5
C2—C3—C14	112.39 (13)	C5—C15—H15A	109.5
C4—C3—H3	108.3	C5—C15—H15B	109.5
C2—C3—H3	108.3	H15A—C15—H15B	109.5
C14—C3—H3	108.3	C5—C15—H15C	109.5
O1—C4—C3	121.28 (12)	H15A—C15—H15C	109.5
O1—C4—C5	121.92 (13)	H15B—C15—H15C	109.5
C3—C4—C5	116.79 (11)	C17—C16—C21	117.74 (12)
C4—C5—C15	111.63 (11)	C17—C16—C6	122.14 (11)
C4—C5—C6	111.35 (11)	C21—C16—C6	120.11 (12)
C15—C5—C6	111.25 (12)	C16—C17—C18	121.77 (12)
C4—C5—H5	107.5	C16—C17—H17	119.1
C15—C5—H5	107.5	C18—C17—H17	119.1
C6—C5—H5	107.5	C19—C18—C17	119.31 (12)
N1—C6—C16	111.47 (11)	C19—C18—H18	120.3
N1—C6—C5	110.78 (10)	C17—C18—H18	120.3
C16—C6—C5	112.22 (10)	O4—C19—C20	115.68 (12)
N1—C6—H6	107.4	O4—C19—C18	124.68 (12)
C16—C6—H6	107.4	C20—C19—C18	119.63 (12)
C5—C6—H6	107.4	C21—C20—C19	120.17 (12)
O2—C7—O2'	109.5 (3)	C21—C20—H20	119.9
O2—C7—N1	124.4 (3)	C19—C20—H20	119.9
O2'—C7—N1	126.1 (3)	C20—C21—C16	121.34 (13)
O2—C7—H7	117.8	C20—C21—H21	119.3
N1—C7—H7	117.8	C16—C21—H21	119.3
O2'—C7—H7'	116.8	O3—C22—H22A	109.5
N1—C7—H7'	117.2	O3—C22—H22B	109.5
C13—C8—C9	116.86 (13)	H22A—C22—H22B	109.5
C13—C8—C2	120.31 (12)	O3—C22—H22C	109.5
C9—C8—C2	122.79 (12)	H22A—C22—H22C	109.5
C10—C9—C8	121.23 (14)	H22B—C22—H22C	109.5
C10—C9—H9	119.4	O4—C23—H23A	109.5
C8—C9—H9	119.4	O4—C23—H23B	109.5
C9—C10—C11	121.01 (14)	H23A—C23—H23B	109.5
C9—C10—H10	119.5	O4—C23—H23C	109.5

C11—C10—H10	119.5	H23A—C23—H23C	109.5
O3—C11—C12	125.11 (13)	H23B—C23—H23C	109.5
O3—C11—C10	115.89 (13)		
C7—N1—C2—C8	−109.80 (19)	N1—C2—C8—C9	−140.48 (13)
C6—N1—C2—C8	86.30 (14)	C3—C2—C8—C9	−15.60 (18)
C7—N1—C2—C3	121.58 (18)	C13—C8—C9—C10	−1.5 (2)
C6—N1—C2—C3	−42.33 (16)	C2—C8—C9—C10	−179.28 (14)
N1—C2—C3—C4	58.59 (14)	C8—C9—C10—C11	0.5 (3)
C8—C2—C3—C4	−68.32 (14)	C22—O3—C11—C12	0.1 (2)
N1—C2—C3—C14	−63.02 (15)	C22—O3—C11—C10	178.86 (15)
C8—C2—C3—C14	170.07 (11)	C9—C10—C11—O3	−177.91 (14)
C2—C3—C4—O1	160.79 (13)	C9—C10—C11—C12	0.9 (2)
C14—C3—C4—O1	−75.37 (17)	O3—C11—C12—C13	177.39 (14)
C2—C3—C4—C5	−20.74 (16)	C10—C11—C12—C13	−1.3 (2)
C14—C3—C4—C5	103.09 (14)	C11—C12—C13—C8	0.3 (2)
O1—C4—C5—C15	19.05 (19)	C9—C8—C13—C12	1.1 (2)
C3—C4—C5—C15	−159.40 (12)	C2—C8—C13—C12	178.93 (13)
O1—C4—C5—C6	144.05 (13)	N1—C6—C16—C17	60.77 (17)
C3—C4—C5—C6	−34.40 (15)	C5—C6—C16—C17	−64.16 (17)
C7—N1—C6—C16	57.6 (2)	N1—C6—C16—C21	−120.35 (14)
C2—N1—C6—C16	−138.34 (12)	C5—C6—C16—C21	114.73 (15)
C7—N1—C6—C5	−176.69 (17)	C21—C16—C17—C18	1.7 (2)
C2—N1—C6—C5	−12.62 (16)	C6—C16—C17—C18	−179.42 (13)
C4—C5—C6—N1	51.38 (14)	C16—C17—C18—C19	0.0 (2)
C15—C5—C6—N1	176.60 (11)	C23—O4—C19—C20	173.65 (15)
C4—C5—C6—C16	176.69 (10)	C23—O4—C19—C18	−5.9 (2)
C15—C5—C6—C16	−58.09 (15)	C17—C18—C19—O4	177.91 (13)
C2—N1—C7—O2	−150.5 (3)	C17—C18—C19—C20	−1.7 (2)
C6—N1—C7—O2	13.6 (4)	O4—C19—C20—C21	−177.95 (15)
C2—N1—C7—O2'	32.2 (4)	C18—C19—C20—C21	1.7 (2)
C6—N1—C7—O2'	−163.7 (3)	C19—C20—C21—C16	0.0 (3)
N1—C2—C8—C13	41.78 (17)	C17—C16—C21—C20	−1.7 (2)
C3—C2—C8—C13	166.66 (12)	C6—C16—C21—C20	179.39 (14)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C3—H3···O1 ⁱ	0.98	2.48	3.429 (2)	163
C15—H15C···O2 ⁱⁱ	0.96	2.53	3.240 (4)	131
C20—H20···O1 ⁱⁱⁱ	0.93	2.51	3.432 (2)	171

Symmetry codes: (i) $-x+2, -y, -z$; (ii) $x-1/2, -y+1/2, z-1/2$; (iii) $-x+3/2, y+1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

