

N'-[*E*]-4-Hydroxy-3-methoxybenzylidene]benzohydrazide

Zahid Shafiq,^a Muhammad Yaqub,^a M. Nawaz Tahir,^{b*} Abid Hussain^a and M. Saeed Iqbal^c

^aDepartment of Chemistry, Bahauddin Zakariya University, Multan 60800, Pakistan,

^bDepartment of Physics, University of Sargodha, Sargodha, Pakistan, and

^cDepartment of Chemistry, Government College University, Lahore, Pakistan

Correspondence e-mail: dmntahir_uos@yahoo.com


Received 23 October 2009; accepted 23 October 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; disorder in main residue; R factor = 0.048; wR factor = 0.151; data-to-parameter ratio = 15.9.

In the title compound, $\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_3$, the phenyl ring is disordered over two set of sites with an occupancy ratio of 0.810 (3):0.190 (3); the dihedral angle between the two components is 72.3 (4)°. The benzene and phenyl rings are oriented at dihedral angles of 69.18 (8) and 26.0 (5)° (major and minor orientations, respectively), and an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond occurs. In the crystal, molecules are linked by $\text{N}-\text{H}\cdots\text{O}$, $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions, generating a three-dimensional network.

Related literature

For related structures, see: Shafiq *et al.* (2009*a,b*); Shi (2005).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_3$
 $M_r = 270.28$
 Tetragonal, $I4_1/a$
 $a = 18.6994$ (8) Å

$c = 15.7223$ (12) Å
 $V = 5497.6$ (5) Å³
 $Z = 16$
 Mo $K\alpha$ radiation

$\mu = 0.09$ mm⁻¹
 $T = 296$ K

$0.28 \times 0.24 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.976$, $T_{\max} = 0.979$

15310 measured reflections
 3393 independent reflections
 1656 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.151$
 $S = 0.99$
 3393 reflections

214 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.14$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\cdots\text{O}3$	0.82	2.21	2.653 (2)	115
$\text{N}1-\text{H}1\cdots\text{O}1^{\text{i}}$	0.86	2.14	2.8633 (19)	142
$\text{O}2-\text{H}2\cdots\text{O}1^{\text{ii}}$	0.82	2.05	2.773 (2)	146
$\text{C}2\text{A}-\text{H}2\text{A}\cdots\text{O}1^{\text{i}}$	0.93	2.43	3.165 (2)	136

Symmetry codes: (i) $-y + \frac{1}{4}, x - \frac{1}{4}, z - \frac{1}{4}$; (ii) $y - \frac{1}{4}, -x + \frac{1}{4}, -z + \frac{5}{4}$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

AH gratefully acknowledges the Higher Education Commission, Islamabad, Pakistan, for providing him with a Scholarship under the Indigenous PhD Program (PIN 063-121531-PS3-127).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5175).

References

- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837-838.
 Shafiq, Z., Yaqub, M., Tahir, M. N., Hussain, A. & Iqbal, M. S. (2009*a*). *Acta Cryst.* **E65**, o2496.
 Shafiq, Z., Yaqub, M., Tahir, M. N., Hussain, A. & Iqbal, M. S. (2009*b*). *Acta Cryst.* **E65**, o2501.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112-122.
 Shi, J. (2005). *Acta Cryst.* **E61**, o3917-o3918.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148-155.

supplementary materials

Acta Cryst. (2009). E65, o2898 [doi:10.1107/S1600536809044122]

N'-[(*E*)-4-Hydroxy-3-methoxybenzylidene]benzohydrazide

Z. Shafiq, M. Yaqub, M. N. Tahir, A. Hussain and M. S. Iqbal

Comment

Recently we have reported the crystal structures of (II) *N'*-[(*E*)-(4-Bromo-2-thienyl)methylene]isonicotinohydrazide (Shafiq *et al.*, 2009*a*) and (III) *N'*-[(*E*)-(4-Bromo-2-thienyl)methylidene]benzohydrazide 0.06-hydrate (Shafiq *et al.*, 2009*b*). The title compound (I, Fig. 1), has been prepared in continuation of synthesizing various hydrazide derivatives.

The crystal structure of (IV) (*E*)-*N*-Benzoyl-*N'*-(3-hydroxy-4-methoxybenzylidene)hydrazine (Shi, 2005) has been published which differs from (I) due to positional change of hydroxy and methoxy.

In the title compound benzene ring of benzohydrazide is disordered over two set of sites with occupancy ratio of 0.810 (3):0.190 (3). The majority and minority groups A (C1A—C6A) and B (C1B—C6B) respectively, are oriented at a dihedral angle of 72.27 (36)°. The benzene ring C (C9—C14) of 4-Hydroxy-3-methoxyphenyl is of course planar. The dihedral angle between A/C and B/C is 69.18 (8)° and 25.98 (51)°, respectively. The molecules are stabilized in the form of three dimensional polymeric network due to strong intra as well as intermolecular H-bondings (Table 1, Fig.2).

Experimental

To a hot stirred solution of benzoic hydrazide (1.36 g, 0.01 mol) in ethanol (15 ml) was added vanillin (1.52 g, 0.01 mol). The resultant mixture was then heated under reflux. After an hour precipitates were formed. The reaction mixture was further heated about 30 min for the completion of the reaction which was monitored through TLC. The reaction mixture was cooled to room temperature, filtered and washed with hot ethanol. Colourless prisms of (I) were obtained by recrystallization of the crude product in 1,4-dioxan:ethanol (1:1) after four days.

Refinement

The disordered phenyl rings A (C1A—C6A) and B (C1B—C6B) were refined using AFIX 66 and all atoms have independent anisotropic thermal parameters.

The H-atoms were positioned geometrically (O—H = 0.82 Å, N—H = 0.86 Å, C—H = 0.93–0.96 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

Figures




Fig. 1. View of (I) showing the major disorder component. The displacement ellipsoids are drawn at the 50% probability level. H-atoms are shown by circles of arbitrary radius.




Fig. 2. View of (I) showing the minor disorder component. The displacement ellipsoids are drawn at the 50% probability level. H-atoms are shown by circles of arbitrary radius.




Fig. 3. The partial packing of (I), which shows that molecules form three dimensional polymeric network. H-atoms not involved in H-bondings have been omitted for clarity.

N'-[(*E*)-4-Hydroxy-3-methoxybenzylidene]benzohydrazide

Crystal data

$C_{15}H_{14}N_2O_3$

$M_r = 270.28$

Tetragonal, $I4_1/a$

Hall symbol: $-I\ 4ad$

$a = 18.6994$ (8) Å

$b = 18.6994$ (8) Å

$c = 15.7223$ (12) Å

$\alpha = 90^\circ$

$\beta = 90^\circ$

$\gamma = 90^\circ$

$V = 5497.6$ (5) Å³

$Z = 16$

$F_{000} = 2272$

$D_x = 1.306$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3393 reflections

$\theta = 1.7\text{--}28.3^\circ$

$\mu = 0.09$ mm⁻¹

$T = 296$ K

Prism, colourless

$0.28 \times 0.24 \times 0.22$ mm

Data collection

Bruker Kappa APEXII CCD diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

Detector resolution: 7.40 pixels mm⁻¹

$T = 296$ K

ω scans

Absorption correction: multi-scan (SADABS; Bruker, 2005)

$T_{\min} = 0.976$, $T_{\max} = 0.979$

15310 measured reflections

3393 independent reflections

1656 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 28.3^\circ$

$\theta_{\min} = 1.7^\circ$

$h = -24 \rightarrow 24$

$k = -24 \rightarrow 24$

$l = -20 \rightarrow 20$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.048$

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$wR(F^2) = 0.151$	$w = 1/[\sigma^2(F_o^2) + (0.071P)^2 + 0.721P]$
$S = 0.99$	where $P = (F_o^2 + 2F_c^2)/3$
3393 reflections	$(\Delta/\sigma)_{\max} < 0.001$
214 parameters	$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	$\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$
	Extinction coefficient: ?

Special details

Geometry. Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	0.31015 (8)	0.09452 (8)	0.50043 (7)	0.0639 (5)	
O2	-0.07897 (8)	-0.13841 (8)	0.63057 (9)	0.0715 (6)	
O3	-0.00552 (9)	-0.02825 (9)	0.69117 (8)	0.0751 (6)	
N1	0.23647 (9)	0.04528 (9)	0.40397 (9)	0.0577 (6)	
N2	0.19267 (9)	0.01533 (10)	0.46558 (9)	0.0578 (6)	
C1A	0.32949 (10)	0.12512 (10)	0.35611 (10)	0.0491 (15)	0.813 (3)
C2A	0.29168 (9)	0.15525 (12)	0.28897 (12)	0.0678 (10)	0.813 (3)
C3A	0.32797 (12)	0.19079 (13)	0.22442 (11)	0.0813 (15)	0.813 (3)
C4A	0.40207 (12)	0.19620 (12)	0.22701 (11)	0.0810 (16)	0.813 (3)
C5A	0.43989 (9)	0.16606 (11)	0.29416 (13)	0.0760 (12)	0.813 (3)
C6A	0.40360 (10)	0.13052 (10)	0.35871 (10)	0.0619 (10)	0.813 (3)
C7	0.29070 (11)	0.08778 (10)	0.42620 (10)	0.0495 (7)	
C8	0.14721 (12)	-0.02955 (12)	0.43724 (12)	0.0584 (7)	
C9	0.09173 (11)	-0.06060 (11)	0.49065 (11)	0.0550 (7)	
C10	0.05165 (12)	-0.11712 (12)	0.46193 (12)	0.0638 (8)	
C11	-0.00483 (12)	-0.14403 (11)	0.51003 (13)	0.0636 (8)	
C12	-0.02232 (11)	-0.11260 (11)	0.58606 (12)	0.0559 (7)	
C13	0.01804 (11)	-0.05554 (11)	0.61574 (11)	0.0564 (7)	
C14	0.07470 (11)	-0.02999 (11)	0.56916 (11)	0.0576 (7)	
C15	0.03358 (16)	0.02801 (17)	0.72880 (14)	0.0990 (11)	
C2B	0.3203 (5)	0.1020 (4)	0.2723 (5)	0.059 (4)	0.187 (3)
C3B	0.3506 (6)	0.1424 (5)	0.2075 (4)	0.076 (5)	0.187 (3)
C4B	0.3772 (7)	0.2103 (5)	0.2246 (5)	0.078 (7)	0.187 (3)
C5B	0.3737 (6)	0.2377 (4)	0.3067 (6)	0.092 (6)	0.187 (3)
C6B	0.3434 (5)	0.1973 (4)	0.3716 (4)	0.070 (5)	0.187 (3)
C1B	0.3167 (5)	0.1294 (4)	0.3544 (4)	0.055 (7)	0.187 (3)

supplementary materials

H5A	0.48947	0.16967	0.29590	0.0913*	0.813 (3)
H6A	0.42890	0.11036	0.40363	0.0743*	0.813 (3)
H10	0.06233	-0.13774	0.40962	0.0766*	
H11	-0.03064	-0.18328	0.49061	0.0763*	
H14	0.10184	0.00784	0.58997	0.0691*	
H15A	0.03617	0.06754	0.68996	0.1486*	
H15B	0.08102	0.01179	0.74192	0.1486*	
H15C	0.01014	0.04298	0.78013	0.1486*	
H8	0.14959	-0.04294	0.38034	0.0701*	
H1	0.22866	0.03642	0.35111	0.0692*	
H2	-0.08451	-0.11459	0.67387	0.1072*	
H2A	0.24210	0.15164	0.28723	0.0814*	0.813 (3)
H3A	0.30267	0.21096	0.17949	0.0978*	0.813 (3)
H4A	0.42635	0.21997	0.18383	0.0970*	0.813 (3)
H2B	0.30242	0.05661	0.26082	0.0709*	0.187 (3)
H3B	0.35296	0.12407	0.15254	0.0912*	0.187 (3)
H4B	0.39750	0.23733	0.18123	0.0932*	0.187 (3)
H5B	0.39151	0.28312	0.31819	0.1097*	0.187 (3)
H6B	0.34098	0.21565	0.42646	0.0839*	0.187 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0871 (11)	0.0700 (10)	0.0346 (6)	-0.0024 (8)	-0.0072 (6)	-0.0015 (6)
O2	0.0768 (11)	0.0677 (11)	0.0699 (9)	-0.0116 (8)	0.0145 (8)	0.0004 (7)
O3	0.0816 (11)	0.0916 (12)	0.0521 (8)	-0.0187 (9)	0.0146 (7)	-0.0128 (7)
N1	0.0632 (11)	0.0790 (12)	0.0309 (7)	-0.0057 (9)	0.0011 (7)	0.0057 (7)
N2	0.0605 (11)	0.0739 (12)	0.0390 (8)	0.0006 (10)	0.0052 (7)	0.0098 (8)
C1A	0.064 (3)	0.050 (3)	0.0334 (19)	0.0030 (19)	-0.0007 (14)	-0.0007 (18)
C2A	0.0692 (19)	0.083 (2)	0.0512 (14)	-0.0108 (16)	-0.0127 (13)	0.0175 (14)
C3A	0.099 (3)	0.092 (3)	0.0528 (16)	-0.015 (2)	-0.0081 (18)	0.0242 (17)
C4A	0.105 (3)	0.068 (3)	0.070 (2)	-0.017 (2)	0.0224 (19)	-0.0005 (19)
C5A	0.064 (2)	0.077 (2)	0.087 (2)	-0.0071 (16)	0.0126 (16)	-0.0008 (16)
C6A	0.0558 (19)	0.0681 (19)	0.0618 (15)	0.0009 (13)	-0.0030 (12)	0.0011 (12)
C7	0.0602 (13)	0.0538 (12)	0.0346 (9)	0.0093 (10)	-0.0013 (8)	0.0004 (8)
C8	0.0648 (14)	0.0671 (14)	0.0433 (10)	0.0071 (11)	0.0044 (9)	0.0017 (9)
C9	0.0589 (13)	0.0566 (13)	0.0494 (10)	0.0046 (10)	0.0034 (9)	0.0041 (9)
C10	0.0740 (15)	0.0623 (15)	0.0552 (11)	0.0051 (12)	0.0086 (11)	-0.0063 (10)
C11	0.0727 (16)	0.0500 (13)	0.0680 (13)	-0.0021 (11)	0.0014 (11)	-0.0046 (10)
C12	0.0618 (14)	0.0516 (13)	0.0544 (11)	0.0014 (10)	0.0027 (10)	0.0070 (9)
C13	0.0640 (14)	0.0615 (14)	0.0437 (10)	0.0030 (11)	0.0025 (9)	0.0010 (9)
C14	0.0619 (14)	0.0634 (14)	0.0474 (10)	-0.0053 (10)	0.0008 (9)	0.0013 (9)
C15	0.110 (2)	0.127 (2)	0.0599 (14)	-0.0388 (18)	0.0150 (13)	-0.0343 (14)
C2B	0.053 (7)	0.058 (8)	0.067 (7)	-0.013 (6)	0.004 (5)	0.002 (6)
C3B	0.081 (10)	0.089 (11)	0.058 (7)	0.012 (9)	0.008 (6)	0.007 (7)
C4B	0.098 (15)	0.084 (14)	0.051 (9)	-0.001 (11)	0.026 (8)	0.008 (8)
C5B	0.094 (11)	0.063 (9)	0.118 (12)	-0.014 (8)	-0.003 (8)	-0.001 (8)
C6B	0.084 (9)	0.066 (9)	0.060 (6)	-0.011 (7)	0.000 (6)	-0.005 (6)

C1B 0.029 (7) 0.066 (15) 0.071 (14) -0.016 (7) -0.013 (7) -0.013 (11)

Geometric parameters (Å, °)

O1—C7	1.229 (2)	C9—C14	1.397 (3)
O2—C12	1.358 (2)	C9—C10	1.372 (3)
O3—C13	1.364 (2)	C10—C11	1.393 (3)
O3—C15	1.411 (3)	C11—C12	1.372 (3)
O2—H2	0.8200	C12—C13	1.388 (3)
N1—N2	1.387 (2)	C13—C14	1.374 (3)
N1—C7	1.335 (3)	C2A—H2A	0.9300
N2—C8	1.275 (3)	C2B—H2B	0.9300
N1—H1	0.8600	C3A—H3A	0.9300
C1A—C6A	1.390 (3)	C3B—H3B	0.9300
C1A—C7	1.493 (2)	C4A—H4A	0.9300
C1A—C2A	1.390 (3)	C4B—H4B	0.9300
C1B—C2B	1.390 (10)	C5A—H5A	0.9300
C1B—C6B	1.391 (11)	C5B—H5B	0.9300
C1B—C7	1.455 (7)	C6A—H6A	0.9300
C2A—C3A	1.390 (3)	C6B—H6B	0.9300
C2B—C3B	1.389 (12)	C8—H8	0.9300
C3A—C4A	1.390 (3)	C10—H10	0.9300
C3B—C4B	1.390 (14)	C11—H11	0.9300
C4A—C5A	1.390 (3)	C14—H14	0.9300
C4B—C5B	1.390 (12)	C15—H15B	0.9600
C5A—C6A	1.390 (3)	C15—H15C	0.9600
C5B—C6B	1.390 (12)	C15—H15A	0.9600
C8—C9	1.456 (3)		
C13—O3—C15	118.43 (18)	C12—C13—C14	120.50 (17)
C12—O2—H2	109.00	O3—C13—C12	113.85 (18)
N2—N1—C7	120.41 (14)	C9—C14—C13	120.28 (19)
N1—N2—C8	114.55 (15)	C1A—C2A—H2A	120.00
N2—N1—H1	120.00	C3A—C2A—H2A	120.00
C7—N1—H1	120.00	C1B—C2B—H2B	120.00
C2A—C1A—C6A	120.00 (16)	C3B—C2B—H2B	120.00
C2A—C1A—C7	120.21 (17)	C4A—C3A—H3A	120.00
C6A—C1A—C7	119.78 (15)	C2A—C3A—H3A	120.00
C2B—C1B—C6B	120.0 (7)	C4B—C3B—H3B	120.00
C2B—C1B—C7	122.6 (6)	C2B—C3B—H3B	120.00
C6B—C1B—C7	117.2 (5)	C3A—C4A—H4A	120.00
C1A—C2A—C3A	120.00 (17)	C5A—C4A—H4A	120.00
C1B—C2B—C3B	120.0 (7)	C5B—C4B—H4B	120.00
C2A—C3A—C4A	120.01 (18)	C3B—C4B—H4B	120.00
C2B—C3B—C4B	120.1 (7)	C6A—C5A—H5A	120.00
C3A—C4A—C5A	120.00 (18)	C4A—C5A—H5A	120.00
C3B—C4B—C5B	120.0 (8)	C4B—C5B—H5B	120.00
C4A—C5A—C6A	120.00 (17)	C6B—C5B—H5B	120.00
C4B—C5B—C6B	120.0 (8)	C1A—C6A—H6A	120.00
C1A—C6A—C5A	120.00 (16)	C5A—C6A—H6A	120.00

supplementary materials

C1B—C6B—C5B	120.0 (6)	C5B—C6B—H6B	120.00
O1—C7—C1A	120.62 (18)	C1B—C6B—H6B	120.00
O1—C7—C1B	125.7 (3)	C9—C8—H8	119.00
O1—C7—N1	122.31 (17)	N2—C8—H8	119.00
N1—C7—C1A	117.02 (14)	C9—C10—H10	120.00
N1—C7—C1B	111.7 (3)	C11—C10—H10	120.00
N2—C8—C9	122.43 (17)	C12—C11—H11	120.00
C8—C9—C10	120.45 (17)	C10—C11—H11	120.00
C8—C9—C14	120.57 (18)	C13—C14—H14	120.00
C10—C9—C14	118.80 (18)	C9—C14—H14	120.00
C9—C10—C11	120.91 (18)	O3—C15—H15C	109.00
C10—C11—C12	119.94 (19)	O3—C15—H15B	109.00
C11—C12—C13	119.53 (19)	H15B—C15—H15C	109.00
O2—C12—C11	118.86 (18)	H15A—C15—H15B	109.00
O2—C12—C13	121.61 (17)	H15A—C15—H15C	109.00
O3—C13—C14	125.62 (18)	O3—C15—H15A	109.00
C15—O3—C13—C12	177.4 (2)	C4A—C5A—C6A—C1A	0.0 (3)
C15—O3—C13—C14	-4.5 (3)	N2—C8—C9—C10	-169.1 (2)
C7—N1—N2—C8	-173.59 (19)	N2—C8—C9—C14	15.8 (3)
N2—N1—C7—O1	10.0 (3)	C8—C9—C10—C11	-175.5 (2)
N2—N1—C7—C1A	-172.51 (17)	C14—C9—C10—C11	-0.3 (3)
N1—N2—C8—C9	-173.17 (18)	C8—C9—C14—C13	174.08 (19)
C6A—C1A—C2A—C3A	0.0 (3)	C10—C9—C14—C13	-1.1 (3)
C7—C1A—C2A—C3A	179.15 (19)	C9—C10—C11—C12	1.9 (3)
C2A—C1A—C6A—C5A	0.0 (3)	C10—C11—C12—O2	178.01 (19)
C7—C1A—C6A—C5A	-179.15 (17)	C10—C11—C12—C13	-2.1 (3)
C2A—C1A—C7—O1	-140.5 (2)	O2—C12—C13—O3	-1.2 (3)
C2A—C1A—C7—N1	42.0 (3)	O2—C12—C13—C14	-179.40 (18)
C6A—C1A—C7—O1	38.6 (3)	C11—C12—C13—O3	178.91 (18)
C6A—C1A—C7—N1	-138.89 (19)	C11—C12—C13—C14	0.7 (3)
C1A—C2A—C3A—C4A	0.0 (3)	O3—C13—C14—C9	-177.09 (19)
C2A—C3A—C4A—C5A	0.0 (3)	C12—C13—C14—C9	0.9 (3)
C3A—C4A—C5A—C6A	0.0 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2 \cdots O3	0.82	2.21	2.653 (2)	115
N1—H1 \cdots O1 ⁱ	0.86	2.14	2.8633 (19)	142
O2—H2 \cdots O1 ⁱⁱ	0.82	2.05	2.773 (2)	146
C2A—H2A \cdots O1 ⁱ	0.93	2.43	3.165 (2)	136

Symmetry codes: (i) $-y+1/4, x-1/4, z-1/4$; (ii) $y-1/4, -x+1/4, -z+5/4$.

Fig. 1




Fig. 2




Fig. 3

