

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3,19-Diacetyl-12-nitromethyl-14-deoxyandrographolide

Juan Jia and Cheng Yao*


College of Science, Nanjing University of Technology, Xinmofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: yaocheng@njut.edu.cn

Received 15 October 2009; accepted 23 October 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.072; wR factor = 0.199; data-to-parameter ratio = 8.0.In the crystal of the title compound, $\text{C}_{24}\text{H}_{33}\text{NO}_9$, intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules.

Related literature

For general background, see: Thunuguntla *et al.* (2004). For bond-length data, see: Allen *et al.* (1987).

Experimental

Crystal data

 $\text{C}_{24}\text{H}_{33}\text{NO}_9$ $M_r = 479.51$ Monoclinic, $P2_1$
 $a = 10.533$ (2) Å
 $b = 12.756$ (3) Å
 $c = 10.659$ (2) Å
 $\beta = 117.04$ (3)°
 $V = 1275.6$ (4) Å³ $Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 293$ K
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4
diffractometer
Absorption correction: none
2565 measured reflections
2431 independent reflections1945 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$
3 standard reflections
every 200 reflections
intensity decay: 1%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.199$
 $S = 1.00$
2431 reflections
302 parameters14 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.69$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.48$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C15}-\text{H15A}\cdots\text{O8}^{\text{i}}$	0.97	2.59	3.312 (18)	131
$\text{C22}-\text{H22A}\cdots\text{O9}^{\text{ii}}$	0.97	2.47	3.388 (13)	158
$\text{C22}-\text{H22B}\cdots\text{O1}^{\text{iii}}$	0.97	2.51	3.301 (13)	139
$\text{C24}-\text{H24B}\cdots\text{O3}^{\text{iv}}$	0.97	2.56	3.376 (9)	142

Symmetry codes: (i) $x - 1, y, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + 1$; (iii) $-x + 1, y - \frac{1}{2}, -z + 1$; (iv) $x + 1, y, z$.Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5144).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Thunuguntla, S. S. R., Nyavanandi, V. K. & Nanduri, S. (2004). *Tetrahedron Lett.* **45**, 9357–9360.

supplementary materials

Acta Cryst. (2009). E65, o2965 [doi:10.1107/S160053680904402X]

3,19-Diacetyl-12-nitromethyl-14-deoxyandrographolide

J. Jia and C. Yao

Experimental

Andrographolide (15 g) and nitromethane (20 ml) in dry methanol were stirred in the presence of sodium methoxide at room temperature for 3 h. After confirming the completion of reaction, the mixture was washed with brine. The organic phase was evaporated in vacuo, and the residue was recrystallized by ethyl acetate. The product (10 g) and acetic anhydride (20 ml) was refluxed for 10 min. After confirming the completion of reaction, the mixture was washed with brine. The organic phase was evaporated in vacuo to afford corresponding product by flash chromatography. Colourless blocks of (I) were recrystallised from ethyl acetate.

Refinement

The H atoms were positioned geometrically (C—H = 0.93–0.97 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$.

The deepest difference hole is 0.12 Å from the N atom and the highest difference peak is 0.18 Å from atom O9.

Figures




Fig. 1. The molecular structure of (I) showing 30% displacement ellipsoids.

3,19-Diacetyl-12-nitromethyl-14-deoxyandrographolide

Crystal data

$\text{C}_{24}\text{H}_{33}\text{NO}_9$

$M_r = 479.51$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 10.533$ (2) Å

$b = 12.756$ (3) Å

$c = 10.659$ (2) Å

$\beta = 117.04$ (3)°

$V = 1275.6$ (4) Å³

$Z = 2$

$F_{000} = 512$

$D_x = 1.248$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$

$\mu = 0.10$ mm⁻¹

$T = 293$ K

Block, colourless

$0.30 \times 0.20 \times 0.20$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer	$R_{\text{int}} = 0.024$
Radiation source: fine-focus sealed tube	$\theta_{\text{max}} = 25.3^\circ$
Monochromator: graphite	$\theta_{\text{min}} = 2.2^\circ$
$T = 293$ K	$h = -11 \rightarrow 12$
$\omega/2\theta$ scans	$k = -15 \rightarrow 0$
Absorption correction: none	$l = -12 \rightarrow 0$
2565 measured reflections	3 standard reflections
2431 independent reflections	every 200 reflections
1945 reflections with $I > 2\sigma(I)$	intensity decay: 1%

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.072$	H-atom parameters constrained
$wR(F^2) = 0.199$	$w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 1.120P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
2431 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
302 parameters	$\Delta\rho_{\text{max}} = 0.69 \text{ e } \text{\AA}^{-3}$
14 restraints	$\Delta\rho_{\text{min}} = -0.48 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: SHELXL97 (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.063 (8)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N	0.7992 (9)	-0.3512 (6)	0.7140 (12)	0.1177 (19)
O1	0.1814 (7)	0.3398 (4)	0.6418 (6)	0.0915 (17)
C1	0.0804 (12)	0.4065 (7)	0.7867 (12)	0.108 (3)

H1A	0.0769	0.4728	0.7427	0.162*
H1B	0.1387	0.4127	0.8866	0.162*
H1C	-0.0143	0.3859	0.7679	0.162*
O2	0.1466 (6)	0.2330 (3)	0.7887 (5)	0.0719 (13)
C2	0.1425 (7)	0.3259 (5)	0.7293 (8)	0.0645 (16)
O3	-0.1429 (6)	-0.0145 (5)	0.8828 (7)	0.0935 (18)
C3	-0.1806 (8)	0.1612 (7)	0.9391 (9)	0.084 (2)
H3A	-0.2668	0.1342	0.9365	0.126*
H3B	-0.2033	0.2160	0.8704	0.126*
H3C	-0.1207	0.1887	1.0312	0.126*
O4	0.0165 (4)	0.1100 (3)	0.9106 (4)	0.0555 (10)
C4	-0.1040 (7)	0.0749 (5)	0.9066 (6)	0.0587 (15)
C5	0.1112 (7)	0.0355 (5)	0.8929 (6)	0.0559 (14)
H5A	0.0813	-0.0350	0.9011	0.067*
H5B	0.2070	0.0454	0.9681	0.067*
C6	0.1141 (6)	0.0463 (5)	0.7499 (5)	0.0470 (12)
O5	0.3609 (8)	-0.3524 (5)	0.7817 (8)	0.115 (2)
C7	0.1806 (6)	-0.0527 (4)	0.7167 (5)	0.0424 (12)
H7A	0.1659	-0.0412	0.6201	0.051*
O6	0.4623 (8)	-0.1819 (6)	0.3207 (5)	0.106 (2)
C8	0.3456 (6)	-0.0689 (4)	0.8036 (5)	0.0456 (12)
O7	0.4337 (11)	-0.3352 (6)	0.4088 (7)	0.145 (3)
O8	0.8651 (14)	-0.3399 (8)	0.6600 (15)	0.218 (7)
C9	0.4175 (7)	0.0340 (5)	0.7949 (8)	0.0644 (16)
H9A	0.5187	0.0294	0.8580	0.077*
H9B	0.4063	0.0420	0.6999	0.077*
O9	0.7740 (7)	-0.4391 (5)	0.7441 (9)	0.1177 (19)
C10	0.3570 (7)	0.1318 (5)	0.8330 (8)	0.0655 (17)
H10A	0.3776	0.1280	0.9313	0.079*
H10B	0.4038	0.1937	0.8209	0.079*
C11	0.2003 (7)	0.1420 (4)	0.7444 (7)	0.0558 (14)
H11A	0.1817	0.1527	0.6465	0.067*
C12	-0.0425 (6)	0.0581 (5)	0.6332 (6)	0.0596 (15)
H12A	-0.0433	0.0659	0.5433	0.089*
H12B	-0.0848	0.1188	0.6523	0.089*
H12C	-0.0959	-0.0032	0.6321	0.089*
C13	0.3944 (7)	-0.0991 (5)	0.9598 (6)	0.0595 (15)
H13A	0.3676	-0.0447	1.0053	0.089*
H13B	0.4961	-0.1075	1.0066	0.089*
H13C	0.3498	-0.1637	0.9640	0.089*
C14	0.0996 (6)	-0.1543 (4)	0.7078 (6)	0.0522 (13)
H14A	-0.0017	-0.1426	0.6505	0.063*
H14B	0.1150	-0.1743	0.8014	0.063*
C15	0.1489 (7)	-0.2431 (5)	0.6436 (7)	0.0616 (16)
H15A	0.1009	-0.3075	0.6453	0.074*
H15B	0.1234	-0.2267	0.5462	0.074*
C16	0.3051 (6)	-0.2581 (5)	0.7227 (6)	0.0524 (13)
C17	0.3904 (6)	-0.1603 (4)	0.7310 (6)	0.0467 (12)
H17A	0.3592	-0.1379	0.6334	0.056*

supplementary materials

C18	0.5504 (6)	-0.1759 (5)	0.7935 (6)	0.0554 (13)
H18A	0.5837	-0.2095	0.8847	0.066*
H18B	0.5959	-0.1078	0.8086	0.066*
C19	0.5979 (6)	-0.2425 (5)	0.7016 (6)	0.0526 (14)
H19A	0.5479	-0.3098	0.6836	0.063*
C20	0.5592 (6)	-0.1911 (5)	0.5610 (6)	0.0524 (13)
C21	0.4825 (11)	-0.2478 (8)	0.4296 (8)	0.087 (2)
C22	0.5278 (9)	-0.0844 (7)	0.3765 (9)	0.088 (2)
H22A	0.4587	-0.0279	0.3415	0.105*
H22B	0.6036	-0.0701	0.3511	0.105*
C23	0.5853 (8)	-0.0952 (6)	0.5302 (8)	0.0693 (17)
H23A	0.6326	-0.0428	0.5957	0.083*
C24	0.7548 (7)	-0.2639 (5)	0.7812 (8)	0.0649 (16)
H24A	0.7784	-0.2830	0.8774	0.078*
H24B	0.8072	-0.2008	0.7838	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N	0.109 (3)	0.059 (2)	0.196 (6)	0.008 (3)	0.079 (4)	-0.003 (3)
O1	0.128 (5)	0.059 (3)	0.100 (4)	0.001 (3)	0.062 (4)	0.019 (3)
C1	0.147 (8)	0.051 (4)	0.152 (9)	0.012 (5)	0.090 (7)	0.001 (5)
O2	0.112 (4)	0.036 (2)	0.090 (3)	0.001 (2)	0.066 (3)	0.002 (2)
C2	0.068 (4)	0.037 (3)	0.083 (4)	0.001 (3)	0.030 (3)	0.007 (3)
O3	0.114 (4)	0.073 (3)	0.120 (4)	-0.029 (3)	0.077 (4)	-0.020 (3)
C3	0.072 (4)	0.097 (6)	0.093 (5)	0.009 (4)	0.046 (4)	-0.015 (5)
O4	0.066 (2)	0.047 (2)	0.062 (2)	-0.0022 (19)	0.037 (2)	-0.0078 (18)
C4	0.069 (4)	0.061 (4)	0.054 (3)	-0.009 (3)	0.035 (3)	-0.007 (3)
C5	0.072 (4)	0.055 (3)	0.051 (3)	0.006 (3)	0.036 (3)	-0.001 (3)
C6	0.061 (3)	0.043 (3)	0.040 (3)	-0.001 (3)	0.026 (2)	0.000 (2)
O5	0.157 (6)	0.081 (4)	0.128 (5)	0.010 (4)	0.082 (5)	0.007 (4)
C7	0.049 (3)	0.044 (3)	0.036 (2)	-0.002 (2)	0.021 (2)	-0.001 (2)
O6	0.155 (6)	0.095 (4)	0.065 (3)	-0.007 (4)	0.048 (3)	0.000 (3)
C8	0.051 (3)	0.043 (3)	0.044 (3)	-0.002 (2)	0.022 (2)	-0.005 (2)
O7	0.241 (9)	0.085 (5)	0.084 (4)	-0.049 (6)	0.054 (5)	-0.028 (4)
O8	0.294 (13)	0.122 (7)	0.410 (18)	-0.038 (8)	0.310 (15)	-0.057 (9)
C9	0.062 (3)	0.044 (3)	0.092 (5)	-0.008 (3)	0.039 (3)	-0.005 (3)
O9	0.109 (3)	0.059 (2)	0.196 (6)	0.008 (3)	0.079 (4)	-0.003 (3)
C10	0.075 (4)	0.038 (3)	0.088 (5)	-0.010 (3)	0.040 (4)	-0.011 (3)
C11	0.078 (4)	0.036 (3)	0.062 (3)	0.002 (3)	0.040 (3)	0.000 (3)
C12	0.065 (3)	0.058 (4)	0.053 (3)	0.008 (3)	0.023 (3)	0.002 (3)
C13	0.072 (4)	0.060 (4)	0.044 (3)	0.009 (3)	0.023 (3)	-0.006 (3)
C14	0.053 (3)	0.048 (3)	0.061 (3)	-0.003 (3)	0.030 (3)	-0.007 (3)
C15	0.059 (4)	0.054 (4)	0.073 (4)	-0.010 (3)	0.031 (3)	-0.019 (3)
C16	0.065 (4)	0.044 (3)	0.056 (3)	0.001 (3)	0.034 (3)	-0.007 (2)
C17	0.051 (3)	0.045 (3)	0.046 (3)	0.003 (2)	0.024 (2)	-0.001 (2)
C18	0.056 (3)	0.054 (3)	0.054 (3)	0.003 (3)	0.023 (3)	-0.003 (3)
C19	0.053 (3)	0.045 (3)	0.062 (3)	0.004 (3)	0.029 (3)	0.001 (3)

C20	0.055 (3)	0.044 (3)	0.066 (3)	0.007 (3)	0.034 (3)	0.001 (3)
C21	0.114 (6)	0.082 (6)	0.069 (5)	-0.005 (5)	0.044 (4)	-0.006 (4)
C22	0.104 (6)	0.083 (6)	0.099 (5)	0.020 (5)	0.066 (5)	0.033 (5)
C23	0.076 (4)	0.059 (4)	0.085 (5)	0.003 (3)	0.048 (4)	0.000 (3)
C24	0.058 (4)	0.047 (3)	0.085 (4)	0.006 (3)	0.029 (3)	0.005 (3)

Geometric parameters (Å, °)

N—O8	1.094 (10)	C9—H9B	0.9700
N—O9	1.228 (10)	C10—C11	1.488 (9)
N—C24	1.508 (10)	C10—H10A	0.9700
O1—C2	1.190 (8)	C10—H10B	0.9700
C1—C2	1.492 (11)	C11—H11A	0.9800
C1—H1A	0.9600	C12—H12A	0.9600
C1—H1B	0.9600	C12—H12B	0.9600
C1—H1C	0.9600	C12—H12C	0.9600
O2—C2	1.335 (7)	C13—H13A	0.9600
O2—C11	1.460 (7)	C13—H13B	0.9600
O3—C4	1.200 (8)	C13—H13C	0.9600
C3—C4	1.495 (10)	C14—C15	1.531 (8)
C3—H3A	0.9600	C14—H14A	0.9700
C3—H3B	0.9600	C14—H14B	0.9700
C3—H3C	0.9600	C15—C16	1.481 (9)
O4—C4	1.327 (7)	C15—H15A	0.9700
O4—C5	1.451 (7)	C15—H15B	0.9700
C5—C6	1.544 (7)	C16—C17	1.517 (8)
C5—H5A	0.9700	C17—C18	1.518 (7)
C5—H5B	0.9700	C17—H17A	0.9800
C6—C11	1.538 (8)	C18—C19	1.541 (8)
C6—C12	1.557 (8)	C18—H18A	0.9700
C6—C7	1.561 (8)	C18—H18B	0.9700
O5—C16	1.361 (9)	C19—C24	1.500 (8)
C7—C14	1.530 (8)	C19—C20	1.512 (8)
C7—C8	1.568 (7)	C19—H19A	0.9800
C7—H7A	0.9800	C20—C23	1.328 (9)
O6—C21	1.369 (10)	C20—C21	1.452 (10)
O6—C22	1.416 (12)	C22—C23	1.473 (11)
C8—C9	1.539 (8)	C22—H22A	0.9700
C8—C13	1.551 (8)	C22—H22B	0.9700
C8—C17	1.585 (7)	C23—H23A	0.9300
O7—C21	1.205 (12)	C24—H24A	0.9700
C9—C10	1.538 (9)	C24—H24B	0.9700
C9—H9A	0.9700		
O8—N—O9	121.4 (10)	C6—C12—H12A	109.5
O8—N—C24	124.1 (9)	C6—C12—H12B	109.5
O9—N—C24	113.5 (9)	H12A—C12—H12B	109.5
C2—C1—H1A	109.5	C6—C12—H12C	109.5
C2—C1—H1B	109.5	H12A—C12—H12C	109.5
H1A—C1—H1B	109.5	H12B—C12—H12C	109.5

supplementary materials

C2—C1—H1C	109.5	C8—C13—H13A	109.5
H1A—C1—H1C	109.5	C8—C13—H13B	109.5
H1B—C1—H1C	109.5	H13A—C13—H13B	109.5
C2—O2—C11	119.1 (5)	C8—C13—H13C	109.5
O1—C2—O2	123.8 (7)	H13A—C13—H13C	109.5
O1—C2—C1	126.2 (7)	H13B—C13—H13C	109.5
O2—C2—C1	110.1 (6)	C7—C14—C15	111.2 (5)
C4—C3—H3A	109.5	C7—C14—H14A	109.4
C4—C3—H3B	109.5	C15—C14—H14A	109.4
H3A—C3—H3B	109.5	C7—C14—H14B	109.4
C4—C3—H3C	109.5	C15—C14—H14B	109.4
H3A—C3—H3C	109.5	H14A—C14—H14B	108.0
H3B—C3—H3C	109.5	C16—C15—C14	110.9 (5)
C4—O4—C5	118.6 (5)	C16—C15—H15A	109.5
O3—C4—O4	123.9 (6)	C14—C15—H15A	109.5
O3—C4—C3	125.5 (7)	C16—C15—H15B	109.5
O4—C4—C3	110.7 (6)	C14—C15—H15B	109.5
O4—C5—C6	113.2 (5)	H15A—C15—H15B	108.1
O4—C5—H5A	108.9	O5—C16—C15	120.8 (6)
C6—C5—H5A	108.9	O5—C16—C17	125.5 (6)
O4—C5—H5B	108.9	C15—C16—C17	113.7 (5)
C6—C5—H5B	108.9	C16—C17—C18	115.3 (5)
H5A—C5—H5B	107.8	C16—C17—C8	109.5 (4)
C11—C6—C5	112.6 (5)	C18—C17—C8	113.6 (4)
C11—C6—C12	108.8 (5)	C16—C17—H17A	105.9
C5—C6—C12	108.0 (4)	C18—C17—H17A	105.9
C11—C6—C7	107.5 (4)	C8—C17—H17A	105.9
C5—C6—C7	111.3 (4)	C17—C18—C19	114.2 (5)
C12—C6—C7	108.6 (4)	C17—C18—H18A	108.7
C14—C7—C6	113.9 (4)	C19—C18—H18A	108.7
C14—C7—C8	111.2 (4)	C17—C18—H18B	108.7
C6—C7—C8	117.7 (4)	C19—C18—H18B	108.7
C14—C7—H7A	104.1	H18A—C18—H18B	107.6
C6—C7—H7A	104.1	C24—C19—C20	111.7 (5)
C8—C7—H7A	104.1	C24—C19—C18	109.5 (5)
C21—O6—C22	109.0 (6)	C20—C19—C18	111.9 (5)
C9—C8—C13	110.1 (5)	C24—C19—H19A	107.9
C9—C8—C7	107.2 (4)	C20—C19—H19A	107.9
C13—C8—C7	113.8 (4)	C18—C19—H19A	107.9
C9—C8—C17	109.5 (4)	C23—C20—C21	107.9 (6)
C13—C8—C17	108.3 (4)	C23—C20—C19	130.6 (6)
C7—C8—C17	107.7 (4)	C21—C20—C19	121.5 (6)
C10—C9—C8	113.7 (5)	O7—C21—O6	121.6 (8)
C10—C9—H9A	108.8	O7—C21—C20	129.9 (8)
C8—C9—H9A	108.8	O6—C21—C20	108.4 (7)
C10—C9—H9B	108.8	O6—C22—C23	105.1 (6)
C8—C9—H9B	108.8	O6—C22—H22A	110.7
H9A—C9—H9B	107.7	C23—C22—H22A	110.7
C11—C10—C9	112.1 (5)	O6—C22—H22B	110.7

C11—C10—H10A	109.2	C23—C22—H22B	110.7
C9—C10—H10A	109.2	H22A—C22—H22B	108.8
C11—C10—H10B	109.2	C20—C23—C22	109.5 (7)
C9—C10—H10B	109.2	C20—C23—H23A	125.2
H10A—C10—H10B	107.9	C22—C23—H23A	125.2
O2—C11—C10	110.2 (5)	C19—C24—N	111.5 (6)
O2—C11—C6	107.2 (4)	C19—C24—H24A	109.3
C10—C11—C6	114.2 (5)	N—C24—H24A	109.3
O2—C11—H11A	108.4	C19—C24—H24B	109.3
C10—C11—H11A	108.4	N—C24—H24B	109.3
C6—C11—H11A	108.4	H24A—C24—H24B	108.0
C11—O2—C2—O1	-0.7 (10)	C7—C14—C15—C16	55.1 (7)
C11—O2—C2—C1	178.3 (7)	C14—C15—C16—O5	122.3 (7)
C5—O4—C4—O3	-3.5 (9)	C14—C15—C16—C17	-57.0 (7)
C5—O4—C4—C3	175.1 (5)	O5—C16—C17—C18	8.8 (8)
C4—O4—C5—C6	108.6 (6)	C15—C16—C17—C18	-172.0 (5)
O4—C5—C6—C11	75.4 (6)	O5—C16—C17—C8	-120.8 (6)
O4—C5—C6—C12	-44.7 (7)	C15—C16—C17—C8	58.5 (6)
O4—C5—C6—C7	-163.8 (4)	C9—C8—C17—C16	-172.8 (5)
C11—C6—C7—C14	-175.9 (5)	C13—C8—C17—C16	67.1 (6)
C5—C6—C7—C14	60.3 (6)	C7—C8—C17—C16	-56.5 (5)
C12—C6—C7—C14	-58.4 (6)	C9—C8—C17—C18	56.7 (6)
C11—C6—C7—C8	51.4 (5)	C13—C8—C17—C18	-63.4 (6)
C5—C6—C7—C8	-72.4 (6)	C7—C8—C17—C18	173.0 (4)
C12—C6—C7—C8	168.9 (4)	C16—C17—C18—C19	67.9 (6)
C14—C7—C8—C9	174.9 (5)	C8—C17—C18—C19	-164.6 (5)
C6—C7—C8—C9	-51.2 (6)	C17—C18—C19—C24	-173.0 (5)
C14—C7—C8—C13	-63.1 (6)	C17—C18—C19—C20	62.6 (7)
C6—C7—C8—C13	70.8 (6)	C24—C19—C20—C23	-70.8 (8)
C14—C7—C8—C17	57.1 (5)	C18—C19—C20—C23	52.4 (8)
C6—C7—C8—C17	-169.0 (4)	C24—C19—C20—C21	108.2 (7)
C13—C8—C9—C10	-73.4 (7)	C18—C19—C20—C21	-128.6 (7)
C7—C8—C9—C10	50.9 (7)	C22—O6—C21—O7	178.3 (11)
C17—C8—C9—C10	167.6 (5)	C22—O6—C21—C20	2.0 (9)
C8—C9—C10—C11	-56.1 (8)	C23—C20—C21—O7	-176.9 (12)
C2—O2—C11—C10	89.9 (7)	C19—C20—C21—O7	3.9 (16)
C2—O2—C11—C6	-145.2 (5)	C23—C20—C21—O6	-0.9 (9)
C9—C10—C11—O2	177.3 (5)	C19—C20—C21—O6	179.9 (6)
C9—C10—C11—C6	56.5 (7)	C21—O6—C22—C23	-2.2 (9)
C5—C6—C11—O2	-51.7 (6)	C21—C20—C23—C22	-0.5 (8)
C12—C6—C11—O2	68.0 (6)	C19—C20—C23—C22	178.6 (6)
C7—C6—C11—O2	-174.6 (5)	O6—C22—C23—C20	1.7 (8)
C5—C6—C11—C10	70.7 (6)	C20—C19—C24—N	-70.7 (7)
C12—C6—C11—C10	-169.6 (5)	C18—C19—C24—N	164.7 (6)
C7—C6—C11—C10	-52.3 (6)	O8—N—C24—C19	110.7 (14)
C6—C7—C14—C15	167.4 (5)	O9—N—C24—C19	-80.2 (10)
C8—C7—C14—C15	-56.9 (6)		

supplementary materials

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C15—H15A \cdots O8 ⁱ	0.97	2.59	3.312 (18)	131
C22—H22A \cdots O9 ⁱⁱ	0.97	2.47	3.388 (13)	158
C22—H22B \cdots O1 ⁱⁱⁱ	0.97	2.51	3.301 (13)	139
C24—H24B \cdots O3 ^{iv}	0.97	2.56	3.376 (9)	142

Symmetry codes: (i) $x-1, y, z$; (ii) $-x+1, y+1/2, -z+1$; (iii) $-x+1, y-1/2, -z+1$; (iv) $x+1, y, z$.

Fig. 1

