

## Hexaaquazinc(II) bis(4-hydroxybenzenesulfonate) dihydrate

Shan Gao<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>College of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 14 November 2009; accepted 14 November 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.030;  $wR$  factor = 0.082; data-to-parameter ratio = 13.9.

The asymmetric unit of the title hydrated molecular salt,  $[\text{Zn}(\text{H}_2\text{O})_6](\text{C}_6\text{H}_5\text{O}_4\text{S})_2 \cdot 2\text{H}_2\text{O}$ , contains two half-cations, two anions and two uncoordinated water molecules. Both cations are completed by crystallographic inversion symmetry, generating almost regular  $\text{ZnO}_6$  octahedra. In the crystal, the cations, anions and uncoordinated water molecules are linked by  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds, forming a three-dimensional network.

### Related literature

For the isostructural cobalt and nickel analogs of the title compound, see: Du *et al.* (2007) and Kosnic *et al.* (1992), respectively.



### Experimental

#### Crystal data



$M_r = 555.82$

Monoclinic,  $P2_1/c$

$a = 11.7957(5)\text{ \AA}$

$b = 7.2590(4)\text{ \AA}$

$c = 25.3992(11)\text{ \AA}$

$\beta = 94.340(1)^\circ$

$V = 2168.57(18)\text{ \AA}^3$

$Z = 4$

Mo  $K\alpha$  radiation

$\mu = 1.40\text{ mm}^{-1}$

$T = 293\text{ K}$

$0.19 \times 0.19 \times 0.15\text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID IP diffractometer

Absorption correction: multi-scan (*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.777$ ,  $T_{\max} = 0.817$

20669 measured reflections

4940 independent reflections

3319 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.040$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$

$wR(F^2) = 0.082$

$S = 1.03$

4940 reflections

355 parameters

18 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.33\text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.40\text{ e \AA}^{-3}$

**Table 1**  
Selected bond lengths (Å).

Zn1—O2W	2.0654 (16)	Zn2—O4W	2.0763 (15)
Zn1—O3W	2.0692 (16)	Zn2—O6W	2.0780 (16)
Zn1—O1W	2.1087 (15)	Zn2—O5W	2.0849 (16)

**Table 2**  
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1w—H11···O1	0.84 (1)	1.93 (1)	2.763 (2)	169 (3)
O1w—H12···O7w	0.85 (1)	1.86 (1)	2.707 (2)	175 (2)
O2w—H21···O2	0.85 (1)	1.95 (1)	2.792 (2)	172 (3)
O2w—H22···O4 <sup>i</sup>	0.85 (1)	1.92 (1)	2.745 (2)	166 (3)
O3w—H31···O3 <sup>ii</sup>	0.84 (1)	1.97 (1)	2.800 (2)	168 (3)
O3w—H32···O7w <sup>ii</sup>	0.84 (1)	1.95 (1)	2.790 (2)	174 (3)
O4w—H41···O5	0.84 (1)	1.91 (1)	2.742 (2)	174 (3)
O4w—H42···O1w <sup>iii</sup>	0.85 (1)	2.08 (1)	2.914 (2)	167 (3)
O5w—H51···O6	0.85 (1)	1.91 (1)	2.755 (2)	175 (3)
O5w—H52···O8 <sup>iv</sup>	0.85 (1)	1.98 (1)	2.819 (2)	168 (3)
O6w—H61···O8w	0.85 (1)	1.85 (1)	2.698 (2)	170 (3)
O6w—H62···O7 <sup>v</sup>	0.84 (1)	2.03 (1)	2.8401 (19)	163 (3)
O4—H4···O7 <sup>i</sup>	0.85 (1)	1.93 (1)	2.777 (2)	175 (3)
O8—H8···O3 <sup>i</sup>	0.85 (1)	1.95 (1)	2.788 (2)	172 (3)

Symmetry codes: (i)  $-x + 1, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (ii)  $-x + 1, -y + 2, -z + 1$ ; (iii)  $-x + 1, -y + 1, -z + 1$ ; (iv)  $-x, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (v)  $-x, -y + 2, -z + 1$ .

Data collection: *RAPID-AUTO* (Rigaku, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalClear* (Rigaku/MSC, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the Key Project of Natural Science Foundation of Heilongjiang Province (No. ZD200903), the Scientific Fund of Remarkable Teachers of Heilongjiang Province (No. 1054G036), Heilongjiang University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5229).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Du, J.-M., Li, Q., Li, W., Lin, H.-M. & Guo, G.-C. (2007). *Acta Cryst. E63*, m2597.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Kosnic, J., McClymont, E. L., Hodder, R. A. & Squatrito, P. J. (1992). *Inorg. Chim. Acta*, **201**, 143–151.
- Rigaku (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Rigaku/MSC (2002). *CrystalClear*. Rigaku/MSC Inc., The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Westrip, S. P. (2009). *publCIF*. In preparation.

## **supplementary materials**

*Acta Cryst.* (2009). E65, m1633 [doi:10.1107/S1600536809048375]

## Hexaaquazinc(II) bis(4-hydroxybenzenesulfonate) dihydrate

S. Gao and S. W. Ng

### Experimental

One millimolar quantities each of zinc dichloride and *p*-hydroxylbenzenesulfonic acid were dissolved in water; colourless prisms of (I) were isolated after a few days.

### Refinement

Carbon-bound H-atoms were placed in calculated positions ( $C-H = 0.93 \text{ \AA}$ ) and were included in the refinement in the riding model approximation, with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The water H-atoms were located in a difference Fourier map, and were refined with a distance restraint of  $O-H = 0.85 \pm 0.01 \text{ \AA}$ ; their  $U_{\text{iso}}$  values were refined.

### Figures




Fig. 1. The molecular structure of (I) shown at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius. Unlabelled atoms bonded to Zn1 and Zn2 are generated by the symmetry operations  $(1-x, 1-y, 1-z)$  and  $(-x, 1-y, 1-z)$ , respectively.

## Hexaaquazinc(II) bis(4-hydroxybenzenesulfonate) dihydrate

### Crystal data

$[\text{Zn}(\text{H}_2\text{O})_6](\text{C}_6\text{H}_5\text{O}_4\text{S})_2 \cdot 2\text{H}_2\text{O}$	$F_{000} = 1152$
$M_r = 555.82$	$D_x = 1.702 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 12326 reflections
$a = 11.7957 (5) \text{ \AA}$	$\theta = 3.2\text{--}27.5^\circ$
$b = 7.2590 (4) \text{ \AA}$	$\mu = 1.40 \text{ mm}^{-1}$
$c = 25.3992 (11) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 94.340 (1)^\circ$	Prism, colorless
$V = 2168.57 (18) \text{ \AA}^3$	$0.19 \times 0.19 \times 0.15 \text{ mm}$
$Z = 4$	

### Data collection

Rigaku R-AXIS RAPID IP diffractometer	4940 independent reflections
--	------------------------------

# supplementary materials

---

Radiation source: fine-focus sealed tube	3319 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.040$
$T = 293 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
$\omega$ scan	$\theta_{\text{min}} = 3.2^\circ$
Absorption correction: Multi-scan (ABSCOR; Higashi, 1995)	$h = -15 \rightarrow 15$
$T_{\text{min}} = 0.777, T_{\text{max}} = 0.817$	$k = -9 \rightarrow 9$
20669 measured reflections	$l = -29 \rightarrow 32$

## Refinement

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.030$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.082$	$w = 1/[\sigma^2(F_o^2) + (0.0405P)^2 + 0.0996P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\text{max}} = 0.001$
4940 reflections	$\Delta\rho_{\text{max}} = 0.33 \text{ e \AA}^{-3}$
355 parameters	$\Delta\rho_{\text{min}} = -0.40 \text{ e \AA}^{-3}$
18 restraints	Extinction correction: none
Primary atom site location: structure-invariant direct methods	

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn1	0.5000	0.5000	0.5000	0.02424 (10)
Zn2	0.0000	0.5000	0.5000	0.02415 (10)
S1	0.60479 (4)	0.93447 (8)	0.365706 (19)	0.02769 (13)
S2	0.09496 (4)	0.94376 (8)	0.36609 (2)	0.02778 (13)
O1	0.68413 (13)	0.8168 (2)	0.39683 (6)	0.0436 (4)
O2	0.48651 (12)	0.8933 (2)	0.37348 (6)	0.0367 (4)
O3	0.63007 (14)	1.1293 (2)	0.37462 (6)	0.0410 (4)
O4	0.68090 (14)	0.8205 (2)	0.14198 (5)	0.0405 (4)
O5	0.17678 (13)	0.8315 (2)	0.39749 (6)	0.0393 (4)
O6	-0.02227 (12)	0.8971 (2)	0.37348 (6)	0.0385 (4)
O7	0.11697 (13)	1.1408 (2)	0.37451 (5)	0.0359 (4)
O8	0.17676 (14)	0.8359 (2)	0.14229 (6)	0.0412 (4)
O1W	0.64682 (13)	0.6615 (2)	0.49333 (6)	0.0300 (3)

O2W	0.43907 (16)	0.5631 (3)	0.42376 (6)	0.0425 (4)
O3W	0.41407 (15)	0.7258 (2)	0.52682 (7)	0.0412 (4)
O4W	0.15569 (14)	0.5586 (3)	0.47077 (8)	0.0463 (5)
O5W	-0.08685 (14)	0.5987 (2)	0.43118 (6)	0.0361 (4)
O6W	-0.01038 (14)	0.7610 (2)	0.53290 (6)	0.0349 (4)
O7W	0.64137 (15)	0.9889 (2)	0.54413 (7)	0.0371 (4)
O8W	0.16394 (15)	0.9973 (3)	0.52453 (8)	0.0410 (4)
C1	0.62536 (16)	0.8899 (3)	0.29904 (7)	0.0243 (4)
C2	0.73038 (17)	0.8248 (3)	0.28587 (8)	0.0290 (5)
H2	0.7871	0.7971	0.3121	0.035*
C3	0.74935 (17)	0.8018 (3)	0.23319 (8)	0.0296 (5)
H3	0.8194	0.7591	0.2239	0.035*
C4	0.66492 (17)	0.8420 (3)	0.19444 (7)	0.0271 (5)
C5	0.55995 (17)	0.9073 (3)	0.20759 (8)	0.0323 (5)
H5	0.5031	0.9340	0.1813	0.039*
C6	0.54101 (17)	0.9321 (3)	0.26028 (8)	0.0299 (5)
H6	0.4714	0.9771	0.2695	0.036*
C7	0.11767 (16)	0.8998 (3)	0.29945 (8)	0.0254 (4)
C8	0.21941 (18)	0.8220 (3)	0.28634 (8)	0.0315 (5)
H8A	0.2734	0.7847	0.3128	0.038*
C9	0.23990 (18)	0.8004 (3)	0.23389 (8)	0.0329 (5)
H9	0.3080	0.7487	0.2250	0.039*
C10	0.15988 (18)	0.8550 (3)	0.19468 (8)	0.0283 (5)
C11	0.05795 (17)	0.9337 (3)	0.20744 (8)	0.0305 (5)
H11A	0.0040	0.9707	0.1810	0.037*
C12	0.03764 (17)	0.9561 (3)	0.26017 (8)	0.0283 (5)
H12A	-0.0300	1.0092	0.2691	0.034*
H4	0.74446 (14)	0.769 (4)	0.1384 (11)	0.068 (9)*
H8	0.23555 (15)	0.769 (3)	0.1402 (11)	0.060 (9)*
H11	0.659 (2)	0.694 (3)	0.4625 (5)	0.048 (8)*
H12	0.642 (2)	0.762 (2)	0.5103 (9)	0.056 (8)*
H21	0.459 (3)	0.657 (3)	0.4072 (11)	0.084 (11)*
H22	0.3924 (17)	0.494 (3)	0.4062 (9)	0.048 (8)*
H31	0.409 (2)	0.761 (4)	0.5581 (5)	0.054 (8)*
H32	0.393 (2)	0.813 (3)	0.5066 (9)	0.065 (10)*
H41	0.167 (2)	0.639 (3)	0.4480 (8)	0.058 (8)*
H42	0.2171 (14)	0.511 (3)	0.4836 (10)	0.065 (10)*
H51	-0.063 (2)	0.690 (3)	0.4144 (10)	0.073 (10)*
H52	-0.122 (2)	0.518 (3)	0.4119 (9)	0.047 (8)*
H61	0.0480 (18)	0.831 (4)	0.5340 (13)	0.102 (13)*
H62	-0.037 (2)	0.768 (4)	0.5627 (6)	0.072 (10)*
H71	0.7069 (12)	1.027 (4)	0.5550 (10)	0.053 (9)*
H72	0.603 (2)	1.004 (4)	0.5705 (8)	0.084 (12)*
H81	0.205 (2)	1.039 (4)	0.5508 (8)	0.059 (9)*
H82	0.121 (3)	1.077 (4)	0.5089 (13)	0.112 (15)*

## supplementary materials

---

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Zn1	0.02731 (18)	0.0253 (2)	0.02019 (17)	-0.00203 (14)	0.00222 (14)	-0.00013 (14)
Zn2	0.02819 (18)	0.0218 (2)	0.02263 (18)	-0.00079 (14)	0.00322 (14)	0.00139 (14)
S1	0.0340 (3)	0.0312 (3)	0.0185 (3)	-0.0047 (2)	0.0060 (2)	0.0007 (2)
S2	0.0349 (3)	0.0284 (3)	0.0206 (3)	-0.0069 (2)	0.0056 (2)	0.0014 (2)
O1	0.0478 (9)	0.0573 (12)	0.0259 (8)	0.0082 (8)	0.0034 (7)	0.0106 (8)
O2	0.0347 (8)	0.0444 (10)	0.0323 (8)	-0.0053 (7)	0.0119 (7)	0.0013 (7)
O3	0.0598 (10)	0.0359 (10)	0.0293 (8)	-0.0147 (8)	0.0151 (8)	-0.0076 (7)
O4	0.0484 (10)	0.0537 (12)	0.0198 (7)	0.0188 (8)	0.0061 (7)	-0.0021 (7)
O5	0.0461 (9)	0.0444 (11)	0.0272 (8)	-0.0023 (8)	0.0004 (7)	0.0092 (7)
O6	0.0368 (8)	0.0438 (11)	0.0362 (9)	-0.0103 (7)	0.0122 (7)	0.0016 (8)
O7	0.0511 (9)	0.0290 (9)	0.0292 (8)	-0.0110 (7)	0.0131 (7)	-0.0067 (7)
O8	0.0520 (11)	0.0486 (11)	0.0235 (8)	0.0167 (9)	0.0068 (8)	-0.0010 (7)
O1W	0.0377 (8)	0.0267 (9)	0.0262 (8)	-0.0057 (7)	0.0065 (7)	0.0017 (7)
O2W	0.0637 (12)	0.0380 (11)	0.0237 (9)	-0.0182 (9)	-0.0102 (8)	0.0058 (8)
O3W	0.0605 (11)	0.0352 (11)	0.0288 (9)	0.0136 (9)	0.0087 (8)	-0.0025 (8)
O4W	0.0312 (9)	0.0519 (12)	0.0574 (12)	0.0040 (9)	0.0129 (9)	0.0285 (10)
O5W	0.0480 (10)	0.0315 (10)	0.0274 (9)	-0.0078 (8)	-0.0060 (8)	0.0057 (8)
O6W	0.0509 (10)	0.0255 (9)	0.0297 (9)	-0.0042 (8)	0.0112 (8)	-0.0049 (7)
O7W	0.0358 (9)	0.0393 (10)	0.0370 (9)	-0.0047 (8)	0.0078 (8)	-0.0056 (8)
O8W	0.0379 (10)	0.0417 (11)	0.0431 (11)	-0.0049 (8)	0.0017 (9)	-0.0040 (9)
C1	0.0297 (11)	0.0248 (12)	0.0189 (9)	-0.0033 (9)	0.0048 (9)	0.0011 (8)
C2	0.0312 (11)	0.0305 (13)	0.0247 (10)	0.0054 (9)	-0.0023 (9)	-0.0004 (9)
C3	0.0303 (11)	0.0314 (13)	0.0274 (11)	0.0086 (9)	0.0053 (9)	-0.0025 (9)
C4	0.0372 (12)	0.0241 (12)	0.0202 (10)	0.0030 (9)	0.0037 (9)	-0.0026 (9)
C5	0.0308 (11)	0.0413 (15)	0.0241 (11)	0.0051 (10)	-0.0015 (9)	-0.0007 (10)
C6	0.0260 (10)	0.0362 (13)	0.0278 (11)	0.0016 (10)	0.0045 (9)	-0.0003 (9)
C7	0.0308 (11)	0.0221 (12)	0.0235 (10)	-0.0046 (9)	0.0024 (9)	-0.0016 (9)
C8	0.0347 (12)	0.0315 (13)	0.0277 (11)	0.0052 (10)	-0.0019 (10)	0.0004 (9)
C9	0.0343 (12)	0.0332 (13)	0.0316 (12)	0.0089 (10)	0.0050 (10)	-0.0027 (10)
C10	0.0381 (12)	0.0251 (12)	0.0221 (10)	0.0008 (9)	0.0057 (9)	-0.0004 (9)
C11	0.0308 (11)	0.0332 (13)	0.0267 (11)	0.0052 (10)	-0.0036 (9)	0.0004 (9)
C12	0.0273 (11)	0.0285 (12)	0.0296 (12)	0.0000 (9)	0.0046 (9)	-0.0021 (9)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Zn1—O2W	2.0654 (16)	O4W—H41	0.840 (10)
Zn1—O2W <sup>i</sup>	2.0654 (16)	O4W—H42	0.847 (10)
Zn1—O3W	2.0692 (16)	O5W—H51	0.846 (10)
Zn1—O3W <sup>i</sup>	2.0692 (16)	O5W—H52	0.849 (10)
Zn1—O1W <sup>i</sup>	2.1087 (15)	O6W—H61	0.854 (10)
Zn1—O1W	2.1087 (15)	O6W—H62	0.842 (10)
Zn2—O4W <sup>ii</sup>	2.0763 (15)	O7W—H71	0.847 (10)
Zn2—O4W	2.0763 (15)	O7W—H72	0.846 (10)
Zn2—O6W	2.0780 (16)	O8W—H81	0.849 (10)

Zn2—O6W <sup>ii</sup>	2.0780 (16)	O8W—H82	0.848 (10)
Zn2—O5W	2.0849 (16)	C1—C6	1.380 (3)
Zn2—O5W <sup>ii</sup>	2.0849 (16)	C1—C2	1.390 (3)
S1—O2	1.4549 (14)	C2—C3	1.383 (3)
S1—O1	1.4555 (17)	C2—H2	0.9300
S1—O3	1.4598 (17)	C3—C4	1.377 (3)
S1—C1	1.7587 (19)	C3—H3	0.9300
S2—O6	1.4495 (15)	C4—C5	1.390 (3)
S2—O5	1.4542 (16)	C5—C6	1.385 (3)
S2—O7	1.4666 (16)	C5—H5	0.9300
S2—C7	1.763 (2)	C6—H6	0.9300
O4—C4	1.369 (2)	C7—C12	1.382 (3)
O4—H4	0.851 (10)	C7—C8	1.389 (3)
O8—C10	1.367 (2)	C8—C9	1.381 (3)
O8—H8	0.849 (10)	C8—H8A	0.9300
O1W—H11	0.843 (10)	C9—C10	1.378 (3)
O1W—H12	0.849 (10)	C9—H9	0.9300
O2W—H21	0.845 (10)	C10—C11	1.391 (3)
O2W—H22	0.846 (10)	C11—C12	1.388 (3)
O3W—H31	0.840 (10)	C11—H11A	0.9300
O3W—H32	0.841 (10)	C12—H12A	0.9300
O2W—Zn1—O2W <sup>i</sup>	180.0	Zn1—O3W—H31	128.5 (19)
O2W—Zn1—O3W	89.34 (7)	Zn1—O3W—H32	121.7 (19)
O2W <sup>i</sup> —Zn1—O3W	90.66 (7)	H31—O3W—H32	108 (3)
O2W—Zn1—O3W <sup>i</sup>	90.66 (7)	Zn2—O4W—H41	125.1 (18)
O2W <sup>i</sup> —Zn1—O3W <sup>i</sup>	89.34 (7)	Zn2—O4W—H42	122.1 (18)
O3W—Zn1—O3W <sup>i</sup>	180.0	H41—O4W—H42	112 (2)
O2W—Zn1—O1W <sup>i</sup>	88.23 (6)	Zn2—O5W—H51	122 (2)
O2W <sup>i</sup> —Zn1—O1W <sup>i</sup>	91.77 (6)	Zn2—O5W—H52	115.8 (18)
O3W—Zn1—O1W <sup>i</sup>	89.32 (7)	H51—O5W—H52	114 (3)
O3W <sup>i</sup> —Zn1—O1W <sup>i</sup>	90.68 (7)	Zn2—O6W—H61	119 (2)
O2W—Zn1—O1W	91.77 (6)	Zn2—O6W—H62	117 (2)
O2W <sup>i</sup> —Zn1—O1W	88.23 (6)	H61—O6W—H62	107 (3)
O3W—Zn1—O1W	90.68 (7)	H71—O7W—H72	104 (3)
O3W <sup>i</sup> —Zn1—O1W	89.32 (7)	H81—O8W—H82	114 (3)
O1W <sup>i</sup> —Zn1—O1W	180.0	C6—C1—C2	120.75 (18)
O4W <sup>ii</sup> —Zn2—O4W	180.0	C6—C1—S1	120.10 (15)
O4W <sup>ii</sup> —Zn2—O6W	87.85 (7)	C2—C1—S1	118.98 (16)
O4W—Zn2—O6W	92.15 (7)	C3—C2—C1	119.08 (19)
O4W <sup>ii</sup> —Zn2—O6W <sup>ii</sup>	92.15 (7)	C3—C2—H2	120.5
O4W—Zn2—O6W <sup>ii</sup>	87.85 (7)	C1—C2—H2	120.5
O6W—Zn2—O6W <sup>ii</sup>	180.0	C4—C3—C2	120.31 (18)
O4W <sup>ii</sup> —Zn2—O5W	88.76 (7)	C4—C3—H3	119.8
O4W—Zn2—O5W	91.24 (7)	C2—C3—H3	119.8
O6W—Zn2—O5W	89.07 (7)	O4—C4—C3	121.75 (18)

## supplementary materials

---

O6W <sup>ii</sup> —Zn2—O5W	90.93 (7)	O4—C4—C5	117.61 (19)
O4W <sup>ii</sup> —Zn2—O5W <sup>ii</sup>	91.24 (7)	C3—C4—C5	120.64 (18)
O4W—Zn2—O5W <sup>ii</sup>	88.76 (7)	C6—C5—C4	119.2 (2)
O6W—Zn2—O5W <sup>ii</sup>	90.93 (7)	C6—C5—H5	120.4
O6W <sup>ii</sup> —Zn2—O5W <sup>ii</sup>	89.07 (7)	C4—C5—H5	120.4
O5W—Zn2—O5W <sup>ii</sup>	180.0	C1—C6—C5	120.01 (19)
O2—S1—O1	112.93 (10)	C1—C6—H6	120.0
O2—S1—O3	111.38 (10)	C5—C6—H6	120.0
O1—S1—O3	111.66 (10)	C12—C7—C8	120.14 (19)
O2—S1—C1	107.25 (9)	C12—C7—S2	119.64 (15)
O1—S1—C1	106.56 (9)	C8—C7—S2	120.03 (16)
O3—S1—C1	106.63 (9)	C9—C8—C7	119.7 (2)
O6—S2—O5	113.46 (9)	C9—C8—H8A	120.2
O6—S2—O7	111.65 (9)	C7—C8—H8A	120.2
O5—S2—O7	111.33 (10)	C10—C9—C8	120.29 (19)
O6—S2—C7	107.33 (9)	C10—C9—H9	119.9
O5—S2—C7	106.41 (9)	C8—C9—H9	119.9
O7—S2—C7	106.16 (9)	O8—C10—C9	122.13 (18)
C4—O4—H4	109.9 (19)	O8—C10—C11	117.43 (19)
C10—O8—H8	107.3 (19)	C9—C10—C11	120.44 (18)
Zn1—O1W—H11	115.8 (17)	C10—C11—C12	119.2 (2)
Zn1—O1W—H12	110.2 (17)	C10—C11—H11A	120.4
H11—O1W—H12	105 (2)	C12—C11—H11A	120.4
Zn1—O2W—H21	124 (2)	C7—C12—C11	120.23 (19)
Zn1—O2W—H22	121.9 (19)	C7—C12—H12A	119.9
H21—O2W—H22	114 (3)	C11—C12—H12A	119.9
O2—S1—C1—C6	37.7 (2)	O6—S2—C7—C12	44.6 (2)
O1—S1—C1—C6	158.94 (18)	O5—S2—C7—C12	166.37 (17)
O3—S1—C1—C6	-81.67 (19)	O7—S2—C7—C12	-74.94 (19)
O2—S1—C1—C2	-146.98 (17)	O6—S2—C7—C8	-140.39 (17)
O1—S1—C1—C2	-25.8 (2)	O5—S2—C7—C8	-18.6 (2)
O3—S1—C1—C2	93.60 (18)	O7—S2—C7—C8	100.09 (18)
C6—C1—C2—C3	-0.3 (3)	C12—C7—C8—C9	-0.3 (3)
S1—C1—C2—C3	-175.53 (16)	S2—C7—C8—C9	-175.31 (17)
C1—C2—C3—C4	-0.4 (3)	C7—C8—C9—C10	-0.2 (3)
C2—C3—C4—O4	-179.9 (2)	C8—C9—C10—O8	-179.9 (2)
C2—C3—C4—C5	0.5 (3)	C8—C9—C10—C11	0.4 (3)
O4—C4—C5—C6	-179.5 (2)	O8—C10—C11—C12	-179.8 (2)
C3—C4—C5—C6	0.1 (3)	C9—C10—C11—C12	-0.1 (3)
C2—C1—C6—C5	0.9 (3)	C8—C7—C12—C11	0.6 (3)
S1—C1—C6—C5	176.08 (17)	S2—C7—C12—C11	175.62 (17)
C4—C5—C6—C1	-0.8 (3)	C10—C11—C12—C7	-0.4 (3)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $-x, -y+1, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O1w—H11 $\cdots$ O1	0.84 (1)	1.93 (1)	2.763 (2)	169 (3)

## supplementary materials

---

O1w—H12···O7w	0.85 (1)	1.86 (1)	2.707 (2)	175 (2)
O2w—H21···O2	0.85 (1)	1.95 (1)	2.792 (2)	172 (3)
O2w—H22···O4 <sup>iii</sup>	0.85 (1)	1.92 (1)	2.745 (2)	166 (3)
O3w—H31···O3 <sup>iv</sup>	0.84 (1)	1.97 (1)	2.800 (2)	168 (3)
O3w—H32···O7w <sup>iv</sup>	0.84 (1)	1.95 (1)	2.790 (2)	174 (3)
O4w—H41···O5	0.84 (1)	1.91 (1)	2.742 (2)	174 (3)
O4w—H42···O1w <sup>i</sup>	0.85 (1)	2.08 (1)	2.914 (2)	167 (3)
O5w—H51···O6	0.85 (1)	1.91 (1)	2.755 (2)	175 (3)
O5w—H52···O8 <sup>v</sup>	0.85 (1)	1.98 (1)	2.819 (2)	168 (3)
O6w—H61···O8w	0.85 (1)	1.85 (1)	2.698 (2)	170 (3)
O6w—H62···O7 <sup>vi</sup>	0.84 (1)	2.03 (1)	2.8401 (19)	163 (3)
O4—H4···O7 <sup>iii</sup>	0.85 (1)	1.93 (1)	2.777 (2)	175 (3)
O8—H8···O3 <sup>iii</sup>	0.85 (1)	1.95 (1)	2.788 (2)	172 (3)

Symmetry codes: (iii)  $-x+1, y-1/2, -z+1/2$ ; (iv)  $-x+1, -y+2, -z+1$ ; (i)  $-x+1, -y+1, -z+1$ ; (v)  $-x, y-1/2, -z+1/2$ ; (vi)  $-x, -y+2, -z+1$ .

## supplementary materials

---

Fig. 1

