

## Dichloridobis(4-chlorobenzyl)(4,4'-dimethyl-2,2'-bipyridine- $\kappa^2 N,N'$ )tin(IV)

Thy Chun Keng, Kong Mun Lo and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 18 November 2009; accepted 18 November 2009

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  
 $R$  factor = 0.031;  $wR$  factor = 0.095; data-to-parameter ratio = 20.9.

The  $\text{Sn}^{IV}$  atom in the title compound,  $[\text{Sn}(\text{C}_7\text{H}_6\text{Cl})_2\text{Cl}_2(\text{C}_{12}\text{H}_{12}\text{N}_2)]$ , is coordinated by the bidentate *N*-heterocycle molecule, two chlorobenzyl anions and two  $\text{Cl}^-$  anions in a distorted *trans*- $\text{C}_2\text{SnCl}_2\text{N}_2$  octahedral geometry [ $\text{C}-\text{Sn}-\text{C} = 178.4(1)^\circ$ ]. In the molecular structure, the two benzene rings are oriented at a dihedral angle of  $39.62(17)^\circ$ .

### Related literature

For the synthesis of bis(4-chlorobenzyl)tin dichloride, see: Sisido *et al.* (1961). For the 1,10-phenanthroline adduct, see: Tan *et al.* (2009).



### Experimental

#### Crystal data

$[\text{Sn}(\text{C}_7\text{H}_6\text{Cl})_2\text{Cl}_2(\text{C}_{12}\text{H}_{12}\text{N}_2)]$	$V = 2772.6(3)\text{ \AA}^3$
$M_r = 624.96$	$Z = 4$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
$a = 11.4035(6)\text{ \AA}$	$\mu = 1.32\text{ mm}^{-1}$
$b = 14.6804(8)\text{ \AA}$	$T = 293\text{ K}$
$c = 16.9270(9)\text{ \AA}$	$0.30 \times 0.28 \times 0.20\text{ mm}$
$\beta = 101.9182(9)^\circ$	

#### Data collection

Bruker SMART APEX	15766 measured reflections
diffractometer	6263 independent reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	4883 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.017$
	$T_{\min} = 0.692$ , $T_{\max} = 0.778$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$	300 parameters
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\max} = 0.78\text{ e \AA}^{-3}$
6263 reflections	$\Delta\rho_{\min} = -0.49\text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the University of Malaya (grant No. RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2682).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Sisido, K., Takeda, Y. & Kinugawa, Z. (1961). *J. Am. Chem. Soc.* **83**, 538–541.  
Tan, C. L., Lo, K. M. & Ng, S. W. (2009). *Acta Cryst. E* **65**, m717.  
Westrip, S. P. (2009). *publCIF*. In preparation.

## **supplementary materials**

*Acta Cryst.* (2009). E65, m1676 [doi:10.1107/S1600536809049381]

## Dichloridobis(4-chlorobenzyl)(4,4'-dimethyl-2,2'-bipyridine- $\kappa^2N,N'$ )tin(IV)

T. C. Keng, K. M. Lo and S. W. Ng

### Experimental

Di(4-chlorobenzyl)tin dichloride was synthesized by the reaction of 4-chlorobenzyl chloride and metallic tin (Sisido *et al.*). The diorganotin compound (0.44 g, 1 mmol) and 4,4'-dimethyl-2,2'-bipyridine (0.18 g, 1 mmol) was heated in chloroform (30 ml) for 1 h. Faint-yellow crystals separated upon slow evaporation of the solvent.

### Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.93–0.97 Å) and were included in the refinement in the riding model approximation, with  $U(H)$  set to 1.2–1.5 $U(C)$ .

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{SnCl}_2(\text{C}_{10}\text{H}_{12}\text{N}_2)(\text{C}_7\text{H}_6\text{Cl})_2$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

## Dichloridobis(4-chlorobenzyl)(4,4'-dimethyl-2,2'-bipyridine- $\kappa^2N,N'$ )tin(IV)

### Crystal data

$[\text{Sn}(\text{C}_7\text{H}_6\text{Cl})_2\text{Cl}_2(\text{C}_{12}\text{H}_{12}\text{N}_2)]$	$F(000) = 1248$
$M_r = 624.96$	$D_x = 1.497 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 6177 reflections
$a = 11.4035 (6) \text{ \AA}$	$\theta = 2.4\text{--}28.3^\circ$
$b = 14.6804 (8) \text{ \AA}$	$\mu = 1.32 \text{ mm}^{-1}$
$c = 16.9270 (9) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 101.9182 (9)^\circ$	Block, pale yellow
$V = 2772.6 (3) \text{ \AA}^3$	$0.30 \times 0.28 \times 0.20 \text{ mm}$
$Z = 4$	

### Data collection

Bruker SMART APEX diffractometer	6263 independent reflections
Radiation source: fine-focus sealed tube	4883 reflections with $I > 2\sigma(I)$

# supplementary materials

---

graphite	$R_{\text{int}} = 0.017$
$\omega$ scans	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	$h = -14 \rightarrow 14$
$T_{\text{min}} = 0.692, T_{\text{max}} = 0.778$	$k = -19 \rightarrow 18$
15766 measured reflections	$l = -21 \rightarrow 17$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.031$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.095$	H-atom parameters constrained
$S = 1.00$	$w = 1/[\sigma^2(F_o^2) + (0.0523P)^2 + 1.1119P]$ where $P = (F_o^2 + 2F_c^2)/3$
6263 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
300 parameters	$\Delta\rho_{\text{max}} = 0.78 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.49 \text{ e \AA}^{-3}$

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.278374 (15)	0.231209 (14)	0.536683 (11)	0.05372 (8)
Cl1	0.06174 (7)	0.19394 (9)	0.52457 (7)	0.0993 (3)
Cl2	0.28579 (8)	0.31620 (8)	0.40783 (5)	0.0894 (3)
Cl3	0.35723 (13)	-0.23372 (9)	0.67467 (9)	0.1119 (4)
Cl4	0.68248 (17)	0.53534 (12)	0.79043 (9)	0.1696 (8)
N1	0.4862 (2)	0.24492 (17)	0.57811 (15)	0.0558 (5)
N2	0.34094 (19)	0.16803 (15)	0.66356 (12)	0.0513 (5)
C1	0.3049 (3)	0.1063 (2)	0.47620 (17)	0.0686 (8)
H1A	0.2382	0.0985	0.4309	0.082*
H1B	0.3768	0.1127	0.4545	0.082*
C2	0.3163 (3)	0.0226 (2)	0.52556 (17)	0.0606 (7)
C3	0.2208 (3)	-0.0357 (2)	0.5251 (2)	0.0721 (8)
H3	0.1468	-0.0219	0.4926	0.087*
C4	0.2322 (3)	-0.1133 (3)	0.5713 (2)	0.0788 (9)
H4	0.1670	-0.1518	0.5696	0.095*
C5	0.3412 (3)	-0.1333 (2)	0.6199 (2)	0.0726 (8)
C6	0.4370 (3)	-0.0777 (2)	0.62247 (19)	0.0680 (8)
H6	0.5105	-0.0920	0.6555	0.082*
C7	0.4249 (3)	-0.0004 (2)	0.57607 (18)	0.0639 (7)
H7	0.4907	0.0377	0.5784	0.077*
C8	0.2463 (3)	0.3566 (2)	0.5963 (2)	0.0728 (8)
H8A	0.2064	0.3988	0.5553	0.087*
H8B	0.1911	0.3434	0.6313	0.087*
C9	0.3519 (3)	0.4035 (2)	0.64484 (19)	0.0664 (8)

C10	0.3852 (3)	0.3904 (2)	0.7278 (2)	0.0735 (8)
H10	0.3389	0.3529	0.7535	0.088*
C11	0.4853 (4)	0.4318 (3)	0.7727 (2)	0.0858 (11)
H11	0.5059	0.4230	0.8282	0.103*
C12	0.5538 (4)	0.4856 (3)	0.7352 (2)	0.0943 (12)
C13	0.5227 (4)	0.5011 (3)	0.6537 (2)	0.0985 (13)
H13	0.5691	0.5394	0.6288	0.118*
C14	0.4235 (4)	0.4599 (2)	0.6093 (2)	0.0797 (9)
H14	0.4035	0.4699	0.5539	0.096*
C15	0.5562 (3)	0.2800 (2)	0.5314 (2)	0.0705 (9)
H15	0.5227	0.2934	0.4778	0.085*
C16	0.6750 (3)	0.2967 (3)	0.5597 (2)	0.0813 (10)
H16	0.7213	0.3205	0.5254	0.098*
C17	0.7262 (3)	0.2785 (3)	0.6385 (3)	0.0825 (10)
C18	0.6537 (3)	0.2412 (2)	0.6866 (2)	0.0689 (8)
H18	0.6858	0.2277	0.7404	0.083*
C19	0.5348 (2)	0.22410 (18)	0.65515 (16)	0.0516 (6)
C20	0.8561 (4)	0.2981 (5)	0.6727 (3)	0.139 (2)
H20A	0.8984	0.3037	0.6295	0.209*
H20B	0.8899	0.2492	0.7077	0.209*
H20C	0.8629	0.3540	0.7028	0.209*
C21	0.4549 (2)	0.17973 (18)	0.70150 (15)	0.0528 (6)
C22	0.4934 (3)	0.1484 (2)	0.77971 (18)	0.0733 (9)
H22	0.5725	0.1579	0.8060	0.088*
C23	0.4162 (4)	0.1035 (3)	0.81914 (19)	0.0831 (10)
C24	0.3023 (3)	0.0899 (2)	0.7781 (2)	0.0772 (9)
H24	0.2483	0.0580	0.8019	0.093*
C25	0.2669 (3)	0.1234 (2)	0.70125 (18)	0.0632 (7)
H25	0.1879	0.1147	0.6744	0.076*
C26	0.4588 (5)	0.0702 (4)	0.9050 (2)	0.138 (2)
H26A	0.4559	0.0049	0.9060	0.206*
H26B	0.4079	0.0946	0.9385	0.206*
H26C	0.5397	0.0902	0.9249	0.206*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sn1	0.04228 (11)	0.06803 (14)	0.04599 (12)	0.00021 (8)	-0.00217 (8)	0.00089 (8)
Cl1	0.0438 (4)	0.1234 (8)	0.1241 (8)	-0.0074 (4)	0.0018 (4)	-0.0088 (7)
Cl2	0.0821 (6)	0.1201 (8)	0.0552 (5)	-0.0123 (5)	-0.0105 (4)	0.0282 (5)
Cl3	0.1097 (9)	0.1036 (8)	0.1256 (10)	-0.0010 (6)	0.0316 (8)	0.0320 (7)
Cl4	0.2037 (16)	0.1603 (14)	0.1127 (10)	-0.1049 (13)	-0.0418 (10)	0.0234 (9)
N1	0.0450 (12)	0.0726 (15)	0.0471 (13)	-0.0018 (10)	0.0032 (10)	0.0021 (10)
N2	0.0498 (11)	0.0597 (13)	0.0425 (11)	-0.0029 (9)	0.0052 (9)	-0.0037 (10)
C1	0.0699 (18)	0.087 (2)	0.0462 (15)	0.0005 (16)	0.0043 (13)	-0.0122 (15)
C2	0.0583 (15)	0.0721 (19)	0.0503 (15)	0.0011 (13)	0.0084 (12)	-0.0151 (13)
C3	0.0534 (17)	0.088 (2)	0.0686 (19)	-0.0050 (15)	-0.0018 (14)	-0.0151 (18)
C4	0.0645 (19)	0.085 (2)	0.086 (2)	-0.0156 (17)	0.0133 (17)	-0.012 (2)

## supplementary materials

---

C5	0.074 (2)	0.076 (2)	0.071 (2)	-0.0009 (16)	0.0231 (16)	-0.0028 (17)
C6	0.0580 (17)	0.082 (2)	0.0625 (18)	0.0071 (15)	0.0097 (14)	-0.0047 (16)
C7	0.0520 (15)	0.079 (2)	0.0605 (17)	-0.0022 (13)	0.0127 (13)	-0.0135 (15)
C8	0.0664 (18)	0.072 (2)	0.077 (2)	0.0131 (15)	0.0066 (16)	-0.0032 (17)
C9	0.078 (2)	0.0556 (16)	0.0646 (18)	0.0083 (14)	0.0125 (15)	-0.0035 (14)
C10	0.094 (2)	0.0649 (19)	0.0637 (19)	0.0016 (17)	0.0216 (17)	-0.0017 (15)
C11	0.123 (3)	0.074 (2)	0.0553 (18)	-0.008 (2)	0.0051 (19)	0.0007 (16)
C12	0.123 (3)	0.076 (2)	0.073 (2)	-0.032 (2)	-0.005 (2)	0.0002 (19)
C13	0.133 (4)	0.079 (2)	0.079 (2)	-0.038 (2)	0.012 (2)	0.009 (2)
C14	0.104 (3)	0.069 (2)	0.0620 (19)	-0.0090 (19)	0.0083 (18)	0.0058 (16)
C15	0.0569 (17)	0.100 (3)	0.0549 (17)	-0.0075 (15)	0.0106 (14)	0.0061 (16)
C16	0.0580 (18)	0.111 (3)	0.079 (2)	-0.0180 (18)	0.0228 (17)	-0.002 (2)
C17	0.0478 (17)	0.110 (3)	0.085 (3)	-0.0116 (16)	0.0041 (16)	-0.013 (2)
C18	0.0496 (16)	0.091 (2)	0.0591 (18)	-0.0032 (14)	-0.0051 (13)	-0.0041 (16)
C19	0.0464 (13)	0.0581 (15)	0.0456 (14)	0.0008 (11)	-0.0012 (11)	-0.0047 (11)
C20	0.054 (2)	0.222 (6)	0.133 (4)	-0.037 (3)	-0.001 (2)	-0.004 (4)
C21	0.0567 (14)	0.0535 (15)	0.0434 (13)	-0.0033 (11)	-0.0006 (11)	-0.0032 (11)
C22	0.080 (2)	0.076 (2)	0.0520 (17)	-0.0178 (16)	-0.0151 (14)	0.0107 (15)
C23	0.112 (3)	0.079 (2)	0.0510 (17)	-0.026 (2)	-0.0015 (18)	0.0113 (16)
C24	0.101 (3)	0.074 (2)	0.0589 (18)	-0.0274 (18)	0.0211 (18)	-0.0006 (16)
C25	0.0643 (17)	0.0681 (18)	0.0575 (17)	-0.0124 (14)	0.0131 (13)	-0.0064 (14)
C26	0.184 (5)	0.145 (5)	0.066 (2)	-0.062 (4)	-0.016 (3)	0.040 (3)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Sn1—C1	2.151 (3)	C10—H10	0.9300
Sn1—C8	2.166 (3)	C11—C12	1.357 (5)
Sn1—N2	2.313 (2)	C11—H11	0.9300
Sn1—N1	2.337 (2)	C12—C13	1.371 (5)
Sn1—Cl1	2.4970 (9)	C13—C14	1.363 (5)
Sn1—Cl2	2.5293 (8)	C13—H13	0.9300
Cl3—C5	1.731 (4)	C14—H14	0.9300
Cl4—C12	1.730 (4)	C15—C16	1.363 (5)
N1—C19	1.343 (4)	C15—H15	0.9300
N1—C15	1.337 (4)	C16—C17	1.366 (5)
N2—C25	1.331 (4)	C16—H16	0.9300
N2—C21	1.336 (3)	C17—C18	1.388 (5)
C1—C2	1.476 (5)	C17—C20	1.502 (5)
C1—H1A	0.9700	C18—C19	1.374 (4)
C1—H1B	0.9700	C18—H18	0.9300
C2—C3	1.384 (4)	C19—C21	1.472 (4)
C2—C7	1.394 (4)	C20—H20A	0.9600
C3—C4	1.372 (5)	C20—H20B	0.9600
C3—H3	0.9300	C20—H20C	0.9600
C4—C5	1.374 (5)	C21—C22	1.384 (4)
C4—H4	0.9300	C22—C23	1.378 (5)
C5—C6	1.358 (5)	C22—H22	0.9300
C6—C7	1.371 (5)	C23—C24	1.356 (5)
C6—H6	0.9300	C23—C26	1.515 (5)

C7—H7	0.9300	C24—C25	1.371 (4)
C8—C9	1.480 (5)	C24—H24	0.9300
C8—H8A	0.9700	C25—H25	0.9300
C8—H8B	0.9700	C26—H26A	0.9600
C9—C10	1.391 (5)	C26—H26B	0.9600
C9—C14	1.385 (5)	C26—H26C	0.9600
C10—C11	1.375 (5)		
C1—Sn1—C8	178.42 (12)	C9—C10—H10	119.4
C1—Sn1—N2	93.06 (10)	C12—C11—C10	119.4 (3)
C8—Sn1—N2	87.58 (11)	C12—C11—H11	120.3
C1—Sn1—N1	88.99 (11)	C10—C11—H11	120.3
C8—Sn1—N1	92.59 (11)	C11—C12—C13	120.9 (4)
N2—Sn1—N1	69.92 (8)	C11—C12—Cl4	119.9 (3)
C1—Sn1—Cl1	90.48 (9)	C13—C12—Cl4	119.2 (3)
C8—Sn1—Cl1	88.01 (9)	C14—C13—C12	119.5 (4)
N2—Sn1—Cl1	95.66 (6)	C14—C13—H13	120.2
N1—Sn1—Cl1	165.52 (7)	C12—C13—H13	120.2
C1—Sn1—Cl2	88.68 (9)	C13—C14—C9	121.5 (3)
C8—Sn1—Cl2	91.21 (10)	C13—C14—H14	119.2
N2—Sn1—Cl2	160.01 (6)	C9—C14—H14	119.2
N1—Sn1—Cl2	90.22 (6)	N1—C15—C16	122.2 (3)
Cl1—Sn1—Cl2	104.24 (4)	N1—C15—H15	118.9
C19—N1—C15	119.0 (3)	C16—C15—H15	118.9
C19—N1—Sn1	117.50 (18)	C15—C16—C17	120.1 (3)
C15—N1—Sn1	123.2 (2)	C15—C16—H16	120.0
C25—N2—C21	118.9 (2)	C17—C16—H16	120.0
C25—N2—Sn1	122.54 (19)	C16—C17—C18	117.6 (3)
C21—N2—Sn1	118.53 (17)	C16—C17—C20	121.8 (4)
C2—C1—Sn1	116.38 (19)	C18—C17—C20	120.5 (4)
C2—C1—H1A	108.2	C19—C18—C17	120.3 (3)
Sn1—C1—H1A	108.2	C19—C18—H18	119.9
C2—C1—H1B	108.2	C17—C18—H18	119.9
Sn1—C1—H1B	108.2	N1—C19—C18	120.8 (3)
H1A—C1—H1B	107.3	N1—C19—C21	116.2 (2)
C3—C2—C7	116.8 (3)	C18—C19—C21	123.0 (3)
C3—C2—C1	122.4 (3)	C17—C20—H20A	109.5
C7—C2—C1	120.8 (3)	C17—C20—H20B	109.5
C4—C3—C2	122.0 (3)	H20A—C20—H20B	109.5
C4—C3—H3	119.0	C17—C20—H20C	109.5
C2—C3—H3	119.0	H20A—C20—H20C	109.5
C3—C4—C5	119.1 (3)	H20B—C20—H20C	109.5
C3—C4—H4	120.4	N2—C21—C22	120.1 (3)
C5—C4—H4	120.4	N2—C21—C19	116.6 (2)
C6—C5—C4	120.9 (3)	C22—C21—C19	123.2 (3)
C6—C5—Cl3	120.0 (3)	C23—C22—C21	120.9 (3)
C4—C5—Cl3	119.1 (3)	C23—C22—H22	119.5
C5—C6—C7	119.6 (3)	C21—C22—H22	119.5
C5—C6—H6	120.2	C24—C23—C22	117.5 (3)
C7—C6—H6	120.2	C24—C23—C26	121.9 (4)

## supplementary materials

---

C6—C7—C2	121.6 (3)	C22—C23—C26	120.6 (4)
C6—C7—H7	119.2	C23—C24—C25	119.8 (3)
C2—C7—H7	119.2	C23—C24—H24	120.1
C9—C8—Sn1	117.3 (2)	C25—C24—H24	120.1
C9—C8—H8A	108.0	N2—C25—C24	122.7 (3)
Sn1—C8—H8A	108.0	N2—C25—H25	118.7
C9—C8—H8B	108.0	C24—C25—H25	118.7
Sn1—C8—H8B	108.0	C23—C26—H26A	109.5
H8A—C8—H8B	107.2	C23—C26—H26B	109.5
C10—C9—C14	117.3 (3)	H26A—C26—H26B	109.5
C10—C9—C8	121.2 (3)	C23—C26—H26C	109.5
C14—C9—C8	121.4 (3)	H26A—C26—H26C	109.5
C11—C10—C9	121.3 (3)	H26B—C26—H26C	109.5
C11—C10—H10	119.4		
C1—Sn1—N1—C15	82.9 (3)	Sn1—C8—C9—C10	−95.9 (3)
N2—Sn1—N1—C15	176.5 (3)	C14—C9—C10—C11	−0.1 (5)
Cl1—Sn1—N1—C15	170.9 (2)	C8—C9—C10—C11	178.0 (3)
Cl2—Sn1—N1—C15	−5.8 (3)	C9—C10—C11—C12	−0.8 (6)
C1—Sn1—N1—C19	−103.4 (2)	C10—C11—C12—C13	1.7 (7)
C8—Sn1—N1—C19	76.7 (2)	C10—C11—C12—Cl4	−178.0 (3)
N2—Sn1—N1—C19	−9.79 (19)	C11—C12—C13—C14	−1.8 (7)
Cl1—Sn1—N1—C19	−15.4 (4)	Cl4—C12—C13—C14	178.0 (4)
Cl2—Sn1—N1—C19	167.9 (2)	C12—C13—C14—C9	0.9 (7)
C1—Sn1—N2—C25	−86.3 (2)	C10—C9—C14—C13	0.0 (6)
C8—Sn1—N2—C25	92.2 (2)	C8—C9—C14—C13	−178.1 (4)
N1—Sn1—N2—C25	−174.1 (2)	C19—N1—C15—C16	−1.1 (5)
Cl1—Sn1—N2—C25	4.5 (2)	Sn1—N1—C15—C16	172.5 (3)
Cl2—Sn1—N2—C25	179.13 (18)	N1—C15—C16—C17	−0.7 (6)
C1—Sn1—N2—C21	96.2 (2)	C15—C16—C17—C18	1.4 (6)
C8—Sn1—N2—C21	−85.3 (2)	C15—C16—C17—C20	−178.4 (4)
N1—Sn1—N2—C21	8.36 (19)	C16—C17—C18—C19	−0.3 (6)
Cl1—Sn1—N2—C21	−173.05 (19)	C20—C17—C18—C19	179.5 (4)
Cl2—Sn1—N2—C21	1.6 (3)	C15—N1—C19—C18	2.2 (4)
C8—Sn1—C1—C2	−97 (5)	Sn1—N1—C19—C18	−171.8 (2)
N2—Sn1—C1—C2	16.6 (2)	C15—N1—C19—C21	−175.8 (3)
N1—Sn1—C1—C2	86.5 (2)	Sn1—N1—C19—C21	10.3 (3)
Cl1—Sn1—C1—C2	−79.0 (2)	C17—C18—C19—N1	−1.5 (5)
Cl2—Sn1—C1—C2	176.7 (2)	C17—C18—C19—C21	176.3 (3)
Sn1—C1—C2—C3	97.9 (3)	C25—N2—C21—C22	−2.2 (4)
Sn1—C1—C2—C7	−81.2 (3)	Sn1—N2—C21—C22	175.5 (2)
C7—C2—C3—C4	−0.8 (5)	C25—N2—C21—C19	176.1 (2)
C1—C2—C3—C4	−179.9 (3)	Sn1—N2—C21—C19	−6.3 (3)
C2—C3—C4—C5	0.6 (5)	N1—C19—C21—N2	−2.7 (4)
C3—C4—C5—C6	−0.3 (5)	C18—C19—C21—N2	179.4 (3)
C3—C4—C5—Cl3	−177.1 (3)	N1—C19—C21—C22	175.5 (3)
C4—C5—C6—C7	0.3 (5)	C18—C19—C21—C22	−2.4 (5)
Cl3—C5—C6—C7	177.0 (2)	N2—C21—C22—C23	1.2 (5)
C5—C6—C7—C2	−0.5 (5)	C19—C21—C22—C23	−176.9 (3)
C3—C2—C7—C6	0.7 (4)	C21—C22—C23—C24	1.0 (6)

## supplementary materials

---

C1—C2—C7—C6	179.9 (3)	C21—C22—C23—C26	-179.4 (4)
N2—Sn1—C8—C9	65.4 (3)	C22—C23—C24—C25	-2.2 (6)
N1—Sn1—C8—C9	-4.4 (3)	C26—C23—C24—C25	178.2 (4)
Cl1—Sn1—C8—C9	161.1 (3)	C21—N2—C25—C24	0.9 (4)
Cl2—Sn1—C8—C9	-94.6 (3)	Sn1—N2—C25—C24	-176.6 (2)
Sn1—C8—C9—C14	82.1 (4)	C23—C24—C25—N2	1.4 (5)

## supplementary materials

---

Fig. 1

