

2-(4-Chlorophenyl)imidazo[1,2-a]-pyridine-3-carbaldehyde

Yun-Hui Li,^{a*} Wen-Yu Liu,^a Yang Gao^b and Yu-Peng Wang^a

^aSchool of Chemistry and Environmental Engineering, Changchun University of Science and Technology, Changchun 130022, People's Republic of China, and ^bJilin College of Transportation Vocation and Technology, Changchun 130012, People's Republic of China

Correspondence e-mail: lyunhui@163.com


Received 31 October 2009; accepted 14 November 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.036; wR factor = 0.089; data-to-parameter ratio = 12.9.

The asymmetric unit of the title compound, $\text{C}_{14}\text{H}_9\text{ClN}_2\text{O}$, contains two molecules with dihedral angles of 33.52 (11) and 34.58 (11)° between their benzene rings and imidazole ring systems. In the crystal, $\text{C}-\text{H}\cdots\text{N}$ and $\text{C}-\text{H}\cdots\text{O}$ interactions are observed. The crystal examined was found to be a racemic twin.

Related literature

For the synthesis, see: Burkholder *et al.* (2001).



Experimental

Crystal data

$\text{C}_{14}\text{H}_9\text{ClN}_2\text{O}$

$M_r = 256.68$

Orthorhombic, $Pna2_1$
 $a = 21.3367$ (13) Å
 $b = 7.2391$ (4) Å
 $c = 15.1748$ (10) Å
 $V = 2343.9$ (2) Å³

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.31$ mm⁻¹
 $T = 293$ K
 $0.35 \times 0.29 \times 0.28$ mm

Data collection

Bruker SMART APEX CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2002)
 $T_{\min} = 0.870$, $T_{\max} = 0.894$

12428 measured reflections
 4198 independent reflections
 3564 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.089$
 $S = 1.03$
 4198 reflections
 325 parameters
 1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.22$ e Å⁻³
 $\Delta\rho_{\min} = -0.18$ e Å⁻³
 Absolute structure: Flack (1983),
 1800 Friedel pairs
 Flack parameter: 0.40 (5)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C11}-\text{H11}\cdots\text{O2}^i$	0.93	2.54	3.442 (3)	163
$\text{C12}-\text{H12}\cdots\text{N4}^{ii}$	0.93	2.59	3.518 (4)	172

Symmetry codes: (i) $x + \frac{1}{2}, -y + \frac{3}{2}, z$; (ii) $-x + \frac{3}{2}, y - \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 2002); cell refinement: *SAINT* (Bruker, 2002); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Changchun University of Science and Technology for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5204).

References

- Bruker (2002). *SMART, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Burkholder, C., Dolbier, W. R., Medebielle, M. & Ait-Mohand, S. (2001). *Tetrahedron Lett.* **42**, 3077–3080.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, o3192 [doi:10.1107/S1600536809048302]

2-(4-Chlorophenyl)imidazo[1,2-*a*]pyridine-3-carbaldehyde

Y.-H. Li, W.-Y. Liu, Y. Gao and Y.-P. Wang

Experimental

To a solution of 2.0 mmol 2-(4-chlorophenyl)H-imidazo[1,2-*a*]pyridine (Burkholder *et al.*, 2001) in DMF (10 ml) was added phosphoryl trichloride (2.2 mmol) in one portion at room temperature under stirring. The mixture was heated to 353 K and stirred for 5.0 h. After the intermediate was consumed (monitored by TLC), the reaction mixture was extracted, filtered and concentrated in vacuo. The pure product was obtained through silica gel chromatography, and yellow blocks of (I) were obtained by slow evaporation of an ethyl acetate/petroleum ether (1:1) solution at room temperature.

Refinement

All H atoms were generated geometrically and refined as riding atoms with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. Molecular structure of (I) with displacement ellipsoids drawn at the 30% probability level for non-H atoms.

2-(4-Chlorophenyl)imidazo[1,2-*a*]pyridine-3-carbaldehyde

Crystal data

$\text{C}_{14}\text{H}_9\text{ClN}_2\text{O}$

$M_r = 256.68$

Orthorhombic, $Pna2_1$

Hall symbol: P 2c -2n

$a = 21.3367$ (13) Å

$b = 7.2391$ (4) Å

$c = 15.1748$ (10) Å

$V = 2343.9$ (2) Å³

$Z = 8$

$F(000) = 1056$

$D_x = 1.455$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4265 reflections

$\theta = 1.9$ – 26.0°

$\mu = 0.31$ mm⁻¹

$T = 293$ K

Block, yellow

$0.35 \times 0.29 \times 0.28$ mm

Data collection

Bruker APEX CCD
diffractometer

Radiation source: fine-focus sealed tube

4198 independent reflections

3564 reflections with $I > 2\sigma(I)$

supplementary materials

graphite	$R_{\text{int}} = 0.026$
ω scans	$\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2002)	$h = -26 \rightarrow 26$
$T_{\text{min}} = 0.870$, $T_{\text{max}} = 0.894$	$k = -8 \rightarrow 8$
12428 measured reflections	$l = -15 \rightarrow 18$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.036$	H-atom parameters constrained
$wR(F^2) = 0.089$	$w = 1/[\sigma^2(F_o^2) + (0.0461P)^2 + 0.2155P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
4198 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
325 parameters	$\Delta\rho_{\text{max}} = 0.22 \text{ e } \text{\AA}^{-3}$
1 restraint	$\Delta\rho_{\text{min}} = -0.18 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), 1800 Friedel pairs
	Flack parameter: 0.40 (5)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.38891 (3)	0.69710 (12)	0.18830 (6)	0.0741 (3)
C12	0.86516 (3)	1.12252 (11)	0.50704 (6)	0.0673 (2)
O2	0.48266 (8)	1.0458 (3)	0.48167 (13)	0.0652 (6)
O1	0.76678 (8)	0.5045 (3)	0.22142 (13)	0.0620 (5)
N1	0.73924 (8)	0.6319 (3)	0.39935 (13)	0.0373 (4)
N2	0.64367 (8)	0.7006 (3)	0.45277 (14)	0.0434 (5)
N4	0.60866 (9)	1.2130 (3)	0.24891 (14)	0.0403 (5)
C6	0.57814 (10)	0.6745 (3)	0.32064 (17)	0.0387 (6)
N3	0.51184 (8)	1.1689 (3)	0.30384 (13)	0.0368 (4)
C7	0.63898 (10)	0.6637 (3)	0.36576 (16)	0.0345 (5)
C21	0.61298 (10)	1.1699 (3)	0.33495 (17)	0.0370 (5)

C11	0.83204 (12)	0.6244 (4)	0.48197 (19)	0.0507 (7)
H11	0.8750	0.6038	0.4857	0.061*
C17	0.79144 (11)	1.1336 (4)	0.45777 (19)	0.0467 (6)
C14	0.70483 (10)	0.6813 (3)	0.47280 (16)	0.0400 (6)
C20	0.67494 (10)	1.1586 (3)	0.37871 (17)	0.0376 (6)
C8	0.69729 (10)	0.6195 (3)	0.32942 (16)	0.0360 (5)
C28	0.54724 (10)	1.2129 (3)	0.23030 (16)	0.0384 (6)
C18	0.78521 (11)	1.0860 (4)	0.37049 (19)	0.0489 (6)
H18	0.8198	1.0467	0.3382	0.059*
C22	0.55419 (10)	1.1401 (3)	0.37297 (16)	0.0380 (5)
C19	0.72678 (10)	1.0975 (3)	0.33157 (17)	0.0423 (6)
H19	0.7221	1.0638	0.2728	0.051*
C2	0.51455 (11)	0.7437 (4)	0.1929 (2)	0.0505 (6)
H2	0.5110	0.7846	0.1350	0.061*
C9	0.71397 (11)	0.5482 (3)	0.24447 (17)	0.0434 (6)
H9	0.6819	0.5340	0.2035	0.052*
C5	0.52395 (11)	0.6241 (3)	0.36600 (18)	0.0448 (6)
H5	0.5268	0.5846	0.4242	0.054*
C24	0.44781 (11)	1.1572 (3)	0.30118 (19)	0.0462 (6)
H24	0.4251	1.1274	0.3515	0.055*
C27	0.51591 (12)	1.2473 (4)	0.15048 (19)	0.0492 (6)
H27	0.5384	1.2786	0.1001	0.059*
C13	0.73559 (11)	0.7049 (4)	0.55341 (19)	0.0507 (7)
H13	0.7133	0.7401	0.6033	0.061*
C10	0.80304 (10)	0.6045 (3)	0.40399 (19)	0.0446 (6)
H10	0.8257	0.5727	0.3539	0.054*
C26	0.45240 (12)	1.2345 (4)	0.14768 (19)	0.0526 (7)
H26	0.4314	1.2554	0.0949	0.063*
C1	0.57237 (11)	0.7353 (4)	0.23404 (18)	0.0439 (6)
H1	0.6080	0.7709	0.2032	0.053*
C16	0.74146 (11)	1.1959 (4)	0.5060 (2)	0.0497 (6)
H16	0.7466	1.2286	0.5649	0.060*
C4	0.46599 (12)	0.6322 (4)	0.32548 (19)	0.0497 (6)
H4	0.4300	0.5988	0.3561	0.060*
C23	0.53638 (11)	1.0703 (3)	0.45701 (17)	0.0463 (6)
H23	0.5683	1.0410	0.4963	0.056*
C15	0.68309 (11)	1.2097 (4)	0.46614 (18)	0.0468 (6)
H15	0.6491	1.2536	0.4983	0.056*
C12	0.79878 (12)	0.6758 (4)	0.5579 (2)	0.0513 (7)
H12	0.8197	0.6899	0.6112	0.062*
C25	0.41833 (12)	1.1899 (4)	0.22393 (19)	0.0514 (7)
H25	0.3749	1.1827	0.2212	0.062*
C3	0.46229 (12)	0.6899 (4)	0.23968 (19)	0.0485 (7)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0499 (4)	0.0888 (6)	0.0835 (6)	0.0081 (4)	-0.0283 (4)	-0.0004 (5)

supplementary materials

C12	0.0459 (3)	0.0834 (5)	0.0725 (5)	-0.0041 (3)	-0.0179 (3)	0.0060 (4)
O2	0.0454 (10)	0.0979 (15)	0.0524 (12)	-0.0022 (10)	0.0123 (9)	0.0151 (11)
O1	0.0459 (10)	0.0937 (16)	0.0462 (12)	0.0041 (10)	0.0086 (9)	-0.0156 (10)
N1	0.0354 (10)	0.0400 (11)	0.0365 (12)	-0.0013 (8)	-0.0023 (9)	-0.0030 (8)
N2	0.0392 (10)	0.0512 (13)	0.0399 (14)	0.0004 (9)	0.0005 (10)	-0.0062 (10)
N4	0.0380 (10)	0.0492 (12)	0.0337 (12)	-0.0002 (9)	0.0021 (9)	-0.0003 (9)
C6	0.0399 (13)	0.0367 (12)	0.0393 (15)	0.0020 (10)	-0.0026 (11)	-0.0046 (11)
N3	0.0343 (10)	0.0400 (10)	0.0361 (12)	0.0006 (8)	-0.0024 (9)	-0.0024 (8)
C7	0.0354 (11)	0.0351 (12)	0.0331 (13)	0.0008 (9)	-0.0014 (10)	-0.0026 (10)
C21	0.0393 (12)	0.0365 (12)	0.0354 (14)	0.0017 (10)	0.0030 (11)	-0.0013 (10)
C11	0.0389 (12)	0.0547 (16)	0.0585 (19)	0.0020 (11)	-0.0094 (13)	-0.0043 (13)
C17	0.0409 (13)	0.0476 (15)	0.0517 (18)	-0.0066 (11)	-0.0098 (13)	0.0087 (12)
C14	0.0388 (12)	0.0448 (14)	0.0364 (15)	0.0006 (11)	0.0010 (11)	-0.0068 (10)
C20	0.0357 (12)	0.0385 (13)	0.0384 (15)	-0.0040 (9)	-0.0001 (10)	0.0044 (11)
C8	0.0366 (12)	0.0379 (12)	0.0337 (13)	-0.0032 (9)	-0.0010 (10)	0.0000 (10)
C28	0.0377 (12)	0.0444 (14)	0.0333 (14)	-0.0012 (10)	0.0025 (10)	-0.0032 (10)
C18	0.0387 (13)	0.0547 (17)	0.0533 (18)	0.0038 (11)	0.0035 (12)	0.0056 (13)
C22	0.0352 (12)	0.0413 (13)	0.0375 (14)	0.0011 (9)	0.0005 (11)	-0.0015 (10)
C19	0.0411 (13)	0.0500 (14)	0.0359 (14)	0.0018 (10)	0.0011 (11)	0.0014 (11)
C2	0.0561 (15)	0.0541 (17)	0.0411 (16)	0.0085 (12)	-0.0045 (14)	0.0009 (13)
C9	0.0438 (13)	0.0479 (15)	0.0385 (14)	-0.0032 (11)	0.0001 (11)	-0.0027 (11)
C5	0.0418 (13)	0.0505 (15)	0.0421 (15)	0.0007 (10)	0.0016 (11)	-0.0004 (12)
C24	0.0359 (12)	0.0513 (15)	0.0513 (17)	-0.0028 (11)	0.0038 (12)	-0.0016 (12)
C27	0.0536 (15)	0.0571 (16)	0.0368 (16)	-0.0002 (12)	-0.0031 (13)	-0.0029 (12)
C13	0.0513 (15)	0.0651 (17)	0.0357 (16)	0.0014 (13)	-0.0052 (13)	-0.0103 (13)
C10	0.0341 (12)	0.0499 (15)	0.0497 (16)	0.0014 (11)	0.0010 (11)	-0.0045 (12)
C26	0.0545 (15)	0.0571 (16)	0.0463 (18)	0.0074 (13)	-0.0121 (14)	-0.0076 (13)
C1	0.0408 (13)	0.0475 (15)	0.0435 (17)	0.0017 (11)	0.0008 (11)	0.0011 (11)
C16	0.0509 (14)	0.0592 (16)	0.0389 (16)	-0.0058 (12)	-0.0078 (13)	-0.0015 (13)
C4	0.0378 (13)	0.0559 (16)	0.0555 (18)	-0.0022 (11)	-0.0025 (12)	-0.0025 (13)
C23	0.0425 (13)	0.0539 (15)	0.0424 (15)	-0.0009 (11)	0.0028 (12)	0.0038 (12)
C15	0.0449 (13)	0.0561 (17)	0.0395 (17)	-0.0011 (11)	0.0036 (12)	0.0015 (12)
C12	0.0503 (15)	0.0621 (17)	0.0414 (16)	-0.0041 (13)	-0.0133 (13)	-0.0032 (13)
C25	0.0402 (13)	0.0579 (17)	0.0560 (18)	0.0024 (12)	-0.0091 (13)	-0.0064 (13)
C3	0.0421 (13)	0.0475 (15)	0.0560 (18)	0.0069 (11)	-0.0129 (13)	-0.0065 (13)

Geometric parameters (Å, °)

C11—C3	1.750 (3)	C28—C27	1.406 (4)
C12—C17	1.743 (2)	C18—C19	1.382 (3)
O2—C23	1.219 (3)	C18—H18	0.9300
O1—C9	1.221 (3)	C22—C23	1.424 (3)
N1—C10	1.378 (3)	C19—H19	0.9300
N1—C14	1.382 (3)	C2—C3	1.378 (4)
N1—C8	1.391 (3)	C2—C1	1.384 (3)
N2—C14	1.347 (3)	C2—H2	0.9300
N2—C7	1.351 (3)	C9—H9	0.9300
N4—C28	1.341 (3)	C5—C4	1.382 (3)
N4—C21	1.346 (3)	C5—H5	0.9300

C6—C1	1.391 (4)	C24—C25	1.351 (4)
C6—C5	1.394 (3)	C24—H24	0.9300
C6—C7	1.470 (3)	C27—C26	1.359 (3)
N3—C24	1.369 (3)	C27—H27	0.9300
N3—C28	1.385 (3)	C13—C12	1.366 (3)
N3—C22	1.400 (3)	C13—H13	0.9300
C7—C8	1.398 (3)	C10—H10	0.9300
C21—C22	1.397 (3)	C26—C25	1.404 (4)
C21—C20	1.482 (3)	C26—H26	0.9300
C11—C10	1.343 (4)	C1—H1	0.9300
C11—C12	1.403 (4)	C16—C15	1.389 (3)
C11—H11	0.9300	C16—H16	0.9300
C17—C16	1.370 (4)	C4—C3	1.370 (4)
C17—C18	1.375 (4)	C4—H4	0.9300
C14—C13	1.399 (4)	C23—H23	0.9300
C20—C15	1.388 (4)	C15—H15	0.9300
C20—C19	1.390 (3)	C12—H12	0.9300
C8—C9	1.434 (3)	C25—H25	0.9300
C10—N1—C14	121.4 (2)	C1—C2—H2	120.8
C10—N1—C8	131.7 (2)	O1—C9—C8	125.4 (2)
C14—N1—C8	106.87 (18)	O1—C9—H9	117.3
C14—N2—C7	105.8 (2)	C8—C9—H9	117.3
C28—N4—C21	105.7 (2)	C4—C5—C6	120.8 (3)
C1—C6—C5	118.4 (2)	C4—C5—H5	119.6
C1—C6—C7	122.3 (2)	C6—C5—H5	119.6
C5—C6—C7	119.2 (2)	C25—C24—N3	118.6 (2)
C24—N3—C28	122.3 (2)	C25—C24—H24	120.7
C24—N3—C22	131.1 (2)	N3—C24—H24	120.7
C28—N3—C22	106.61 (18)	C26—C27—C28	119.3 (3)
N2—C7—C8	111.4 (2)	C26—C27—H27	120.4
N2—C7—C6	120.7 (2)	C28—C27—H27	120.4
C8—C7—C6	127.9 (2)	C12—C13—C14	119.2 (2)
N4—C21—C22	112.0 (2)	C12—C13—H13	120.4
N4—C21—C20	120.6 (2)	C14—C13—H13	120.4
C22—C21—C20	127.4 (2)	C11—C10—N1	119.0 (2)
C10—C11—C12	121.3 (2)	C11—C10—H10	120.5
C10—C11—H11	119.4	N1—C10—H10	120.5
C12—C11—H11	119.4	C27—C26—C25	120.4 (3)
C16—C17—C18	121.5 (2)	C27—C26—H26	119.8
C16—C17—C12	119.2 (2)	C25—C26—H26	119.8
C18—C17—C12	119.3 (2)	C2—C1—C6	121.2 (2)
N2—C14—N1	111.1 (2)	C2—C1—H1	119.4
N2—C14—C13	129.7 (2)	C6—C1—H1	119.4
N1—C14—C13	119.2 (2)	C17—C16—C15	119.2 (3)
C15—C20—C19	118.5 (2)	C17—C16—H16	120.4
C15—C20—C21	121.7 (2)	C15—C16—H16	120.4
C19—C20—C21	119.8 (2)	C3—C4—C5	119.2 (3)
N1—C8—C7	104.9 (2)	C3—C4—H4	120.4
N1—C8—C9	123.3 (2)	C5—C4—H4	120.4

supplementary materials

C7—C8—C9	131.1 (2)	O2—C23—C22	125.3 (2)
N4—C28—N3	111.3 (2)	O2—C23—H23	117.3
N4—C28—C27	130.3 (2)	C22—C23—H23	117.3
N3—C28—C27	118.4 (2)	C20—C15—C16	120.7 (2)
C17—C18—C19	118.9 (2)	C20—C15—H15	119.7
C17—C18—H18	120.5	C16—C15—H15	119.7
C19—C18—H18	120.5	C13—C12—C11	120.0 (3)
C21—C22—N3	104.3 (2)	C13—C12—H12	120.0
C21—C22—C23	131.6 (2)	C11—C12—H12	120.0
N3—C22—C23	123.5 (2)	C24—C25—C26	121.0 (2)
C18—C19—C20	121.1 (2)	C24—C25—H25	119.5
C18—C19—H19	119.4	C26—C25—H25	119.5
C20—C19—H19	119.4	C4—C3—C2	122.0 (2)
C3—C2—C1	118.5 (3)	C4—C3—C11	119.0 (2)
C3—C2—H2	120.8	C2—C3—C11	119.1 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C11—H11 \cdots O2 ⁱ	0.93	2.54	3.442 (3)	163
C12—H12 \cdots N4 ⁱⁱ	0.93	2.59	3.518 (4)	172

Symmetry codes: (i) $x+1/2, -y+3/2, z$; (ii) $-x+3/2, y-1/2, z+1/2$.

Fig. 1

