

6-Allyl-3-(6-chloro-3-pyridylmethyl)-6,7-dihydro-3*H*-1,2,3-triazolo[4,5-*d*]-pyrimidin-7-imine

Dong-Feng Pan,^a Jing Xu,^b Jun-Kai Ma,^{b*} Hong Luo^b and Zuan Ma^b

^aDepartment of Oncology, Renmin Hospital, Yunyang Medical College, Shiyan, 442000, People's Republic of China, and ^bInstitute of Medicinal Chemistry, Yunyang Medical College, Shiyan, 442000, People's Republic of China
Correspondence e-mail: youtianya@126.com


Received 12 November 2009; accepted 13 November 2009

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; disorder in main residue; R factor = 0.040; wR factor = 0.109; data-to-parameter ratio = 13.2.

The title compound, $\text{C}_{13}\text{H}_{12}\text{ClN}_7$, crystallizes with two independent molecules in the asymmetric unit, each with similar geometries. The dihedral angles between the triazole and pyrimidine rings are 0.45 (9) and 1.00 (10) $^\circ$ in the two molecules. A number of $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds co-operate with $\text{C}-\text{H}\cdots\pi$ contacts, forming a supramolecular array in the *ab* plane. $\text{C}-\text{H}\cdots\pi$ interactions are also present. One of the vinyl groups was found to be disordered so that the $\text{C}(\text{H})=\text{CH}_2$ atoms were resolved over two positions with the major component having a site occupancy factor of 0.539 (4).

Related literature

For general background to 8-azapurine derivatives, see: Albert (1986). For the biological activity of 8-azapurines, see: Shiokawa *et al.* (1986); Slusarkchik & Zahler (1989); Subramanian & Gerwick (1989); Vince & Hua (1990); Yamamoto *et al.* (1994).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{12}\text{ClN}_7$
 $M_r = 301.75$
Triclinic, $P\bar{1}$

$\alpha = 87.351 (1)^\circ$
 $\beta = 81.752 (1)^\circ$
 $\gamma = 82.917 (1)^\circ$
 $V = 1395.4 (2)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.28\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.48 \times 0.46 \times 0.43\text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector diffractometer
Absorption correction: none
10116 measured reflections

5052 independent reflections
4277 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.109$
 $S = 1.03$
5052 reflections
384 parameters

22 restraints
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.35\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N10—H10A…N1	0.86	2.44	3.292 (3)	173
N3—H3A…N8 ⁱ	0.86	2.45	3.299 (2)	169
C11—H11A…N3	0.97	2.41	2.749 (2)	100
C19—H19…N11	0.93	2.59	2.909 (3)	101
C3—H3…N13	0.93	2.46	3.309 (2)	151
C11—H11A…Cg4 ⁱⁱ	0.97	2.87	3.446 (2)	119
C24—H24A…Cg1 ⁱⁱⁱ	0.97	2.99	3.851 (3)	149

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, -y + 2, -z + 1$; (iii) $-x + 2, -y + 1, -z + 1$. Cg4 and Cg1 are the centroids of the N4—N6/C1/C4 and N11—N13/C14/C17C4 rings, respectively.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors gratefully acknowledge financial support of this work by Yunyang Medical College (grant No. 2007ZQB24).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2573).

References

- Albert, A. (1986). *Adv. Heterocycl. Chem.* **39**, 117–180.
- Bruker (2000). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shiokawa, K., Tsubo, S. & Moriya, K. (1986). EP Patent 192 060.
- Slusarkchik, W. A. & Zahler, R. (1989). EP Patent 335–355.
- Subramanian, M. V. & Gerwick, B. C. (1989). *Inhibition of Acetylacetate Synthase by Triazolopyrimidines, in Biocatalysis in Agricultural Biotechnology*, edited by J. R. Whitaker & P. E. Sonnet. ACS Symposium Series No. 389, pp. 277–288. Washington, DC, USA: American Chemical Society.
- Vince, R. & Hua, M. (1990). US Patent 4 931 559.
- Yamamoto, I., Yabita, G., Tomizawa, M. & Hissasomi, A. (1994). *J. Pestic. Sci.* **19**, 335–339.

supplementary materials

Acta Cryst. (2009). E65, o3195 [doi:10.1107/S1600536809048168]

6-Allyl-3-(6-chloro-3-pyridylmethyl)-6,7-dihydro-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7-imine

D.-F. Pan, J. Xu, J.-K. Ma, H. Luo and Z. Ma

Comment

1,2,3-Triazolo[4,5-*d*]pyrimidines, *i.e.* 8-azapurines (Albert, 1986), have attracted attention because some of these derivatives exhibit anti-viral (Slusarkchik & Zahler, 1989), anti-tumour (Slusarkchik & Zahler, 1989; Vince & Hua, 1990), and herbicidal activities (Subramanian & Gerwick, 1989). Neonicotinoid insecticides, as nicotinic acetylcholine receptor inhibitors, have also attracted increasing attention because of their low toxicity, wide range of activities, and high potency (Shiokawa *et al.*, 1986). It has been found that most biologically active nicotinic compounds contain the 3-aminomethylpyridine group (Yamamoto *et al.*, 1994). Herein, we report the crystal structure of (I), Fig. 1, which was synthesized by introducing pyridine rings into a 1,2,3-triazolo[4,5-*d*]pyrimidine framework.

Several N—H···N hydrogen bonding contacts, together with C—H···N and C—H···π interactions, lead to the formation of supramolecular arrays in the *ab* plane, Table 1 and Fig. 2. In addition π—π stacking interactions stabilize the crystal structure, with the shortest centroid-centroid distance of 3.412 (1) Å occurring between centrosymmetrically related planes through the (N4—N6, C1, C4) rings, symmetry operation: 2-x, 1-y, 1-z.

Experimental

Allylamine (1 mmol) in anhydrous acetonitrile (4 ml) was added dropwise to a solution of ethyl-*N*-3-((6-chloropyridin-3-yl)methyl)-5-cyano-3*H*-1,2,3-triazol-4-yl-formimidate (1 mmol) in anhydrous acetonitrile (8 ml) at room temperature. The mixture was stirred until the reaction was complete (by thin layer chromatography) and the solution concentrated under vacuum. The residue was recrystallized from anhydrous ethanol to give (I) (yield 87%). Colourless crystals were grown from a dichloromethane and petroleum ether (1:3 *v/v*) solution of (I).

Refinement

H atoms were placed in calculated positions, with C—H distances in the range 0.93–0.97 Å and N—H distances of 0.86 Å, and included in the final cycles of refinement using a riding-model approximation, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}$ (carrier atom). A rotating group model was used for the methyl groups. Disorder was noted in the C24-C26 vinyl substituent in that two positions were resolved for the C25 atom. From refinement, the major component had a site occupancy factor = 0.539 (4).

supplementary materials

Figures




Fig. 1. The molecular structures of the two independent molecules in (I), showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. A view of a supramolecular layer in (I). Hydrogen bonds are shown as dashed lines.

6-Allyl-3-(6-chloro-3-pyridylmethyl)-6,7-dihydro-3*H*-1,2,3-triazolo[4,5-*d*]pyrimidin-7-imine

Crystal data

C ₁₃ H ₁₂ ClN ₇	Z = 4
M _r = 301.75	F(000) = 624
Triclinic, P $\bar{1}$	D _x = 1.436 Mg m ⁻³
Hall symbol: -P 1	Mo K α radiation, λ = 0.71073 Å
a = 7.2845 (7) Å	Cell parameters from 5848 reflections
b = 13.2684 (12) Å	θ = 2.8–28.2°
c = 14.7069 (14) Å	μ = 0.28 mm ⁻¹
α = 87.351 (1)°	T = 296 K
β = 81.752 (1)°	Block, colorless
γ = 82.917 (1)°	0.48 × 0.46 × 0.43 mm
V = 1395.4 (2) Å ³	

Data collection

Bruker SMART APEX CCD area-detector diffractometer	4277 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.015$
graphite	$\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.8^\circ$
φ and ω scans	$h = -8 \rightarrow 8$
10116 measured reflections	$k = -15 \rightarrow 16$
5052 independent reflections	$l = -17 \rightarrow 17$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained

$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0474P)^2 + 0.5781P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
5052 reflections	$(\Delta/\sigma)_{\max} = 0.001$
384 parameters	$\Delta\rho_{\max} = 0.35 \text{ e } \text{\AA}^{-3}$
22 restraints	$\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: <i>SHELXL</i> , $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
	Extinction coefficient: 0.050 (2)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken

into account individually in the estimation of e.s.d.'s in distances, angles

and torsion angles; correlations between e.s.d.'s in cell parameters are only

used when they are defined by crystal symmetry. An approximate (isotropic)

treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and

goodness of fit S are based on F^2 , conventional R -factors R are based

on F , with F set to zero for negative F^2 . The threshold expression of

$F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is

not relevant to the choice of reflections for refinement. R -factors based

on F^2 are statistically about twice as large as those based on F , and R -

factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C24	0.7964 (4)	0.39425 (19)	0.67973 (16)	0.0710 (7)	0.539 (4)
H24A	0.9002	0.4288	0.6913	0.085*	0.539 (4)
H24B	0.6844	0.4265	0.7159	0.085*	0.539 (4)
C25	0.8281 (6)	0.2896 (3)	0.7111 (3)	0.0666 (9)	0.539 (4)
H25	0.9341	0.2520	0.6814	0.080*	0.539 (4)
C26	0.7319 (5)	0.2418 (3)	0.7732 (2)	0.1002 (10)	0.539 (4)
H26A	0.6239	0.2747	0.8058	0.120*	0.539 (4)
H26B	0.7692	0.1738	0.7864	0.120*	0.539 (4)
C24'	0.7964 (4)	0.39425 (19)	0.67973 (16)	0.0710 (7)	0.461 (4)
H24C	0.9279	0.3756	0.6840	0.085*	0.461 (4)

supplementary materials

H24D	0.7600	0.4593	0.7086	0.085*	0.461 (4)
C25'	0.6943 (8)	0.3207 (4)	0.7325 (3)	0.0666 (9)	0.461 (4)
H25'	0.5659	0.3385	0.7363	0.080*	0.461 (4)
C26'	0.7319 (5)	0.2418 (3)	0.7732 (2)	0.1002 (10)	0.461 (4)
H26C	0.8561	0.2159	0.7746	0.120*	0.461 (4)
H26D	0.6368	0.2068	0.8034	0.120*	0.461 (4)
C1	0.7678 (2)	1.00162 (12)	0.51743 (12)	0.0346 (4)	
C2	0.7015 (2)	1.02006 (12)	0.42960 (12)	0.0360 (4)	
C3	0.7187 (3)	0.83395 (13)	0.43489 (14)	0.0448 (4)	
H3	0.6996	0.7774	0.4040	0.054*	
C4	0.8029 (2)	0.90450 (12)	0.55236 (12)	0.0362 (4)	
C5	0.9108 (3)	0.83866 (14)	0.70333 (13)	0.0457 (4)	
H5A	1.0253	0.8510	0.7252	0.055*	
H5B	0.9309	0.7719	0.6768	0.055*	
C6	0.5880 (3)	0.80588 (16)	0.77284 (13)	0.0493 (5)	
H6	0.5802	0.7758	0.7179	0.059*	
C7	0.7544 (3)	0.84151 (13)	0.78275 (12)	0.0406 (4)	
C8	0.7665 (3)	0.88164 (17)	0.86595 (15)	0.0588 (5)	
H8	0.8770	0.9047	0.8763	0.071*	
C9	0.6143 (3)	0.88762 (18)	0.93401 (15)	0.0639 (6)	
H9	0.6193	0.9140	0.9910	0.077*	
C10	0.4556 (3)	0.85311 (15)	0.91408 (13)	0.0511 (5)	
C11	0.6112 (3)	0.92790 (16)	0.30226 (13)	0.0484 (5)	
H11A	0.5276	0.9899	0.2960	0.058*	
H11B	0.5401	0.8711	0.3001	0.058*	
C12	0.7634 (3)	0.9228 (2)	0.22394 (16)	0.0681 (6)	
H12	0.8453	0.9719	0.2193	0.082*	
C13	0.7911 (4)	0.8556 (3)	0.1618 (2)	0.1080 (12)	
H13A	0.7121	0.8053	0.1641	0.130*	
H13B	0.8902	0.8573	0.1144	0.130*	
C14	0.7367 (2)	0.51225 (13)	0.45004 (13)	0.0407 (4)	
C15	0.7607 (3)	0.51058 (14)	0.54603 (13)	0.0434 (4)	
C16	0.7703 (3)	0.32703 (15)	0.52893 (15)	0.0523 (5)	
H16	0.7816	0.2638	0.5589	0.063*	
C17	0.7361 (2)	0.42382 (13)	0.40561 (13)	0.0398 (4)	
C18	0.7094 (3)	0.39121 (15)	0.23947 (14)	0.0514 (5)	
H18A	0.5996	0.4168	0.2114	0.062*	
H18B	0.6960	0.3220	0.2611	0.062*	
C19	1.0541 (3)	0.40537 (14)	0.18782 (14)	0.0499 (5)	
H19	1.0650	0.4172	0.2486	0.060*	
C20	0.8801 (3)	0.39059 (13)	0.16747 (13)	0.0440 (4)	
C21	0.8673 (3)	0.37205 (16)	0.07674 (14)	0.0549 (5)	
H21	0.7530	0.3618	0.0598	0.066*	
C22	1.0239 (3)	0.36883 (17)	0.01182 (15)	0.0608 (6)	
H22	1.0184	0.3558	-0.0493	0.073*	
C23	1.1885 (3)	0.38548 (15)	0.04051 (15)	0.0546 (5)	
C11	0.25485 (10)	0.86518 (6)	0.99613 (4)	0.0824 (2)	
C12	1.39020 (10)	0.38267 (5)	-0.04027 (5)	0.0812 (2)	
N1	0.7805 (2)	0.81572 (11)	0.51360 (11)	0.0446 (4)	

N2	0.6786 (2)	0.92572 (11)	0.39244 (10)	0.0399 (3)
N3	0.6629 (2)	1.10214 (11)	0.38456 (11)	0.0450 (4)
H3A	0.6766	1.1591	0.4069	0.054*
N4	0.86503 (19)	0.91590 (10)	0.63304 (10)	0.0384 (3)
N5	0.8663 (2)	1.01637 (11)	0.64796 (10)	0.0412 (4)
N6	0.8078 (2)	1.06876 (11)	0.57735 (10)	0.0386 (3)
N7	0.4377 (2)	0.81185 (14)	0.83706 (11)	0.0541 (4)
N8	0.7516 (2)	0.32692 (11)	0.44197 (12)	0.0503 (4)
N9	0.7752 (2)	0.40853 (12)	0.58091 (11)	0.0487 (4)
N10	0.7707 (3)	0.58234 (13)	0.59890 (12)	0.0575 (5)
H10A	0.7618	0.6440	0.5777	0.069*
N11	0.7175 (2)	0.45313 (11)	0.31812 (11)	0.0451 (4)
N12	0.7058 (3)	0.55636 (12)	0.30882 (12)	0.0545 (4)
N13	0.7175 (3)	0.59204 (12)	0.38966 (12)	0.0515 (4)
N14	1.2090 (2)	0.40379 (13)	0.12549 (12)	0.0551 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C24	0.0985 (19)	0.0654 (15)	0.0573 (14)	-0.0258 (13)	-0.0265 (13)	0.0029 (11)
C25	0.065 (2)	0.079 (2)	0.0500 (19)	0.0091 (18)	-0.0045 (17)	0.0038 (16)
C26	0.150 (3)	0.093 (2)	0.0610 (17)	-0.037 (2)	-0.0126 (18)	0.0071 (16)
C24'	0.0985 (19)	0.0654 (15)	0.0573 (14)	-0.0258 (13)	-0.0265 (13)	0.0029 (11)
C25'	0.065 (2)	0.079 (2)	0.0500 (19)	0.0091 (18)	-0.0045 (17)	0.0038 (16)
C26'	0.150 (3)	0.093 (2)	0.0610 (17)	-0.037 (2)	-0.0126 (18)	0.0071 (16)
C1	0.0295 (8)	0.0307 (8)	0.0413 (9)	-0.0047 (6)	0.0045 (7)	-0.0034 (7)
C2	0.0276 (8)	0.0352 (9)	0.0432 (10)	-0.0045 (6)	0.0037 (7)	-0.0052 (7)
C3	0.0447 (10)	0.0334 (9)	0.0554 (12)	-0.0065 (7)	0.0007 (8)	-0.0110 (8)
C4	0.0312 (8)	0.0321 (8)	0.0428 (10)	-0.0045 (6)	0.0042 (7)	-0.0023 (7)
C5	0.0446 (10)	0.0416 (10)	0.0485 (11)	-0.0001 (8)	-0.0046 (8)	0.0060 (8)
C6	0.0538 (11)	0.0573 (12)	0.0388 (10)	-0.0159 (9)	-0.0041 (8)	-0.0049 (9)
C7	0.0466 (10)	0.0334 (9)	0.0414 (10)	-0.0049 (7)	-0.0070 (8)	0.0050 (7)
C8	0.0591 (13)	0.0651 (14)	0.0568 (13)	-0.0206 (10)	-0.0102 (10)	-0.0094 (10)
C9	0.0783 (15)	0.0719 (15)	0.0449 (12)	-0.0233 (12)	-0.0028 (11)	-0.0168 (10)
C10	0.0619 (12)	0.0475 (11)	0.0421 (11)	-0.0126 (9)	0.0040 (9)	0.0009 (9)
C11	0.0442 (10)	0.0509 (11)	0.0512 (11)	-0.0049 (8)	-0.0076 (8)	-0.0114 (9)
C12	0.0555 (13)	0.1005 (19)	0.0509 (13)	-0.0218 (12)	-0.0025 (10)	-0.0115 (12)
C13	0.0756 (18)	0.169 (3)	0.0780 (19)	-0.0095 (19)	0.0080 (15)	-0.055 (2)
C14	0.0416 (9)	0.0328 (9)	0.0478 (10)	-0.0046 (7)	-0.0048 (8)	-0.0076 (8)
C15	0.0442 (10)	0.0377 (9)	0.0494 (11)	-0.0077 (8)	-0.0063 (8)	-0.0065 (8)
C16	0.0639 (13)	0.0353 (10)	0.0589 (13)	-0.0094 (9)	-0.0093 (10)	-0.0001 (9)
C17	0.0397 (9)	0.0332 (9)	0.0464 (11)	-0.0049 (7)	-0.0040 (8)	-0.0070 (7)
C18	0.0596 (12)	0.0466 (11)	0.0516 (12)	-0.0104 (9)	-0.0122 (9)	-0.0133 (9)
C19	0.0639 (13)	0.0438 (11)	0.0448 (11)	-0.0080 (9)	-0.0150 (9)	-0.0042 (8)
C20	0.0581 (11)	0.0306 (9)	0.0455 (11)	-0.0040 (8)	-0.0139 (9)	-0.0054 (7)
C21	0.0660 (13)	0.0504 (12)	0.0518 (12)	-0.0074 (10)	-0.0178 (10)	-0.0096 (9)
C22	0.0794 (16)	0.0599 (13)	0.0441 (12)	-0.0061 (11)	-0.0110 (11)	-0.0102 (10)
C23	0.0660 (13)	0.0419 (11)	0.0532 (13)	-0.0013 (9)	-0.0040 (10)	0.0004 (9)

supplementary materials

Cl1	0.0861 (5)	0.0912 (5)	0.0643 (4)	-0.0290 (4)	0.0286 (3)	-0.0165 (3)
Cl2	0.0782 (4)	0.0853 (5)	0.0717 (4)	-0.0018 (3)	0.0105 (3)	0.0013 (3)
N1	0.0481 (9)	0.0305 (8)	0.0540 (10)	-0.0051 (6)	-0.0016 (7)	-0.0038 (7)
N2	0.0381 (8)	0.0369 (8)	0.0441 (8)	-0.0049 (6)	-0.0014 (6)	-0.0074 (6)
N3	0.0471 (9)	0.0364 (8)	0.0506 (9)	-0.0031 (7)	-0.0054 (7)	-0.0007 (7)
N4	0.0371 (8)	0.0338 (7)	0.0424 (8)	-0.0049 (6)	0.0010 (6)	0.0006 (6)
N5	0.0408 (8)	0.0376 (8)	0.0442 (9)	-0.0071 (6)	0.0002 (6)	-0.0029 (7)
N6	0.0379 (8)	0.0336 (7)	0.0431 (8)	-0.0062 (6)	0.0008 (6)	-0.0028 (6)
N7	0.0555 (10)	0.0642 (11)	0.0438 (9)	-0.0203 (8)	0.0004 (8)	-0.0022 (8)
N8	0.0652 (11)	0.0330 (8)	0.0537 (10)	-0.0085 (7)	-0.0071 (8)	-0.0074 (7)
N9	0.0599 (10)	0.0410 (9)	0.0482 (9)	-0.0128 (7)	-0.0117 (8)	-0.0012 (7)
N10	0.0796 (12)	0.0424 (9)	0.0541 (10)	-0.0111 (8)	-0.0146 (9)	-0.0116 (8)
N11	0.0544 (9)	0.0351 (8)	0.0473 (9)	-0.0064 (7)	-0.0082 (7)	-0.0094 (7)
N12	0.0752 (12)	0.0367 (9)	0.0527 (10)	-0.0044 (8)	-0.0139 (9)	-0.0055 (7)
N13	0.0693 (11)	0.0337 (8)	0.0525 (10)	-0.0035 (7)	-0.0124 (8)	-0.0076 (7)
N14	0.0595 (11)	0.0495 (10)	0.0570 (11)	-0.0068 (8)	-0.0114 (9)	0.0004 (8)

Geometric parameters (\AA , $^\circ$)

C24—C25	1.445 (5)	C11—H11B	0.9700
C24—N9	1.485 (3)	C12—C13	1.283 (4)
C24—H24A	0.9700	C12—H12	0.9300
C24—H24B	0.9700	C13—H13A	0.9300
C25—C26	1.265 (5)	C13—H13B	0.9300
C25—H25	0.9300	C14—N13	1.357 (2)
C26—H26A	0.9300	C14—C17	1.370 (2)
C26—H26B	0.9300	C14—C15	1.446 (3)
C25'—H25'	0.9300	C15—N10	1.272 (2)
C1—N6	1.366 (2)	C15—N9	1.423 (2)
C1—C4	1.373 (2)	C16—N8	1.305 (3)
C1—C2	1.443 (2)	C16—N9	1.360 (2)
C2—N3	1.271 (2)	C16—H16	0.9300
C2—N2	1.426 (2)	C17—N11	1.348 (2)
C3—N1	1.303 (3)	C17—N8	1.367 (2)
C3—N2	1.361 (2)	C18—N11	1.461 (2)
C3—H3	0.9300	C18—C20	1.512 (3)
C4—N4	1.350 (2)	C18—H18A	0.9700
C4—N1	1.369 (2)	C18—H18B	0.9700
C5—N4	1.465 (2)	C19—N14	1.346 (3)
C5—C7	1.509 (3)	C19—C20	1.383 (3)
C5—H5A	0.9700	C19—H19	0.9300
C5—H5B	0.9700	C20—C21	1.386 (3)
C6—N7	1.337 (3)	C21—C22	1.377 (3)
C6—C7	1.383 (3)	C21—H21	0.9300
C6—H6	0.9300	C22—C23	1.372 (3)
C7—C8	1.375 (3)	C22—H22	0.9300
C8—C9	1.380 (3)	C23—N14	1.317 (3)
C8—H8	0.9300	C23—Cl2	1.750 (2)
C9—C10	1.368 (3)	N3—H3A	0.8600

C9—H9	0.9300	N4—N5	1.362 (2)
C10—N7	1.310 (3)	N5—N6	1.315 (2)
C10—Cl1	1.753 (2)	N10—H10A	0.8600
C11—N2	1.477 (2)	N11—N12	1.364 (2)
C11—C12	1.478 (3)	N12—N13	1.318 (2)
C11—H11A	0.9700		
C25—C24—N9	114.8 (2)	H13A—C13—H13B	120.0
C25—C24—H24A	108.6	N13—C14—C17	109.22 (16)
N9—C24—H24A	108.6	N13—C14—C15	129.91 (16)
C25—C24—H24B	108.6	C17—C14—C15	120.84 (16)
N9—C24—H24B	108.6	N10—C15—N9	119.50 (18)
H24A—C24—H24B	107.6	N10—C15—C14	130.94 (18)
C26—C25—C24	129.2 (4)	N9—C15—C14	109.56 (15)
C26—C25—H25	115.4	N8—C16—N9	127.79 (18)
C24—C25—H25	115.4	N8—C16—H16	116.1
C25—C26—H26A	120.0	N9—C16—H16	116.1
C25—C26—H26B	120.0	N11—C17—N8	127.49 (16)
H26A—C26—H26B	120.0	N11—C17—C14	104.94 (15)
N6—C1—C4	109.19 (15)	N8—C17—C14	127.56 (17)
N6—C1—C2	129.88 (15)	N11—C18—C20	113.36 (16)
C4—C1—C2	120.92 (15)	N11—C18—H18A	108.9
N3—C2—N2	119.22 (16)	C20—C18—H18A	108.9
N3—C2—C1	131.28 (16)	N11—C18—H18B	108.9
N2—C2—C1	109.50 (14)	C20—C18—H18B	108.9
N1—C3—N2	127.88 (17)	H18A—C18—H18B	107.7
N1—C3—H3	116.1	N14—C19—C20	124.30 (18)
N2—C3—H3	116.1	N14—C19—H19	117.8
N4—C4—N1	127.68 (16)	C20—C19—H19	117.8
N4—C4—C1	104.78 (15)	C19—C20—C21	116.96 (19)
N1—C4—C1	127.53 (17)	C19—C20—C18	122.96 (17)
N4—C5—C7	110.31 (14)	C21—C20—C18	120.06 (18)
N4—C5—H5A	109.6	C22—C21—C20	119.9 (2)
C7—C5—H5A	109.6	C22—C21—H21	120.0
N4—C5—H5B	109.6	C20—C21—H21	120.0
C7—C5—H5B	109.6	C23—C22—C21	117.6 (2)
H5A—C5—H5B	108.1	C23—C22—H22	121.2
N7—C6—C7	124.30 (18)	C21—C22—H22	121.2
N7—C6—H6	117.8	N14—C23—C22	125.2 (2)
C7—C6—H6	117.8	N14—C23—Cl2	115.96 (17)
C8—C7—C6	117.12 (18)	C22—C23—Cl2	118.84 (17)
C8—C7—C5	122.66 (18)	C3—N1—C4	110.65 (15)
C6—C7—C5	120.18 (17)	C3—N2—C2	123.51 (15)
C7—C8—C9	119.8 (2)	C3—N2—C11	118.39 (15)
C7—C8—H8	120.1	C2—N2—C11	118.09 (15)
C9—C8—H8	120.1	C2—N3—H3A	119.3
C10—C9—C8	117.16 (19)	C4—N4—N5	110.08 (14)
C10—C9—H9	121.4	C4—N4—C5	129.06 (15)
C8—C9—H9	121.4	N5—N4—C5	120.63 (15)
N7—C10—C9	125.59 (19)	N6—N5—N4	107.99 (14)

supplementary materials

N7—C10—Cl1	115.58 (16)	N5—N6—C1	107.95 (14)
C9—C10—Cl1	118.82 (16)	C10—N7—C6	115.93 (17)
N2—C11—C12	113.26 (16)	C16—N8—C17	110.70 (16)
N2—C11—H11A	108.9	C16—N9—C15	123.52 (17)
C12—C11—H11A	108.9	C16—N9—C24	120.41 (17)
N2—C11—H11B	108.9	C15—N9—C24	116.08 (16)
C12—C11—H11B	108.9	C15—N10—H10A	119.3
H11A—C11—H11B	107.7	C17—N11—N12	109.97 (14)
C13—C12—C11	124.9 (3)	C17—N11—C18	129.37 (16)
C13—C12—H12	117.5	N12—N11—C18	120.66 (16)
C11—C12—H12	117.5	N13—N12—N11	107.63 (15)
C12—C13—H13A	120.0	N12—N13—C14	108.23 (15)
C12—C13—H13B	120.0	C23—N14—C19	116.03 (18)
N9—C24—C25—C26	−120.5 (5)	N3—C2—N2—C11	0.2 (2)
N6—C1—C2—N3	0.1 (3)	C1—C2—N2—C11	179.95 (14)
C4—C1—C2—N3	−179.05 (17)	C12—C11—N2—C3	−89.6 (2)
N6—C1—C2—N2	−179.57 (15)	C12—C11—N2—C2	89.2 (2)
C4—C1—C2—N2	1.3 (2)	N1—C4—N4—N5	−179.62 (15)
N6—C1—C4—N4	−0.42 (17)	C1—C4—N4—N5	0.66 (17)
C2—C1—C4—N4	178.88 (14)	N1—C4—N4—C5	−5.2 (3)
N6—C1—C4—N1	179.86 (15)	C1—C4—N4—C5	175.05 (15)
C2—C1—C4—N1	−0.8 (3)	C7—C5—N4—C4	−101.8 (2)
N7—C6—C7—C8	2.9 (3)	C7—C5—N4—N5	72.1 (2)
N7—C6—C7—C5	−175.14 (18)	C4—N4—N5—N6	−0.68 (18)
N4—C5—C7—C8	−105.4 (2)	C5—N4—N5—N6	−175.62 (14)
N4—C5—C7—C6	72.5 (2)	N4—N5—N6—C1	0.39 (17)
C6—C7—C8—C9	−1.9 (3)	C4—C1—N6—N5	0.02 (18)
C5—C7—C8—C9	176.1 (2)	C2—C1—N6—N5	−179.19 (15)
C7—C8—C9—C10	−0.5 (3)	C9—C10—N7—C6	−1.6 (3)
C8—C9—C10—N7	2.4 (4)	C11—C10—N7—C6	177.57 (15)
C8—C9—C10—Cl1	−176.73 (18)	C7—C6—N7—C10	−1.2 (3)
N2—C11—C12—C13	123.7 (3)	N9—C16—N8—C17	−0.1 (3)
N13—C14—C15—N10	−0.3 (4)	N11—C17—N8—C16	179.08 (19)
C17—C14—C15—N10	177.7 (2)	C14—C17—N8—C16	−0.8 (3)
N13—C14—C15—N9	−179.85 (18)	N8—C16—N9—C15	−0.3 (3)
C17—C14—C15—N9	−1.9 (2)	N8—C16—N9—C24	179.6 (2)
N13—C14—C17—N11	0.4 (2)	N10—C15—N9—C16	−178.4 (2)
C15—C14—C17—N11	−178.00 (16)	C14—C15—N9—C16	1.2 (3)
N13—C14—C17—N8	−179.76 (18)	N10—C15—N9—C24	1.6 (3)
C15—C14—C17—N8	1.9 (3)	C14—C15—N9—C24	−178.73 (18)
N14—C19—C20—C21	0.7 (3)	C25—C24—N9—C16	5.5 (4)
N14—C19—C20—C18	179.01 (18)	C25—C24—N9—C15	−174.5 (3)
N11—C18—C20—C19	27.4 (3)	N8—C17—N11—N12	179.79 (18)
N11—C18—C20—C21	−154.32 (18)	C14—C17—N11—N12	−0.3 (2)
C19—C20—C21—C22	0.1 (3)	N8—C17—N11—C18	−0.3 (3)
C18—C20—C21—C22	−178.26 (19)	C14—C17—N11—C18	179.60 (18)
C20—C21—C22—C23	−0.6 (3)	C20—C18—N11—C17	−110.6 (2)
C21—C22—C23—N14	0.4 (3)	C20—C18—N11—N12	69.3 (2)
C21—C22—C23—Cl2	−179.65 (16)	C17—N11—N12—N13	0.2 (2)

N2—C3—N1—C4	0.0 (3)	C18—N11—N12—N13	-179.76 (17)
N4—C4—N1—C3	-179.58 (17)	N11—N12—N13—C14	0.1 (2)
C1—C4—N1—C3	0.1 (2)	C17—C14—N13—N12	-0.3 (2)
N1—C3—N2—C2	0.6 (3)	C15—C14—N13—N12	177.90 (19)
N1—C3—N2—C11	179.45 (18)	C22—C23—N14—C19	0.3 (3)
N3—C2—N2—C3	179.06 (16)	C12—C23—N14—C19	-179.61 (14)
C1—C2—N2—C3	-1.2 (2)	C20—C19—N14—C23	-0.9 (3)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N10—H10A···N1	0.86	2.44	3.292 (3)	173
N3—H3A···N8 ⁱ	0.86	2.45	3.299 (2)	169.
C11—H11A···N3	0.97	2.41	2.749 (2)	100
C19—H19···N11	0.93	2.59	2.909 (3)	101
C3—H3···N13	0.93	2.46	3.309 (2)	151
C11—H11A···Cg ₄ ⁱⁱ	0.97	2.87	3.446 (2)	119
C24—H24A···Cg ₁ ⁱⁱⁱ	0.97	2.99	3.851 (3)	149

Symmetry codes: (i) $x, y+1, z$; (ii) $-x+1, -y+2, -z+1$; (iii) $-x+2, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

