

catena-Poly[[di- μ -iodido-dicopper(I)-(Cu—Cu)]bis(μ -4,4'-di-3-pyridyl-2,2'-disulfaneyldipyrimidine)]

Hai-Bin Zhu

School of Chemistry and Chemical Engineering, Southeast University, Nanjing 211189, People's Republic of China, and School of Material Science and Engineering, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: zhuhaibin@seu.edu.cn


Received 28 November 2009; accepted 5 December 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.041; wR factor = 0.108; data-to-parameter ratio = 14.1.

The title complex, $[\text{Cu}_2\text{I}_2(\text{C}_{18}\text{H}_{12}\text{N}_6\text{S}_2)_2]_n$, contains a Cu_2I_2 core with a Cu—Cu distance of 2.6935 (14) Å. The Cu^{I} atom is coordinated by two bridging 4,4'-di-3-pyridyl-2,2'-disulfaneyldipyrimidine ligands and two bridging I atoms, forming a double chain.

Related literature

For coordination polymers with 4,4'-dipyridinedisulfide, see: Horikoshi & Mochida (2006). For coordination polymers with 2,2'-dithiobis(4-pyridin-4-yl-pyrimidine), see: Zhu *et al.* (2009). For the structure of free 2,2'-dithiobis(3-pyridin-4-yl-pyrimidine), see: Ji *et al.* (2009).



Experimental

Crystal data

$[\text{Cu}_2\text{I}_2(\text{C}_{18}\text{H}_{12}\text{N}_6\text{S}_2)_2]$	$\gamma = 96.449$ (1) $^\circ$
$M_r = 1133.86$	$V = 1023.66$ (13) Å 3
Triclinic, $P\bar{1}$	$Z = 1$
$a = 8.5561$ (6) Å	Mo $K\alpha$ radiation
$b = 10.7702$ (8) Å	$\mu = 2.80$ mm $^{-1}$
$c = 11.9045$ (8) Å	$T = 298$ K
$\alpha = 98.110$ (1) $^\circ$	$0.19 \times 0.15 \times 0.12$ mm
$\beta = 107.193$ (1) $^\circ$	

Data collection

Bruker APEXII CCD diffractometer	5395 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2001)	3568 independent reflections
$T_{\min} = 0.611$, $T_{\max} = 0.715$	2956 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.097$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$	253 parameters
$wR(F^2) = 0.108$	H-atom parameters constrained
$S = 0.99$	$\Delta\rho_{\max} = 1.28$ e Å $^{-3}$
3568 reflections	$\Delta\rho_{\min} = -1.03$ e Å $^{-3}$

Table 1

Selected bond lengths (Å).

Cu1—I1	2.6550 (7)	Cu1—N1	2.037 (4)
Cu1—I1 ⁱ	2.6579 (8)	Cu1—N6 ⁱⁱ	2.060 (4)

Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $x, y + 1, z - 1$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT-Plus (Bruker, 2007); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008) and DIAMOND (Brandenburg, 1999); software used to prepare material for publication: SHELXTL.

The author acknowledges financial support from the China Postdoctoral Research Fund (20070411010).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2260).

References

- Brandenburg, K. (1999). DIAMOND. Crystal Impact GbR, Bonn, Germany.
Bruker (2001). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2007). APEX2 and SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.
Horikoshi, R. & Mochida, T. (2006). *Coord. Chem. Rev.* **250**, 2595–2609.
Ji, J.-F., Li, L. & Zhu, H.-B. (2009). *Acta Cryst.* **E65**, o1253.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Zhu, H.-B., Wang, H., Kong, F., Gou, S.-H. & Sun, Y.-M. (2009). *J. Mol. Struct.* **936**, 99–103.

supplementary materials

Acta Cryst. (2010). E66, m41 [doi:10.1107/S1600536809052325]

catena-Poly[[di- μ -iodido-dicopper(I)(Cu-Cu)]bis(μ -4,4'-di-3-pyridyl-2,2'-disulfanediyl)dipyrimidine]

H.-B. Zhu

Comment

In recent years, heterocyclic disulfide ligands have received increasing attention because of their conformationally defined dihedral angle (Horikoshi & Mochida, 2006). As continuation of our previous research (Zhu *et al.*, 2009), we report here a copper(I) coordination polymer with a 2,2'-dithiobis(3-pyridin-4-ylpyrimidine) (*L*) ligand.

The Cu^I atom in the title complex has a tetrahedral coordination geometry completed by two N atoms from two different *L* ligands and two bridging I atoms (Fig. 1 and Table 1). The C—S—S—C torsion angle of 81.2 (2)° in *L* is almost identical with the free molecule (Ji *et al.*, 2009). Alternative linkings of one Cu₂I₂ core and two bridging *L* ligands generate a one-dimensional double chain (Fig. 2).

Experimental

A CH₂Cl₂ solution (5 ml) of ligand *L* (0.1 mmol) was slowly added into a CuI (0.1 mmol) solution in acetonitrile (10 ml). The mixture was kept on standing for 3 d to give single crystals suitable for X-ray diffraction analysis.

Refinement

H atoms were positioned geometrically and refined as riding atoms, with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The highest residual electron density was found 0.98 Å from I1 and the deepest hole 0.92 Å from I1.

Figures



Fig. 1. The structure of the title compound with 30% probability displacement ellipsoids. H atoms have been omitted for clarity. [Symmetry codes: (i) $-x + 1, -y + 2, -z$; (ii) $x, y + 1, z - 1$.]



Fig. 2. The one-dimensional double chain viewed along the *a* axis.

catena-Poly[[di- μ -iodido-dicopper(I)(Cu—Cu)]bis(μ -4,4'-di-3-pyridyl-2,2'-disulfanediyl)dipyrimidine]

Crystal data

[Cu₂I₂(C₁₈H₁₂N₆S₂)₂]

$M_r = 1133.86$

Triclinic, $P\bar{1}$

$Z = 1$

$F(000) = 552$

$D_x = 1.839 \text{ Mg m}^{-3}$

supplementary materials

Hall symbol: -P 1
 $a = 8.5561$ (6) Å
 $b = 10.7702$ (8) Å
 $c = 11.9045$ (8) Å
 $\alpha = 98.110$ (1)°
 $\beta = 107.193$ (1)°
 $\gamma = 96.449$ (1)°
 $V = 1023.66$ (13) Å³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 3568 reflections
 $\theta = 2.3$ – 25.5 °
 $\mu = 2.80$ mm⁻¹
 $T = 298$ K
Block, yellow
 $0.19 \times 0.15 \times 0.12$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
graphite
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2001)
 $T_{\min} = 0.611$, $T_{\max} = 0.715$
5395 measured reflections

3568 independent reflections
2956 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.097$
 $\theta_{\max} = 25.0$ °, $\theta_{\min} = 1.8$ °
 $h = -7 \rightarrow 10$
 $k = -12 \rightarrow 12$
 $l = -14 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.108$
 $S = 0.99$
3568 reflections
253 parameters
0 restraints

Primary atom site location: structure-invariant direct methods
Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0541P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.28$ e Å⁻³
 $\Delta\rho_{\min} = -1.03$ e Å⁻³

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
II	0.31696 (4)	0.97725 (3)	0.10525 (3)	0.04697 (15)
Cu1	0.38873 (9)	0.92391 (6)	-0.09670 (6)	0.0503 (2)
S1	0.13540 (17)	0.55124 (12)	0.27577 (10)	0.0447 (3)
S2	0.02067 (16)	0.43336 (14)	0.35509 (12)	0.0498 (3)
N2	0.2094 (5)	0.5182 (3)	0.0809 (3)	0.0340 (8)
C5	0.3021 (5)	0.5258 (4)	-0.0900 (4)	0.0328 (10)
N3	0.0948 (5)	0.3267 (4)	0.1271 (4)	0.0426 (10)
N1	0.3815 (5)	0.7322 (4)	-0.1296 (3)	0.0387 (9)
C9	0.1455 (5)	0.4513 (4)	0.1466 (4)	0.0340 (10)
N5	0.1235 (5)	0.2820 (4)	0.4999 (3)	0.0381 (9)

C1	0.3142 (6)	0.6574 (4)	-0.0705 (4)	0.0356 (10)
H1A	0.2728	0.6955	-0.0127	0.043*
C6	0.2284 (5)	0.4523 (4)	-0.0183 (4)	0.0330 (10)
C10	0.1797 (6)	0.3566 (5)	0.4357 (4)	0.0392 (11)
C3	0.4346 (6)	0.5483 (5)	-0.2380 (4)	0.0436 (11)
H3B	0.4772	0.5130	-0.2963	0.052*
C13	0.2302 (6)	0.2187 (4)	0.5638 (4)	0.0381 (10)
C12	0.3936 (6)	0.2331 (5)	0.5625 (4)	0.0480 (13)
H12A	0.4704	0.1894	0.6067	0.058*
C4	0.3639 (6)	0.4719 (5)	-0.1776 (4)	0.0411 (11)
H4A	0.3572	0.3842	-0.1951	0.049*
N4	0.3312 (5)	0.3777 (4)	0.4277 (3)	0.0453 (10)
C11	0.4371 (6)	0.3147 (5)	0.4933 (5)	0.0515 (13)
H11A	0.5461	0.3261	0.4925	0.062*
C2	0.4419 (6)	0.6772 (5)	-0.2117 (4)	0.0418 (11)
H2B	0.4909	0.7284	-0.2528	0.050*
C8	0.1115 (6)	0.2636 (5)	0.0281 (5)	0.0457 (12)
H8A	0.0766	0.1761	0.0087	0.055*
C7	0.1775 (6)	0.3213 (4)	-0.0466 (4)	0.0416 (11)
H7A	0.1883	0.2739	-0.1148	0.050*
C14	0.1671 (6)	0.1341 (4)	0.6336 (4)	0.0380 (10)
C15	-0.0011 (6)	0.1112 (5)	0.6209 (4)	0.0475 (12)
H15A	-0.0761	0.1500	0.5691	0.057*
C16	-0.0545 (7)	0.0301 (5)	0.6864 (5)	0.0532 (14)
H16A	-0.1663	0.0141	0.6796	0.064*
C18	0.2701 (7)	0.0733 (5)	0.7125 (4)	0.0456 (12)
H18A	0.3827	0.0880	0.7217	0.055*
C17	0.0554 (7)	-0.0264 (5)	0.7604 (4)	0.0503 (13)
H17A	0.0161	-0.0825	0.8023	0.060*
N6	0.2191 (5)	-0.0053 (4)	0.7765 (3)	0.0442 (10)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
II	0.0504 (2)	0.0489 (2)	0.0526 (2)	0.00536 (15)	0.02987 (17)	0.01773 (16)
Cu1	0.0647 (4)	0.0437 (4)	0.0576 (4)	0.0113 (3)	0.0319 (3)	0.0285 (3)
S1	0.0595 (8)	0.0464 (7)	0.0346 (6)	0.0117 (6)	0.0179 (6)	0.0187 (5)
S2	0.0478 (7)	0.0725 (9)	0.0424 (7)	0.0181 (7)	0.0204 (6)	0.0338 (7)
N2	0.039 (2)	0.034 (2)	0.0308 (19)	0.0039 (16)	0.0112 (16)	0.0143 (16)
C5	0.031 (2)	0.036 (2)	0.031 (2)	0.0029 (19)	0.0078 (18)	0.0140 (19)
N3	0.045 (2)	0.040 (2)	0.049 (2)	0.0049 (18)	0.0167 (19)	0.0225 (19)
N1	0.044 (2)	0.038 (2)	0.038 (2)	0.0031 (17)	0.0182 (18)	0.0159 (17)
C9	0.035 (2)	0.040 (3)	0.028 (2)	0.008 (2)	0.0081 (19)	0.0147 (19)
N5	0.042 (2)	0.046 (2)	0.0293 (19)	0.0054 (18)	0.0114 (17)	0.0156 (17)
C1	0.042 (3)	0.035 (2)	0.036 (2)	0.007 (2)	0.018 (2)	0.012 (2)
C6	0.033 (2)	0.034 (2)	0.033 (2)	0.0050 (19)	0.0091 (19)	0.0131 (19)
C10	0.044 (3)	0.045 (3)	0.030 (2)	0.004 (2)	0.012 (2)	0.012 (2)
C3	0.046 (3)	0.053 (3)	0.036 (2)	0.008 (2)	0.018 (2)	0.008 (2)

supplementary materials

C13	0.046 (3)	0.040 (3)	0.029 (2)	0.004 (2)	0.011 (2)	0.012 (2)
C12	0.041 (3)	0.060 (3)	0.047 (3)	0.012 (2)	0.010 (2)	0.030 (3)
C4	0.046 (3)	0.039 (3)	0.039 (3)	0.005 (2)	0.015 (2)	0.008 (2)
N4	0.043 (2)	0.057 (3)	0.039 (2)	0.006 (2)	0.0127 (18)	0.021 (2)
C11	0.038 (3)	0.069 (4)	0.051 (3)	0.001 (3)	0.013 (2)	0.027 (3)
C2	0.046 (3)	0.048 (3)	0.036 (3)	0.000 (2)	0.018 (2)	0.016 (2)
C8	0.051 (3)	0.034 (3)	0.052 (3)	0.004 (2)	0.015 (2)	0.014 (2)
C7	0.056 (3)	0.031 (2)	0.039 (3)	0.006 (2)	0.017 (2)	0.010 (2)
C14	0.046 (3)	0.037 (2)	0.033 (2)	0.005 (2)	0.016 (2)	0.010 (2)
C15	0.046 (3)	0.059 (3)	0.040 (3)	0.004 (2)	0.014 (2)	0.017 (2)
C16	0.050 (3)	0.070 (4)	0.045 (3)	0.001 (3)	0.022 (2)	0.018 (3)
C18	0.049 (3)	0.048 (3)	0.045 (3)	0.004 (2)	0.018 (2)	0.021 (2)
C17	0.064 (3)	0.048 (3)	0.047 (3)	-0.002 (3)	0.029 (3)	0.018 (2)
N6	0.054 (3)	0.042 (2)	0.041 (2)	0.0064 (19)	0.0178 (19)	0.0175 (18)

Geometric parameters (\AA , $^\circ$)

Cu1—I1	2.6550 (7)	C3—H3B	0.9300
Cu1—I1 ⁱ	2.6579 (8)	C13—C12	1.394 (7)
Cu1—N1	2.037 (4)	C13—C14	1.477 (6)
Cu1—N6 ⁱⁱ	2.060 (4)	C12—C11	1.379 (7)
Cu1—Cu1 ⁱ	2.6935 (14)	C12—H12A	0.9300
S1—C9	1.779 (5)	C4—H4A	0.9300
S1—S2	2.0183 (17)	N4—C11	1.327 (6)
S2—C10	1.778 (5)	C11—H11A	0.9300
N2—C9	1.323 (5)	C2—H2B	0.9300
N2—C6	1.352 (6)	C8—C7	1.367 (6)
C5—C1	1.390 (6)	C8—H8A	0.9300
C5—C4	1.389 (6)	C7—H7A	0.9300
C5—C6	1.466 (6)	C14—C18	1.379 (7)
N3—C9	1.329 (6)	C14—C15	1.392 (7)
N3—C8	1.329 (6)	C15—C16	1.376 (7)
N1—C2	1.333 (6)	C15—H15A	0.9300
N1—C1	1.330 (5)	C16—C17	1.349 (7)
N5—C13	1.324 (6)	C16—H16A	0.9300
N5—C10	1.328 (6)	C18—N6	1.336 (6)
C1—H1A	0.9300	C18—H18A	0.9300
C6—C7	1.391 (6)	C17—N6	1.344 (7)
C10—N4	1.325 (6)	C17—H17A	0.9300
C3—C2	1.370 (7)	N6—Cu1 ⁱⁱⁱ	2.060 (4)
C3—C4	1.368 (6)		
Cu1—I1—Cu1 ⁱ	60.92 (3)	C12—C13—C14	123.0 (4)
N1—Cu1—N6 ⁱⁱ	115.68 (16)	C11—C12—C13	117.4 (4)
N1—Cu1—I1	106.36 (10)	C11—C12—H12A	121.3
N6 ⁱⁱ —Cu1—I1	106.25 (12)	C13—C12—H12A	121.3
N1—Cu1—I1 ⁱ	105.02 (11)	C3—C4—C5	119.8 (4)
N6 ⁱⁱ —Cu1—I1 ⁱ	104.97 (12)	C3—C4—H4A	120.1

I1—Cu1—I1 ⁱ	119.08 (3)	C5—C4—H4A	120.1
N1—Cu1—Cu1 ⁱ	122.23 (11)	C10—N4—C11	113.9 (4)
N6 ⁱⁱ —Cu1—Cu1 ⁱ	122.06 (12)	N4—C11—C12	123.3 (5)
I1—Cu1—Cu1 ⁱ	59.59 (2)	N4—C11—H11A	118.3
I1 ⁱ —Cu1—Cu1 ⁱ	59.49 (3)	C12—C11—H11A	118.3
C9—S1—S2	104.18 (16)	N1—C2—C3	122.6 (4)
C10—S2—S1	104.60 (17)	N1—C2—H2B	118.7
C9—N2—C6	116.6 (4)	C3—C2—H2B	118.7
C1—C5—C4	116.6 (4)	C7—C8—N3	123.1 (4)
C1—C5—C6	119.6 (4)	C7—C8—H8A	118.5
C4—C5—C6	123.8 (4)	N3—C8—H8A	118.5
C9—N3—C8	114.2 (4)	C8—C7—C6	118.6 (4)
C2—N1—C1	117.8 (4)	C8—C7—H7A	120.7
C2—N1—Cu1	121.4 (3)	C6—C7—H7A	120.7
C1—N1—Cu1	120.8 (3)	C18—C14—C15	117.0 (4)
N2—C9—N3	128.4 (4)	C18—C14—C13	122.1 (4)
N2—C9—S1	110.9 (3)	C15—C14—C13	120.8 (4)
N3—C9—S1	120.7 (3)	C14—C15—C16	118.8 (5)
C13—N5—C10	116.9 (4)	C14—C15—H15A	120.6
N1—C1—C5	123.9 (4)	C16—C15—H15A	120.6
N1—C1—H1A	118.0	C17—C16—C15	120.0 (5)
C5—C1—H1A	118.0	C17—C16—H16A	120.0
N2—C6—C7	119.1 (4)	C15—C16—H16A	120.0
N2—C6—C5	116.7 (4)	N6—C18—C14	124.5 (5)
C7—C6—C5	124.2 (4)	N6—C18—H18A	117.8
N4—C10—N5	128.4 (4)	C14—C18—H18A	117.8
N4—C10—S2	120.6 (3)	C16—C17—N6	123.1 (5)
N5—C10—S2	111.1 (3)	C16—C17—H17A	118.5
C2—C3—C4	119.3 (4)	N6—C17—H17A	118.5
C2—C3—H3B	120.4	C18—N6—C17	116.6 (4)
C4—C3—H3B	120.4	C18—N6—Cu1 ⁱⁱⁱ	120.4 (4)
N5—C13—C12	120.0 (4)	C17—N6—Cu1 ⁱⁱⁱ	122.7 (3)
N5—C13—C14	117.0 (4)		
Cu1 ⁱ —I1—Cu1—N1	118.18 (12)	C10—N5—C13—C12	0.7 (7)
Cu1 ⁱ —I1—Cu1—N6 ⁱⁱ	-118.04 (12)	C10—N5—C13—C14	-179.0 (4)
Cu1 ⁱ —I1—Cu1—I1 ⁱ	0.0	N5—C13—C12—C11	0.1 (7)
C9—S1—S2—C10	81.1 (2)	C14—C13—C12—C11	179.7 (4)
N6 ⁱⁱ —Cu1—N1—C2	75.1 (4)	C2—C3—C4—C5	-0.6 (7)
I1—Cu1—N1—C2	-167.2 (3)	C1—C5—C4—C3	1.1 (6)
I1 ⁱ —Cu1—N1—C2	-40.1 (4)	C6—C5—C4—C3	-178.3 (4)
Cu1 ⁱ —Cu1—N1—C2	-103.2 (3)	N5—C10—N4—C11	0.0 (8)
N6 ⁱⁱ —Cu1—N1—C1	-104.0 (4)	S2—C10—N4—C11	179.6 (4)
I1—Cu1—N1—C1	13.7 (4)	C10—N4—C11—C12	0.8 (8)
I1 ⁱ —Cu1—N1—C1	140.8 (3)	C13—C12—C11—N4	-0.9 (8)
Cu1 ⁱ —Cu1—N1—C1	77.7 (4)	C1—N1—C2—C3	1.0 (7)

supplementary materials

C6—N2—C9—N3	-1.0 (7)	Cu1—N1—C2—C3	-178.1 (4)
C6—N2—C9—S1	178.0 (3)	C4—C3—C2—N1	-0.5 (7)
C8—N3—C9—N2	-0.2 (7)	C9—N3—C8—C7	1.0 (7)
C8—N3—C9—S1	-179.1 (3)	N3—C8—C7—C6	-0.6 (8)
S2—S1—C9—N2	176.0 (3)	N2—C6—C7—C8	-0.7 (7)
S2—S1—C9—N3	-4.8 (4)	C5—C6—C7—C8	179.6 (4)
C2—N1—C1—C5	-0.5 (7)	N5—C13—C14—C18	-173.4 (4)
Cu1—N1—C1—C5	178.7 (3)	C12—C13—C14—C18	6.9 (7)
C4—C5—C1—N1	-0.5 (7)	N5—C13—C14—C15	7.5 (7)
C6—C5—C1—N1	178.9 (4)	C12—C13—C14—C15	-172.2 (5)
C9—N2—C6—C7	1.4 (6)	C18—C14—C15—C16	-0.2 (7)
C9—N2—C6—C5	-178.8 (4)	C13—C14—C15—C16	178.9 (4)
C1—C5—C6—N2	-14.3 (6)	C14—C15—C16—C17	-0.6 (8)
C4—C5—C6—N2	165.0 (4)	C15—C14—C18—N6	0.0 (7)
C1—C5—C6—C7	165.5 (4)	C13—C14—C18—N6	-179.1 (4)
C4—C5—C6—C7	-15.2 (7)	C15—C16—C17—N6	1.6 (8)
C13—N5—C10—N4	-0.8 (7)	C14—C18—N6—C17	0.9 (7)
C13—N5—C10—S2	179.6 (3)	C14—C18—N6—Cu1 ⁱⁱⁱ	-174.0 (4)
S1—S2—C10—N4	-4.1 (4)	C16—C17—N6—C18	-1.7 (8)
S1—S2—C10—N5	175.5 (3)	C16—C17—N6—Cu1 ⁱⁱⁱ	173.0 (4)

Symmetry codes: (i) $-x+1, -y+2, -z$; (ii) $x, y+1, z-1$; (iii) $x, y-1, z+1$.

Fig. 1




Fig. 2

