

## Diiiodido(1,10-phenanthroline- $\kappa^2 N,N'$ )-platinum(II)

Kwang Ha

School of Applied Chemical Engineering, The Research Institute of Catalysis,  
Chonnam National University, Gwangju 500-757, Republic of Korea  
Correspondence e-mail: hakwang@chonnam.ac.kr


Received 1 December 2009; accepted 10 December 2009

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(C-C) = 0.012$  Å;  
 $R$  factor = 0.033;  $wR$  factor = 0.084; data-to-parameter ratio = 14.8.

In the title complex,  $[PtI_2(C_{12}H_8N_2)]$ , the  $Pt^{2+}$  ion is four-coordinated in a slightly distorted square-planar environment by two N atoms of the chelating 1,10-phenanthroline ligand and two iodide ions. The nearly planar molecules, with a maximum deviation of 0.170 (3) Å from the least-squares plane, are stacked in columns along the  $c$  axis with a  $Pt \cdots Pt$  distance of 4.8510 (6) Å. In the column,  $\pi-\pi$  interactions between adjacent six-membered rings are present, the shortest centroid–centroid distance being 3.703 (5) Å.

### Related literature

For the syntheses of  $[PtX_2(\text{phen})]$  ( $\text{phen} = 1,10\text{-phenanthroline}$ ;  $X = \text{Cl}$ , Br or I), see: Hodges & Rund (1975). For the crystal structure of yellow  $[PtCl_2(\text{phen})]$  which is isotypic to the title complex, see: Grzesiak & Matzger (2007).



### Experimental

#### Crystal data

$[PtI_2(C_{12}H_8N_2)]$

$M_r = 629.09$

Monoclinic,  $P2_1/c$   
 $a = 10.3284$  (9) Å  
 $b = 17.9462$  (16) Å  
 $c = 7.3833$  (7) Å  
 $\beta = 108.569$  (2)°  
 $V = 1297.3$  (2) Å<sup>3</sup>

$Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 15.55$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.32 \times 0.13 \times 0.08$  mm

#### Data collection

Bruker SMART 1000 CCD  
diffractometer  
Absorption correction: multi-scan  
(SADABS; Bruker, 2000)  
 $T_{\min} = 0.422$ ,  $T_{\max} = 1.000$

7331 measured reflections  
2284 independent reflections  
2110 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.084$   
 $S = 1.06$   
2284 reflections

154 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 3.05$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -1.40$  e Å<sup>-3</sup>

**Table 1**  
Selected geometric parameters (Å, °).

Pt1–N2	2.039 (6)	Pt1–I2	2.5774 (7)
Pt1–N1	2.060 (7)	Pt1–I1	2.5847 (6)
N2–Pt1–N1			80.6 (2)

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINT* (Bruker, 2000); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2009–0094056).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2700).

### References

- Bruker (2000). *SADABS*, *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Grzesiak, A. L. & Matzger, A. J. (2007). *Inorg. Chem.* **46**, 453–457.
- Hodges, K. D. & Rund, J. V. (1975). *Inorg. Chem.* **14**, 525–528.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2010). E66, m65 [doi:10.1107/S1600536809053100]

## Diiodido(1,10-phenanthroline- $\kappa^2 N,N'$ )platinum(II)

K. Ha

### Comment

The title complex,  $[\text{PtI}_2(\text{phen})]$  (where phen is 1,10-phenanthroline,  $C_{12}\text{H}_8\text{N}_2$ ), is isomorphous with the yellow form of  $[\text{PtCl}_2(\text{phen})]$ , whereas the orange form of  $[\text{PtCl}_2(\text{phen})]$  crystallized in the orthorhombic space group  $Pca2_1$  (Grzesiak & Matzger, 2007).

In the title complex, the  $\text{Pt}^{2+}$  ion is four-coordinated in a slightly distorted square-planar environment by two N atoms of the chelating 1,10-phenanthroline ligand and two iodide ions (Fig. 1). The main contribution to the distortion is the tight  $\text{N}1-\text{Pt}1-\text{N}2$  chelate angle [ $80.6(2)^\circ$ ], which results in non-linear *trans* arrangement [ $\angle \text{N}1-\text{Pt}1-\text{I}1 = 175.72(17)^\circ$  and  $\angle \text{N}2-\text{Pt}1-\text{I}2 = 175.02(17)^\circ$ ]. The  $\text{Pt}1-\text{N}$  and  $\text{Pt}1-\text{I}$  bond lengths are almost equal, respectively [ $\text{Pt}1-\text{N}: 2.060(7)$  and  $2.039(6)$  Å;  $\text{Pt}1-\text{I} 2.5847(6)$  and  $2.5774(7)$  Å] (Table 1). The complex displays numerous intermolecular  $\pi-\pi$  interactions between adjacent six-membered rings, with a shortest centroid-centroid distance of  $3.703(5)$  Å and the dihedral angle between the ring planes is  $3.4(4)^\circ$ . The nearly planar  $[\text{PtI}_2(\text{phen})]$  molecules, with a largest deviation of  $0.170(3)$  Å from the least-squares plane, stack in columns along the *c* axis with a  $\text{Pt}\cdots\text{Pt}$  distance of  $4.8510(6)$  Å (Fig. 2).

### Experimental

To a solution of  $\text{K}_2\text{PtCl}_4$  (0.2011 g, 0.484 mmol) in  $\text{H}_2\text{O}$  (20 ml) were added KI (1.6010 g, 9.644 mmol) and 1,10-phenanthroline (0.0967 g, 0.537 mmol), and refluxed for 3 h. The precipitate obtained was separated by filtration, washed with water and acetone, and dried at  $100\text{ }^\circ\text{C}$ , to give a dark yellow powder (0.2732 g). Crystals suitable for X-ray analysis were obtained by slow evaporation from an *N,N*-dimethylformamide solution at  $50\text{ }^\circ\text{C}$ .

### Refinement

H atoms were positioned geometrically and allowed to ride on their respective parent atoms [ $\text{C}-\text{H} = 0.95$  Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ]. The maximum and minimum residual electron density peaks of  $3.05$  and  $-1.40$  e Å $^{-3}$ , respectively, were located  $0.97$  and  $0.92$  Å from the Pt Atom.

### Figures




Fig. 1. The structure of the title complex, with displacement ellipsoids drawn at the 50% probability level for non-H atoms.

## supplementary materials

---




Fig. 2. Crystal packing of the title complex.

### Diiodido(1,10-phenanthroline- $\kappa^2 N,N'$ )platinum(II)

#### Crystal data

[PtI <sub>2</sub> (C <sub>12</sub> H <sub>8</sub> N <sub>2</sub> )]	$F(000) = 1112$
$M_r = 629.09$	$D_x = 3.221 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 5463 reflections
$a = 10.3284 (9) \text{ \AA}$	$\theta = 2.3\text{--}28.2^\circ$
$b = 17.9462 (16) \text{ \AA}$	$\mu = 15.55 \text{ mm}^{-1}$
$c = 7.3833 (7) \text{ \AA}$	$T = 200 \text{ K}$
$\beta = 108.569 (2)^\circ$	Needle, yellow
$V = 1297.3 (2) \text{ \AA}^3$	$0.32 \times 0.13 \times 0.08 \text{ mm}$
$Z = 4$	

#### Data collection

Bruker SMART 1000 CCD diffractometer	2284 independent reflections
Radiation source: fine-focus sealed tube graphite	2110 reflections with $I > 2\sigma(I)$
$\varphi$ and $\omega$ scans	$R_{\text{int}} = 0.033$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2000)	$\theta_{\text{max}} = 25.1^\circ, \theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.422, T_{\text{max}} = 1.000$	$h = -11 \rightarrow 12$
7331 measured reflections	$k = -21 \rightarrow 21$
	$l = -8 \rightarrow 8$

#### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.033$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.084$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_{\text{o}}^2) + (0.0524P)^2 + 0.9657P]$

	where $P = (F_o^2 + 2F_c^2)/3$
2284 reflections	$(\Delta/\sigma)_{\max} < 0.001$
154 parameters	$\Delta\rho_{\max} = 3.05 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -1.40 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Pt1	0.71220 (3)	0.337678 (16)	0.39502 (4)	0.02011 (13)
I1	0.74244 (6)	0.48072 (3)	0.41617 (8)	0.03146 (17)
I2	0.45788 (6)	0.35783 (3)	0.20604 (9)	0.03744 (18)
N1	0.7037 (6)	0.2230 (4)	0.3858 (8)	0.0245 (15)
N2	0.9082 (6)	0.3142 (4)	0.5563 (9)	0.0219 (13)
C1	0.6012 (9)	0.1789 (5)	0.2965 (12)	0.034 (2)
H1	0.5167	0.2005	0.2237	0.041*
C2	0.6130 (9)	0.1009 (5)	0.3056 (13)	0.035 (2)
H2	0.5376	0.0709	0.2367	0.042*
C3	0.7305 (10)	0.0681 (6)	0.4114 (11)	0.036 (2)
H3	0.7375	0.0153	0.4195	0.044*
C4	0.8426 (9)	0.1132 (4)	0.5097 (12)	0.0291 (19)
C5	0.9726 (10)	0.0861 (5)	0.6267 (13)	0.034 (2)
H5	0.9862	0.0338	0.6419	0.041*
C6	1.0755 (9)	0.1317 (5)	0.7155 (13)	0.0317 (19)
H6	1.1602	0.1114	0.7921	0.038*
C7	1.0598 (8)	0.2110 (4)	0.6964 (11)	0.0248 (17)
C8	1.1647 (8)	0.2622 (5)	0.7825 (11)	0.0311 (19)
H8	1.2519	0.2453	0.8600	0.037*
C9	1.1388 (9)	0.3359 (4)	0.7527 (12)	0.0300 (19)
H9	1.2091	0.3708	0.8101	0.036*
C10	1.0137 (8)	0.3614 (5)	0.6417 (11)	0.0261 (17)
H10	0.9999	0.4135	0.6236	0.031*
C11	0.9351 (8)	0.2394 (4)	0.5838 (10)	0.0208 (16)
C12	0.8236 (8)	0.1915 (4)	0.4917 (11)	0.0233 (17)

## supplementary materials

---

### *Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Pt1	0.0168 (2)	0.0209 (2)	0.0209 (2)	-0.00133 (10)	0.00349 (14)	-0.00034 (10)
I1	0.0309 (3)	0.0217 (3)	0.0393 (3)	0.0010 (2)	0.0077 (3)	0.0001 (2)
I2	0.0184 (3)	0.0439 (4)	0.0433 (4)	0.0001 (2)	0.0004 (2)	0.0026 (3)
N1	0.024 (4)	0.028 (4)	0.023 (3)	-0.004 (3)	0.010 (3)	-0.002 (3)
N2	0.018 (4)	0.025 (3)	0.022 (3)	0.001 (3)	0.006 (3)	0.001 (3)
C1	0.031 (5)	0.040 (5)	0.032 (5)	-0.019 (4)	0.010 (4)	-0.012 (4)
C2	0.030 (5)	0.034 (5)	0.041 (5)	-0.015 (4)	0.011 (4)	-0.007 (4)
C3	0.059 (7)	0.026 (5)	0.031 (5)	-0.006 (4)	0.025 (5)	-0.005 (3)
C4	0.043 (5)	0.021 (4)	0.031 (4)	-0.005 (4)	0.023 (4)	0.001 (3)
C5	0.048 (6)	0.024 (5)	0.038 (5)	0.009 (4)	0.023 (5)	0.002 (4)
C6	0.028 (5)	0.032 (5)	0.038 (5)	0.003 (4)	0.014 (4)	0.004 (4)
C7	0.019 (4)	0.030 (4)	0.027 (4)	0.007 (3)	0.009 (3)	0.003 (3)
C8	0.020 (4)	0.046 (5)	0.025 (4)	0.010 (4)	0.005 (3)	0.006 (4)
C9	0.026 (5)	0.033 (5)	0.025 (4)	-0.005 (3)	0.000 (4)	-0.001 (3)
C10	0.017 (4)	0.023 (4)	0.032 (4)	-0.008 (3)	-0.001 (3)	-0.003 (3)
C11	0.024 (4)	0.019 (4)	0.021 (4)	-0.003 (3)	0.010 (3)	0.000 (3)
C12	0.033 (5)	0.021 (4)	0.022 (4)	-0.001 (3)	0.016 (4)	0.003 (3)

### *Geometric parameters ( $\text{\AA}$ , $^\circ$ )*

Pt1—N2	2.039 (6)	C4—C12	1.419 (11)
Pt1—N1	2.060 (7)	C4—C5	1.431 (13)
Pt1—I2	2.5774 (7)	C5—C6	1.337 (12)
Pt1—I1	2.5847 (6)	C5—H5	0.9500
N1—C1	1.319 (10)	C6—C7	1.434 (11)
N1—C12	1.360 (10)	C6—H6	0.9500
N2—C10	1.365 (10)	C7—C11	1.389 (11)
N2—C11	1.374 (10)	C7—C8	1.407 (12)
C1—C2	1.405 (13)	C8—C9	1.354 (12)
C1—H1	0.9500	C8—H8	0.9500
C2—C3	1.351 (13)	C9—C10	1.369 (12)
C2—H2	0.9500	C9—H9	0.9500
C3—C4	1.410 (12)	C10—H10	0.9500
C3—H3	0.9500	C11—C12	1.423 (11)
N2—Pt1—N1	80.6 (2)	C6—C5—C4	122.3 (8)
N2—Pt1—I2	175.02 (17)	C6—C5—H5	118.9
N1—Pt1—I2	95.57 (17)	C4—C5—H5	118.9
N2—Pt1—I1	95.23 (18)	C5—C6—C7	120.8 (8)
N1—Pt1—I1	175.72 (17)	C5—C6—H6	119.6
I2—Pt1—I1	88.63 (2)	C7—C6—H6	119.6
C1—N1—C12	118.5 (8)	C11—C7—C8	117.7 (7)
C1—N1—Pt1	129.5 (6)	C11—C7—C6	118.5 (7)
C12—N1—Pt1	112.1 (5)	C8—C7—C6	123.8 (7)
C10—N2—C11	116.4 (7)	C9—C8—C7	118.7 (8)

C10—N2—Pt1	129.8 (5)	C9—C8—H8	120.6
C11—N2—Pt1	113.8 (5)	C7—C8—H8	120.6
N1—C1—C2	122.1 (9)	C8—C9—C10	121.5 (8)
N1—C1—H1	119.0	C8—C9—H9	119.2
C2—C1—H1	119.0	C10—C9—H9	119.2
C3—C2—C1	120.6 (8)	N2—C10—C9	122.1 (8)
C3—C2—H2	119.7	N2—C10—H10	118.9
C1—C2—H2	119.7	C9—C10—H10	118.9
C2—C3—C4	119.2 (9)	N2—C11—C7	123.5 (7)
C2—C3—H3	120.4	N2—C11—C12	115.2 (7)
C4—C3—H3	120.4	C7—C11—C12	121.3 (7)
C3—C4—C12	117.0 (8)	N1—C12—C4	122.6 (7)
C3—C4—C5	125.0 (8)	N1—C12—C11	118.2 (7)
C12—C4—C5	117.9 (7)	C4—C12—C11	119.1 (7)
N2—Pt1—N1—C1	-178.5 (7)	Pt1—N2—C10—C9	177.2 (6)
I2—Pt1—N1—C1	4.8 (7)	C8—C9—C10—N2	0.4 (13)
N2—Pt1—N1—C12	2.5 (5)	C10—N2—C11—C7	2.1 (10)
I2—Pt1—N1—C12	-174.3 (4)	Pt1—N2—C11—C7	-176.9 (6)
N1—Pt1—N2—C10	179.0 (7)	C10—N2—C11—C12	-179.5 (6)
I1—Pt1—N2—C10	0.1 (7)	Pt1—N2—C11—C12	1.6 (8)
N1—Pt1—N2—C11	-2.2 (5)	C8—C7—C11—N2	-1.6 (11)
I1—Pt1—N2—C11	178.8 (4)	C6—C7—C11—N2	179.6 (7)
C12—N1—C1—C2	-0.9 (11)	C8—C7—C11—C12	-179.9 (7)
Pt1—N1—C1—C2	-179.9 (6)	C6—C7—C11—C12	1.2 (10)
N1—C1—C2—C3	1.6 (12)	C1—N1—C12—C4	0.0 (10)
C1—C2—C3—C4	-1.4 (12)	Pt1—N1—C12—C4	179.2 (5)
C2—C3—C4—C12	0.5 (11)	C1—N1—C12—C11	178.4 (7)
C2—C3—C4—C5	179.9 (8)	Pt1—N1—C12—C11	-2.4 (8)
C3—C4—C5—C6	179.5 (7)	C3—C4—C12—N1	0.2 (10)
C12—C4—C5—C6	-1.1 (12)	C5—C4—C12—N1	-179.3 (7)
C4—C5—C6—C7	-0.2 (12)	C3—C4—C12—C11	-178.2 (7)
C5—C6—C7—C11	0.1 (11)	C5—C4—C12—C11	2.4 (10)
C5—C6—C7—C8	-178.6 (8)	N2—C11—C12—N1	0.6 (10)
C11—C7—C8—C9	0.4 (11)	C7—C11—C12—N1	179.1 (6)
C6—C7—C8—C9	179.2 (7)	N2—C11—C12—C4	179.0 (6)
C7—C8—C9—C10	0.1 (12)	C7—C11—C12—C4	-2.5 (10)
C11—N2—C10—C9	-1.5 (11)		

## supplementary materials

**Fig. 1**




Fig. 2

