

## Aqua(6,6'-oxydipicolinato- $\kappa^2O,N,N',O'$ )-copper(II)

**Jingya Sun\* and Xiangdi Tong**

College of Marine Sciences, Zhejiang Ocean University, Zhoushan 316000, People's Republic of China

Correspondence e-mail: jingyasun2009@163.com


Received 30 November 2009; accepted 11 December 2009

 Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(C-C) = 0.005$  Å;  $R$  factor = 0.032;  $wR$  factor = 0.110; data-to-parameter ratio = 10.9.

In the title complex,  $[Cu(C_{12}H_6N_2O_5)(H_2O)]$ , the  $Cu^{II}$  ion is in a slightly distorted square-pyramidal coordination environment with two N and two O atoms from a 6,6'-oxydipicolinate ligand occupying the basal plane and a water ligand in the apical site. The dihedral angle between the two pyridine rings is  $5.51(6)^\circ$ . In the crystal structure, intermolecular  $O-H \cdots O$  hydrogen bonds link molecules into a two-dimensional network. In addition, weak intermolecular  $C-H \cdots O$  and  $C=O(\text{lone pair}) \cdots \pi(\text{ring})$  interactions, with  $O \cdots$  ring-centroid distances of  $3.697(4)$  and  $3.094(4)$  Å, provide additional stabilization.

### Related literature

For intermolecular interactions, see: Choudhury *et al.* (2008). For the applications of picolinic acid compounds, see: Mann *et al.* (1992).



### Experimental

#### Crystal data

 $[Cu(C_{12}H_6N_2O_5)(H_2O)]$   
 $M_r = 339.74$ 

 Monoclinic,  $P2_1/n$ 
 $a = 7.2487(16)$  Å

 $b = 21.055(4)$  Å

 $c = 8.2269(17)$  Å

 $\beta = 110.201(9)^\circ$ 
 $V = 1178.4(4)$  Å<sup>3</sup>
 $Z = 4$ 

 Mo  $K\alpha$  radiation

 $\mu = 1.89$  mm<sup>-1</sup>
 $T = 296$  K

 $0.40 \times 0.35 \times 0.30$  mm

#### Data collection

 Siemens SMART CCD  
 diffractometer

 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)

 $T_{\min} = 0.519$ ,  $T_{\max} = 0.602$ 

6790 measured reflections

2074 independent reflections

 1806 reflections with  $I > 2\sigma(I)$ 
 $R_{\text{int}} = 0.027$ 

#### Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.032$ 
 $wR(F^2) = 0.110$ 
 $S = 1.18$ 

2074 reflections

190 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.48$  e Å<sup>-3</sup>
 $\Delta\rho_{\text{min}} = -0.37$  e Å<sup>-3</sup>
**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$C8-H8 \cdots O4^i$	0.93	2.31	3.229 (4)	171
$C9-H9 \cdots O2^{ii}$	0.93	2.42	3.331 (4)	165
$C4-H4 \cdots O6^{iii}$	0.93	2.54	3.303 (4)	140
$O6-H6B \cdots O4^{iv}$	0.85	1.97	2.772 (3)	157
$O6-H6A \cdots O2^v$	0.85	2.01	2.807 (3)	156

Symmetry codes: (i)  $x, y, z - 1$ ; (ii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{3}{2}$ ; (iii)  $x + \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ ; (iv)  $-x, -y, -z + 2$ ; (v)  $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: PLATON (Spek, 2009) and SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2964).

### References

- Choudhury, S. R., Gamez, P., Robertazzi, A., Chen, C. Y., Lee, H. M. & Mukhopadhyay, S. (2008). *Cryst. Growth Des.* **8**, 3773–3784.
- Mann, Y., Chiment, F., Balasco, A., Cenicola, M. L., Amico, M. D., Parrilo, C., Rossi, F. & Marmo, E. (1992). *Eur. J. Med. Chem.* **27**, 633–639.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

**supplementary materials**

*Acta Cryst.* (2010). E66, m70 [ doi:10.1107/S160053680905346X ]

## Aqua(6,6'-oxydipicolinato- $\kappa^2O,N,N',O'$ )copper(II)

J. Sun and X. Tong

### Comment

Picolinic acid compounds play an vital role in the development of coordination chemistry related to catalysis, magnetism and molecular architectures (Mann *et al.*, 1992). As part of our studies on the synthesis and characterization of these types of compounds, we report here the synthesis and crystal structure of the title compound (I).

The molecular structure of the title compound (I) is shown in Fig. 1. The Cu<sup>II</sup> ion is in a slightly distorted square-pyramidal coordination environment with two N and two O atoms from a 6,6'-oxydipicolinato ligand occupying the basal plane and one water ligand in the apical site. The dihedral angle between the two pyridine rings is 5.51 (6)°. The delocalization of electrons within the carboxylate groups is reflected in the C=O lengths. In the crystal structure, there are intermolecular O—H...O hydrogen bonds involving the carboxyl oxygen atoms and coordinated water molecules (Fig. 2) forming a two-dimensional network (see Table 1 for hydrogen bond geometries). In addition to weak intermolecular C-H...O interactions, further stabilization appears to be provided by weak C=O(lone pair)... $\pi$ (ring) stacking interactions (Choudhury *et al.*, 2008). The relevant distances are C12—O4...Cg1<sup>i</sup> = 3.697 (4) Å, Cg1 is the centroid of the ring defined by the atoms N1/C7-C11 [symmetry code: (i) -x, -y, 2-z] and the angle C12—O4...Cg1<sup>i</sup> is 98.95 (34)°; C1—O2...Cg2<sup>ii</sup> = 3.094 (4) Å, Cg2 is the centroid of the ring defined by the atoms N2/C2-C6 [symmetry code: (ii) 0.5+x, 0.5-y, 0.5+z] and the angle C1—O2...Cg2<sup>ii</sup> is 115.48 (4)° (see Fig. 3).

### Experimental

All reagents were available commercially and were used without further purification. 6,6'-Oxydipicolinic acid (260 mg) was added to 1 mmol (132 mg) of CuCl<sub>2</sub> in 10 ml of water. The suspension was stirred for 4 h and filtered. After leaving the filtrate in air for one week, blue block-shaped crystals of (I) were formed. The crystals were isolated, washed with water three times and dried in a vacuum desiccator using silica gel (Yield 75%). Elemental analysis: found C, 42.05; H, 2.96; N, 8.18%; calc. for C<sub>12</sub>H<sub>8</sub>CuN<sub>2</sub>O<sub>6</sub>; C, 42.17; H, 2.95; N, 8.20%.

### Refinement

H atoms bonded to C atoms were positioned geometrically and refined using a riding-model approximation with C—H = 0.93 Å, and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . H atoms bonded to O atoms were found in difference Fourier maps and included as riding with O—H = 0.85 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$ .

## Figures




Fig. 1. The molecular structure of (I) showing 50% probability displacement ellipsoids and the atom-numbering scheme.




Fig. 2. Part of the crystal structure of (I) showing hydrogen bonds as dashed lines.



Fig. 3. Part of the crystal structure of (I) showing C=O(lone pair)⋯π(ring) stacking interactions as dashed lines.

## Aqua(6,6'-oxydipicolinato-κ<sup>2</sup>O,N,N',O')copper(II)

### Crystal data

[Cu(C<sub>12</sub>H<sub>6</sub>N<sub>2</sub>O<sub>5</sub>)(H<sub>2</sub>O)]

*M<sub>r</sub>* = 339.74

Monoclinic, *P*2<sub>1</sub>/*n*

Hall symbol: -*P* 2<sub>1</sub> *y* *n*

*a* = 7.2487 (16) Å

*b* = 21.055 (4) Å

*c* = 8.2269 (17) Å

β = 110.201 (9)°

*V* = 1178.4 (4) Å<sup>3</sup>

*Z* = 4

*F*(000) = 684

*D<sub>x</sub>* = 1.915 Mg m<sup>-3</sup>

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2501 reflections

θ = 2.8–27.5°

μ = 1.89 mm<sup>-1</sup>

*T* = 296 K

Block, blue

0.40 × 0.35 × 0.30 mm

### Data collection

Siemens SMART CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

φ and ω scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)

*T<sub>min</sub>* = 0.519, *T<sub>max</sub>* = 0.602

6790 measured reflections

2074 independent reflections

1806 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.027

θ<sub>max</sub> = 25.0°, θ<sub>min</sub> = 1.9°

*h* = -8→8

*k* = -24→24

*l* = -9→8

### Refinement

Refinement on *F*<sup>2</sup>

Primary atom site location: structure-invariant direct  
methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.032$$

$$wR(F^2) = 0.110$$

$$S = 1.18$$

2074 reflections

190 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0671P)^2 + 0.1209P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.48 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.37 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cu1	0.33165 (5)	0.109980 (17)	0.96148 (4)	0.02725 (18)
O3	0.2907 (3)	0.05841 (10)	1.1417 (3)	0.0327 (5)
N2	0.4163 (4)	0.14875 (12)	0.7848 (3)	0.0265 (6)
O1	0.4318 (3)	0.18774 (11)	1.0899 (3)	0.0336 (5)
O5	0.3272 (4)	0.06644 (11)	0.5777 (3)	0.0368 (6)
N1	0.2837 (4)	0.03042 (12)	0.8315 (3)	0.0254 (6)
O4	0.2158 (4)	-0.04040 (11)	1.1935 (3)	0.0409 (6)
O2	0.5275 (4)	0.28511 (11)	1.0481 (3)	0.0410 (6)
C12	0.2463 (4)	0.00062 (15)	1.1003 (4)	0.0268 (7)
C1	0.4833 (5)	0.23022 (15)	1.0031 (4)	0.0297 (7)
O6	0.0171 (3)	0.14739 (11)	0.8390 (3)	0.0382 (6)
H6A	-0.0120	0.1623	0.7373	0.046*
H6B	-0.0521	0.1144	0.8010	0.046*
C2	0.4850 (5)	0.20891 (14)	0.8261 (4)	0.0284 (7)
C5	0.4587 (5)	0.15934 (17)	0.5117 (4)	0.0342 (8)
H5	0.4473	0.1419	0.4048	0.041*
C11	0.2346 (4)	-0.01770 (15)	0.9187 (4)	0.0258 (7)
C10	0.1814 (5)	-0.07640 (16)	0.8462 (4)	0.0348 (8)
H10	0.1481	-0.1089	0.9075	0.042*
C6	0.4036 (5)	0.12555 (15)	0.6311 (4)	0.0298 (7)
C8	0.2272 (5)	-0.03744 (16)	0.5894 (4)	0.0358 (8)
H8	0.2241	-0.0430	0.4763	0.043*
C7	0.2808 (5)	0.02011 (15)	0.6721 (4)	0.0280 (7)

## supplementary materials

---

C9	0.1787 (5)	-0.08608 (16)	0.6763 (4)	0.0374 (8)
H9	0.1441	-0.1255	0.6235	0.045*
C3	0.5440 (5)	0.24521 (16)	0.7162 (4)	0.0366 (8)
H3	0.5919	0.2861	0.7470	0.044*
C4	0.5311 (5)	0.21967 (17)	0.5561 (5)	0.0407 (9)
H4	0.5715	0.2435	0.4793	0.049*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.0409 (3)	0.0226 (3)	0.0216 (3)	-0.00293 (15)	0.01511 (19)	-0.00212 (14)
O3	0.0481 (14)	0.0300 (13)	0.0229 (11)	-0.0044 (10)	0.0161 (10)	-0.0017 (9)
N2	0.0309 (14)	0.0253 (14)	0.0242 (13)	0.0005 (11)	0.0105 (10)	0.0002 (11)
O1	0.0470 (14)	0.0288 (12)	0.0267 (11)	-0.0051 (10)	0.0150 (10)	-0.0053 (9)
O5	0.0589 (16)	0.0326 (13)	0.0237 (11)	-0.0090 (11)	0.0203 (11)	-0.0052 (10)
N1	0.0327 (14)	0.0225 (13)	0.0220 (12)	-0.0009 (11)	0.0108 (10)	-0.0010 (10)
O4	0.0594 (16)	0.0381 (14)	0.0303 (12)	-0.0093 (12)	0.0219 (11)	0.0034 (11)
O2	0.0495 (15)	0.0263 (13)	0.0471 (15)	-0.0071 (10)	0.0166 (12)	-0.0104 (11)
C12	0.0283 (16)	0.0302 (18)	0.0215 (15)	0.0000 (13)	0.0082 (13)	0.0033 (13)
C1	0.0275 (16)	0.0285 (19)	0.0323 (16)	0.0031 (13)	0.0094 (13)	-0.0018 (14)
O6	0.0378 (13)	0.0339 (13)	0.0400 (13)	0.0015 (10)	0.0098 (11)	0.0019 (11)
C2	0.0286 (16)	0.0233 (16)	0.0310 (16)	-0.0001 (13)	0.0075 (13)	0.0014 (13)
C5	0.0361 (18)	0.040 (2)	0.0298 (17)	0.0017 (14)	0.0164 (14)	0.0037 (14)
C11	0.0289 (16)	0.0250 (16)	0.0234 (15)	0.0026 (12)	0.0087 (12)	0.0014 (12)
C10	0.048 (2)	0.0268 (18)	0.0324 (18)	-0.0054 (14)	0.0167 (15)	-0.0001 (14)
C6	0.0361 (18)	0.0305 (18)	0.0246 (16)	0.0025 (14)	0.0130 (13)	0.0022 (13)
C8	0.046 (2)	0.036 (2)	0.0277 (16)	-0.0008 (15)	0.0161 (15)	-0.0095 (15)
C7	0.0348 (17)	0.0279 (17)	0.0227 (15)	0.0012 (13)	0.0117 (13)	-0.0001 (13)
C9	0.050 (2)	0.0285 (18)	0.0346 (18)	-0.0074 (16)	0.0160 (16)	-0.0114 (15)
C3	0.0374 (19)	0.0310 (18)	0.042 (2)	-0.0050 (15)	0.0142 (15)	0.0034 (16)
C4	0.043 (2)	0.042 (2)	0.042 (2)	-0.0028 (16)	0.0217 (16)	0.0123 (17)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Cu1—O3	1.942 (2)	O6—H6A	0.8500
Cu1—N2	1.942 (3)	O6—H6B	0.8501
Cu1—O1	1.948 (2)	C2—C3	1.361 (5)
Cu1—N1	1.953 (2)	C5—C4	1.375 (5)
Cu1—O6	2.290 (2)	C5—C6	1.379 (4)
O3—C12	1.275 (4)	C5—H5	0.9300
N2—C6	1.328 (4)	C11—C10	1.368 (5)
N2—C2	1.361 (4)	C10—C9	1.406 (5)
O1—C1	1.278 (4)	C10—H10	0.9300
O5—C7	1.359 (4)	C8—C9	1.363 (5)
O5—C6	1.371 (4)	C8—C7	1.378 (5)
N1—C7	1.323 (4)	C8—H8	0.9300
N1—C11	1.358 (4)	C9—H9	0.9300
O4—C12	1.225 (4)	C3—C4	1.395 (5)
O2—C1	1.222 (4)	C3—H3	0.9300

C12—C11	1.517 (4)	C4—H4	0.9300
C1—C2	1.528 (4)		
O3—Cu1—N2	167.91 (10)	N2—C2—C1	112.9 (3)
O3—Cu1—O1	100.51 (9)	C3—C2—C1	125.2 (3)
N2—Cu1—O1	84.13 (10)	C4—C5—C6	117.7 (3)
O3—Cu1—N1	83.87 (9)	C4—C5—H5	121.1
N2—Cu1—N1	89.62 (10)	C6—C5—H5	121.1
O1—Cu1—N1	168.68 (10)	N1—C11—C10	122.0 (3)
O3—Cu1—O6	97.86 (10)	N1—C11—C12	113.2 (3)
N2—Cu1—O6	92.86 (10)	C10—C11—C12	124.8 (3)
O1—Cu1—O6	94.45 (9)	C11—C10—C9	118.0 (3)
N1—Cu1—O6	95.29 (10)	C11—C10—H10	121.0
C12—O3—Cu1	114.87 (19)	C9—C10—H10	121.0
C6—N2—C2	118.7 (3)	N2—C6—O5	121.8 (3)
C6—N2—Cu1	128.4 (2)	N2—C6—C5	123.1 (3)
C2—N2—Cu1	112.8 (2)	O5—C6—C5	115.1 (3)
C1—O1—Cu1	114.29 (19)	C9—C8—C7	118.8 (3)
C7—O5—C6	128.3 (2)	C9—C8—H8	120.6
C7—N1—C11	118.9 (3)	C7—C8—H8	120.6
C7—N1—Cu1	128.5 (2)	N1—C7—O5	121.8 (3)
C11—N1—Cu1	112.4 (2)	N1—C7—C8	122.6 (3)
O4—C12—O3	126.1 (3)	O5—C7—C8	115.6 (3)
O4—C12—C11	118.5 (3)	C8—C9—C10	119.7 (3)
O3—C12—C11	115.4 (3)	C8—C9—H9	120.2
O2—C1—O1	126.1 (3)	C10—C9—H9	120.2
O2—C1—C2	118.6 (3)	C2—C3—C4	118.5 (3)
O1—C1—C2	115.3 (3)	C2—C3—H3	120.8
Cu1—O6—H6A	115.6	C4—C3—H3	120.8
Cu1—O6—H6B	104.6	C5—C4—C3	120.1 (3)
H6A—O6—H6B	91.5	C5—C4—H4	119.9
N2—C2—C3	121.9 (3)	C3—C4—H4	119.9

Hydrogen-bond geometry (Å, °)

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C8—H8 $\cdots$ O4 <sup>i</sup>	0.93	2.31	3.229 (4)	171
C9—H9 $\cdots$ O2 <sup>ii</sup>	0.93	2.42	3.331 (4)	165
C4—H4 $\cdots$ O6 <sup>iii</sup>	0.93	2.54	3.303 (4)	140
O6—H6B $\cdots$ O4 <sup>iv</sup>	0.85	1.97	2.772 (3)	157
O6—H6A $\cdots$ O2 <sup>v</sup>	0.85	2.01	2.807 (3)	156

Symmetry codes: (i)  $x, y, z-1$ ; (ii)  $-x+1/2, y-1/2, -z+3/2$ ; (iii)  $x+1/2, -y+1/2, z-1/2$ ; (iv)  $-x, -y, -z+2$ ; (v)  $x-1/2, -y+1/2, z-1/2$ .

Fig. 1






Fig. 2




Fig. 3

