

Poly[[tetraaquadi- μ_3 -oxalato- μ_2 -oxalato-diprasedymium(III)] dihydrate]

Jian-Hong Chen, Hua-Cai Fang, Hong-Yang Jia, Shan-Shan Li and Yue-Peng Cai*

School of Chemistry and Environment, South China Normal University, Guangzhou 510631, People's Republic of China

Correspondence e-mail: ypcai8@yahoo.com


Received 9 November 2009; accepted 12 December 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.016; wR factor = 0.035; data-to-parameter ratio = 10.7.

In the title compound, $\{[\text{Pr}_2(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_4]\cdot 2\text{H}_2\text{O}\}_n$, the three-dimensional network structure has the Pr^{III} ion coordinated by nine O atoms in a distorted tricapped trigonal-prismatic geometry. The coordinated and uncoordinated water molecules interact with the carboxylate O atoms to consolidate the network via $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For general background, see: Benson *et al.* (2000).



Experimental

Crystal data

$[\text{Pr}_2(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_4]\cdot 2\text{H}_2\text{O}$

$M_r = 653.98$

Monoclinic, $P2_1/c$

$a = 9.8834(5)\text{ \AA}$

$b = 8.2811(4)\text{ \AA}$

$c = 10.1818(5)\text{ \AA}$

$\beta = 99.053(1)^\circ$

$V = 822.95(7)\text{ \AA}^3$

$Z = 2$

Mo $K\alpha$ radiation

$\mu = 5.95\text{ mm}^{-1}$
 $T = 298\text{ K}$

$0.26 \times 0.22 \times 0.16\text{ mm}$

Data collection

Bruker SMART diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.307$, $T_{\max} = 0.450$

4111 measured reflections
1487 independent reflections
1397 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.016$
 $wR(F^2) = 0.035$
 $S = 1.07$
1487 reflections
139 parameters
9 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.44\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.54\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O7—H7A \cdots O9 ⁱ	0.85 (1)	1.89 (1)	2.732 (3)	171 (3)
O7—H7B \cdots O1 ⁱⁱ	0.85 (1)	1.94 (2)	2.732 (3)	155 (4)
O8—H8A \cdots O2 ⁱⁱⁱ	0.84 (1)	2.06 (1)	2.900 (3)	175 (4)
O8—H8A \cdots O3 ^{iv}	0.84 (1)	2.59 (4)	3.026 (3)	114 (3)
O8—H8B \cdots O9 ^v	0.84 (1)	2.12 (2)	2.949 (4)	167 (6)
O9—H9B \cdots O6 ^{vi}	0.83 (1)	1.99 (1)	2.820 (3)	178 (4)
O9—H9A \cdots O8	0.84 (1)	2.05 (1)	2.881 (4)	173 (4)

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x, -y, -z + 1$; (iii) $-x, -y + 1, -z + 1$; (iv) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (v) $-x + 1, -y + 1, -z + 1$; (vi) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The work was supported by the National Natural Science Foundation of China (No. 20772037) and the NSF of Guangdong Province, China (grant Nos. 9251063101000006 and 06025033).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2682).

References

- Benson, D. A., Karsch-Mizrachi, I., Lipman, D. J., Ostell, J., Rapp, B. A. & Wheeler, D. L. (2000). *Genbank*. *Nucl. Acids Res.* **28**, 15–18.
- Bruker (1998). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (1999). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2010). E66, m82 [doi:10.1107/S1600536809053598]

Poly[[tetraaquadi- μ_3 -oxalato- μ_2 -oxalato-diprasedymium(III)] dihydrate]

J.-H. Chen, H.-C. Fang, H.-Y. Jia, S.-S. Li and Y.-P. Cai

Comment

Though complexes including oxalato and praseodymium(III) have been extensively investigated, the crystal structure of praseodymium oxalate with one-dimensional, two-dimensional and three-dimensional topologies and different solvent molecules have constantly been reported recently. In this paper, we would like to report the synthesis and crystal structure of a three-dimensional network complex including oxalato and praseodymium(III) with one lattice water molecule per unsymmetrical unit. The hydrothermal reaction of Pr_6O_{11} and oxalic acid in H_2O afforded the Pr^{III} title polymeric complex, $[(\text{Pr}(\text{C}_2\text{O}_4)_{1.5}(\text{H}_2\text{O})_2)\cdot\text{H}_2\text{O}]_n$. The Pr^{III} ion is coordinated by nine O atoms from two water molecules and four carboxylate groups of oxalate ligands in an irregular coordination geometry (Figure 1). Two oxalate groups of the oxalate anions bridge two symmetry-related Pr^{III} atoms, giving rise to a layer-like structure extending along [100] (Figure 2). These parallel layers are further connected via chelating coordination of oxalate anions into a three-dimensional network (Figure 3) along [010] plane. Moreover, the coordinated/non-coordinated water molecules interact with the carboxylate oxygen atoms from the layers via $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds (Table 1), which contributes to the additional stability of the structure.

Experimental

A suspension of praseodymium oxide (205 mg, 0.20 mmol) in water (20 ml) was slowly added to a solution of oxalic acid (0.10 mmol) in water (10 ml). The resultant mixture was sealed in a 50 cm³ stainless steel reactor with Teflon liner and kept under autogenous pressure at 100 ° for 78 h, and then cooled to room temperature at a rate of 0.5 °·min⁻¹. Colorless block crystals of the compound suited for single-crystal X-ray diffraction analyses formed with a yield of approximate 65%. The assigned structure was substantiated by elemental analysis; calculated for $\text{C}_3\text{H}_6\text{O}_9\text{Pr}$: C 11.01, H 1.83%; found: C 10.96, H 1.95%.

Refinement

The structure was solved using direct methods followed by Fourier synthesis. Non-H atoms were refined anisotropically. The water H atoms were located in a difference Fourier map, and were refined with distance restraints of $\text{O}-\text{H} = 0.84(0.01)$ and $\text{H}\cdots\text{H} = 1.428(0.002)$ Å, but their U_{iso} values were set equal to 1.5 U_{eq} (parent atom O).

Figures




Fig. 1. The structure of the title compound (I) with 30% probability ellipsoids. Symmetry code a: 1 - x , - y , 1 - z ; b: - x , -1/2 + y , 0.5 - z ; c: - x , - y , 1 - z ; d: x , 0.5 - y , 1/2 + z .

supplementary materials




Fig. 2. A view of one-dimensional chain constructed by hydrogen bonding interactions. Symmetry code a: $x - 1, y, z$.

Poly[[tetraaquadi- μ_3 -oxalato- μ_2 -oxalato-diprasedymium(III)] dihydrate]

Crystal data

[Pr ₂ (C ₂ O ₄) ₃ (H ₂ O) ₄]·2H ₂ O	$F(000) = 620$
$M_r = 653.98$	$D_x = 2.639 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 9.8834 (5) \text{ \AA}$	Cell parameters from 3328 reflections
$b = 8.2811 (4) \text{ \AA}$	$\theta = 3.2\text{--}28.6^\circ$
$c = 10.1818 (5) \text{ \AA}$	$\mu = 5.95 \text{ mm}^{-1}$
$\beta = 99.053 (1)^\circ$	$T = 298 \text{ K}$
$V = 822.95 (7) \text{ \AA}^3$	Block, yellow
$Z = 2$	$0.26 \times 0.22 \times 0.16 \text{ mm}$

Data collection

Bruker SMART diffractometer	1487 independent reflections
Radiation source: fine-focus sealed tube graphite	1397 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.023$
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 25.3^\circ, \theta_{\text{min}} = 2.1^\circ$
$T_{\text{min}} = 0.307, T_{\text{max}} = 0.450$	$h = -11 \rightarrow 11$
4111 measured reflections	$k = -9 \rightarrow 9$
	$l = -12 \rightarrow 7$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.016$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.035$	$w = 1/[\sigma^2(F_o^2) + (0.0087P)^2 + 0.7364P]$
$S = 1.07$	where $P = (F_o^2 + 2F_c^2)/3$
1487 reflections	$(\Delta/\sigma)_{\text{max}} = 0.007$
139 parameters	$\Delta\rho_{\text{max}} = 0.44 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.54 \text{ e \AA}^{-3}$

9 restraints Extinction correction: *SHELXL97* (Sheldrick, 2008),
 $F_c^* = k F_c [1 + 0.001 x F_c^2 \lambda^3 / \sin(2\theta)]^{1/4}$

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.0127 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes)

are estimated using the full covariance matrix. The cell e.s.d.'s are taken

into account individually in the estimation of e.s.d.'s in distances, angles

and torsion angles; correlations between e.s.d.'s in cell parameters are only

used when they are defined by crystal symmetry. An approximate (isotropic)

treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted *R*-factor wR and

goodness of fit S are based on F^2 , conventional *R*-factors R are based

on F , with F set to zero for negative F^2 . The threshold expression of

$F^2 > \sigma(F^2)$ is used only for calculating *R*-factors(gt) etc. and is

not relevant to the choice of reflections for refinement. *R*-factors based

on F^2 are statistically about twice as large as those based on F , and *R*-

factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Pr1	0.185002 (15)	0.098389 (18)	0.437068 (15)	0.01483 (9)
O1	0.0042 (2)	0.3167 (2)	0.4302 (2)	0.0210 (5)
O2	-0.1229 (2)	0.4945 (3)	0.2991 (2)	0.0249 (5)
O3	0.1838 (2)	0.3092 (3)	0.2610 (2)	0.0229 (5)
O4	0.0564 (2)	0.4802 (3)	0.12614 (19)	0.0196 (5)
O5	0.4011 (2)	0.0665 (3)	0.3529 (2)	0.0275 (5)
O6	0.6199 (2)	-0.0090 (3)	0.3924 (2)	0.0287 (5)
O7	0.1928 (2)	-0.2047 (3)	0.4324 (3)	0.0357 (6)
H7A	0.255 (3)	-0.260 (3)	0.406 (4)	0.054*
H7B	0.153 (3)	-0.250 (4)	0.490 (3)	0.054*
O8	0.3072 (3)	0.3550 (3)	0.5421 (2)	0.0296 (5)
H8A	0.252 (3)	0.393 (4)	0.589 (3)	0.044*
O9	0.3963 (3)	0.5983 (3)	0.3724 (3)	0.0353 (6)
H9B	0.393 (4)	0.566 (5)	0.2945 (16)	0.053*
C1	-0.0222 (3)	0.4050 (3)	0.3291 (3)	0.0172 (6)
C2	0.0829 (3)	0.3968 (3)	0.2319 (3)	0.0162 (6)

supplementary materials

C3	0.5054 (3)	0.0165 (4)	0.4261 (3)	0.0200 (7)
H8B	0.3907 (13)	0.358 (8)	0.576 (4)	0.13 (3)*
H9A	0.370 (7)	0.534 (5)	0.427 (3)	0.13 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Pr1	0.01116 (11)	0.01772 (12)	0.01591 (11)	0.00145 (6)	0.00305 (7)	0.00222 (6)
O1	0.0236 (11)	0.0229 (11)	0.0185 (11)	0.0044 (9)	0.0094 (9)	0.0063 (9)
O2	0.0248 (12)	0.0312 (12)	0.0191 (11)	0.0126 (10)	0.0051 (10)	0.0018 (9)
O3	0.0186 (11)	0.0244 (11)	0.0274 (12)	0.0049 (9)	0.0090 (10)	0.0067 (10)
O4	0.0179 (11)	0.0261 (11)	0.0149 (11)	0.0013 (9)	0.0033 (9)	0.0049 (9)
O5	0.0183 (12)	0.0451 (14)	0.0198 (12)	0.0077 (10)	0.0052 (10)	0.0090 (10)
O6	0.0195 (12)	0.0478 (14)	0.0204 (12)	0.0121 (10)	0.0078 (10)	0.0095 (11)
O7	0.0351 (14)	0.0252 (13)	0.0533 (17)	0.0080 (11)	0.0273 (13)	0.0110 (11)
O8	0.0303 (14)	0.0317 (13)	0.0272 (13)	-0.0058 (11)	0.0057 (11)	-0.0034 (10)
O9	0.0397 (15)	0.0410 (15)	0.0274 (14)	0.0031 (12)	0.0118 (13)	-0.0034 (12)
C1	0.0182 (16)	0.0162 (14)	0.0170 (15)	-0.0001 (12)	0.0023 (13)	-0.0043 (12)
C2	0.0188 (16)	0.0156 (14)	0.0141 (15)	-0.0066 (12)	0.0022 (12)	-0.0034 (12)
C3	0.0170 (16)	0.0257 (16)	0.0183 (16)	0.0030 (13)	0.0064 (13)	0.0042 (13)

Geometric parameters (\AA , $^\circ$)

Pr1—O5	2.438 (2)	O4—Pr1 ^v	2.552 (2)
Pr1—O6 ⁱ	2.495 (2)	O4—Pr1 ^{iv}	2.566 (2)
Pr1—O3	2.501 (2)	O5—C3	1.244 (4)
Pr1—O7	2.512 (2)	O6—C3	1.251 (4)
Pr1—O1	2.535 (2)	O6—Pr1 ⁱ	2.495 (2)
Pr1—O2 ⁱⁱ	2.537 (2)	O7—H7A	0.847 (10)
Pr1—O4 ⁱⁱⁱ	2.552 (2)	O7—H7B	0.845 (10)
Pr1—O4 ⁱⁱ	2.566 (2)	O8—H8A	0.842 (10)
Pr1—O8	2.590 (2)	O8—H8B	0.844 (10)
O1—C1	1.257 (4)	O9—H9B	0.832 (10)
O2—C1	1.240 (4)	O9—H9A	0.839 (10)
O2—Pr1 ^{iv}	2.537 (2)	C1—C2	1.545 (4)
O3—C2	1.231 (4)	C3—C3 ⁱ	1.549 (6)
O4—C2	1.271 (4)		
O5—Pr1—O6 ⁱ	65.72 (7)	O3—Pr1—O8	70.89 (8)
O5—Pr1—O3	74.28 (7)	O7—Pr1—O8	144.46 (8)
O6 ⁱ —Pr1—O3	128.45 (7)	O1—Pr1—O8	72.97 (8)
O5—Pr1—O7	81.61 (7)	O2 ⁱⁱ —Pr1—O8	134.18 (7)
O6 ⁱ —Pr1—O7	72.19 (8)	O4 ⁱⁱⁱ —Pr1—O8	98.79 (7)
O3—Pr1—O7	132.90 (8)	O4 ⁱⁱ —Pr1—O8	140.47 (7)
O5—Pr1—O1	136.09 (7)	C1—O1—Pr1	119.53 (18)
O6 ⁱ —Pr1—O1	134.01 (7)	C1—O2—Pr1 ^{iv}	120.07 (19)
O3—Pr1—O1	63.82 (6)	C2—O3—Pr1	119.75 (18)

O7—Pr1—O1	137.35 (7)	C2—O4—Pr1 ^v	116.09 (18)
O5—Pr1—O2 ⁱⁱ	74.00 (7)	C2—O4—Pr1 ^{iv}	118.63 (18)
O6 ⁱ —Pr1—O2 ⁱⁱ	127.17 (7)	Pr1 ^v —O4—Pr1 ^{iv}	117.51 (8)
O3—Pr1—O2 ⁱⁱ	65.42 (7)	C3—O5—Pr1	120.94 (18)
O7—Pr1—O2 ⁱⁱ	69.33 (8)	C3—O6—Pr1 ⁱ	119.5 (2)
O1—Pr1—O2 ⁱⁱ	98.71 (7)	Pr1—O7—H7A	125 (2)
O5—Pr1—O4 ⁱⁱⁱ	143.64 (7)	Pr1—O7—H7B	114 (2)
O6 ⁱ —Pr1—O4 ⁱⁱⁱ	79.71 (7)	H7A—O7—H7B	115.0 (15)
O3—Pr1—O4 ⁱⁱⁱ	140.53 (7)	Pr1—O8—H8A	104 (2)
O7—Pr1—O4 ⁱⁱⁱ	77.27 (7)	Pr1—O8—H8B	124 (4)
O1—Pr1—O4 ⁱⁱⁱ	76.71 (6)	H8A—O8—H8B	115.7 (15)
O2 ⁱⁱ —Pr1—O4 ⁱⁱⁱ	123.63 (6)	H9B—O9—H9A	117.4 (16)
O5—Pr1—O4 ⁱⁱ	134.60 (7)	O2—C1—O1	127.3 (3)
O6 ⁱ —Pr1—O4 ⁱⁱ	130.37 (7)	O2—C1—C2	117.6 (3)
O3—Pr1—O4 ⁱⁱ	100.79 (7)	O1—C1—C2	115.1 (3)
O7—Pr1—O4 ⁱⁱ	69.15 (7)	O3—C2—O4	125.1 (3)
O1—Pr1—O4 ⁱⁱ	68.99 (7)	O3—C2—C1	118.2 (3)
O2 ⁱⁱ —Pr1—O4 ⁱⁱ	63.66 (6)	O4—C2—C1	116.7 (3)
O4 ⁱⁱⁱ —Pr1—O4 ⁱⁱ	62.49 (8)	O5—C3—O6	126.3 (3)
O5—Pr1—O8	81.77 (8)	O5—C3—C3 ⁱ	117.6 (3)
O6 ⁱ —Pr1—O8	72.37 (8)	O6—C3—C3 ⁱ	116.1 (3)

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, y-1/2, -z+1/2$; (iii) $x, -y+1/2, z+1/2$; (iv) $-x, y+1/2, -z+1/2$; (v) $x, -y+1/2, z-1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D—\text{H}\cdots A$	$D—\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D—\text{H}\cdots A$
O7—H7A \cdots O9 ^{vi}	0.85 (1)	1.89 (1)	2.732 (3)	171 (3)
O7—H7B \cdots O1 ^{vii}	0.85 (1)	1.94 (2)	2.732 (3)	155 (4)
O8—H8A \cdots O2 ^{viii}	0.84 (1)	2.06 (1)	2.900 (3)	175 (4)
O8—H8A \cdots O3 ⁱⁱⁱ	0.84 (1)	2.59 (4)	3.026 (3)	114 (3)
O8—H8B \cdots O9 ^{ix}	0.84 (1)	2.12 (2)	2.949 (4)	167 (6)
O9—H9B \cdots O6 ^x	0.83 (1)	1.99 (1)	2.820 (3)	178 (4)
O9—H9A \cdots O8	0.84 (1)	2.05 (1)	2.881 (4)	173 (4)

Symmetry codes: (vi) $x, y-1, z$; (vii) $-x, -y, -z+1$; (viii) $-x, -y+1, -z+1$; (iii) $x, -y+1/2, z+1/2$; (ix) $-x+1, -y+1, -z+1$; (x) $-x+1, y+1/2, -z+1/2$.

supplementary materials

Fig. 1




Fig. 2

