

Quinoxaline-2-carbonitrile

Hoong-Kun Fun,^{a*}‡ Ching Kheng Quah,^{a§} Annada C. Maity,^b Nirmal Kumar Das^b and Shyamaprosad Goswami^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, Bengal Engineering and Science University, Shibpur, Howrah 711 103, India

Correspondence e-mail: hkfun@usm.my


Received 26 November 2009; accepted 28 November 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.001\text{ \AA}$; R factor = 0.047; wR factor = 0.135; data-to-parameter ratio = 21.1.

In the title compound, $\text{C}_9\text{H}_5\text{N}_3$, the quinoxaline ring is essentially planar, with a maximum deviation of $0.012(1)\text{ \AA}$. Short intermolecular distances between the centroids of the 2,3-dihydropyrazine and benzene rings [$3.6490(5)\text{ \AA}$] indicate the existence of $\pi \cdots \pi$ interactions. In the crystal packing, the molecules are linked *via* two pairs of intermolecular $\text{C}-\text{H} \cdots \text{N}$ interactions, forming $R_2^2(8)$ and $R_2^2(10)$ ring motifs; these molecules are further linked into a two-dimensional network parallel to (1 0 2) *via* another $\text{C}-\text{H} \cdots \text{N}$ interaction.

Related literature

For the synthesis of cyano *N*-heterocyclic compounds, see: Goswami *et al.* (2007, 2009). For reference bond lengths, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_9\text{H}_5\text{N}_3$
 $M_r = 155.16$
Monoclinic, $P2_1/c$

$a = 3.8055(1)\text{ \AA}$
 $b = 19.0466(4)\text{ \AA}$
 $c = 10.1845(2)\text{ \AA}$

$\beta = 93.466(1)^\circ$
 $V = 736.84(3)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.09\text{ mm}^{-1}$
 $T = 100\text{ K}$
 $0.39 \times 0.28 \times 0.25\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.966$, $T_{\max} = 0.978$

11604 measured reflections
2716 independent reflections
2183 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.135$
 $S = 1.08$
2716 reflections

129 parameters
All H-atom parameters refined
 $\Delta\rho_{\max} = 0.53\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.23\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{C}2-\text{H} \cdots \text{N}1^i$	0.984 (14)	2.619 (14)	3.5730 (12)	163.4 (12)
$\text{C}4-\text{H}4 \cdots \text{N}2^{ii}$	0.988 (13)	2.593 (13)	3.4268 (12)	142.0 (10)
$\text{C}7-\text{H}7 \cdots \text{N}3^{iii}$	0.998 (14)	2.540 (15)	3.5225 (12)	168.3 (12)

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$; (iii) $-x - 1, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF and CKQ thank Universiti Sains Malaysia (USM) for the Research University Golden Goose Grant (1001/PFIZIK/811012). CKQ thanks USM for a Research Fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2699).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, S1–19.
Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
Goswami, S. P., Maity, A. C., Das, N. K., Sen, D. & Maity, S. P. (2009). *Synth. Commun.* **39**, 407–415.
Goswami, S., Maity, A. C., García-Granda, S. & Torre-Fernández, L. (2007). *Acta Cryst. E63*, o1741–o1742.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

‡ Thomson Reuters ResearcherID: A-3561-2009.
§ Thomson Reuters ResearcherID: A-5525-2009.

supplementary materials

Acta Cryst. (2010). E66, o28 [doi:10.1107/S1600536809051289]

Quinoxaline-2-carbonitrile

H.-K. Fun, C. K. Quah, A. C. Maity, N. K. Das and S. Goswami

Comment

Heterocyclic molecules containing a cyano group are useful as drug intermediates. The development of new pathways leading to efficient synthesis of heterocycles with diversity in skeleton and functional groups is an important field of research in both synthetic and medicinal chemistry. Recently, we have synthesized a number of cyano N-heterocyclic compounds using triselenium dicyanide (TSD) (Goswami *et al.*, 2007, 2009). Herein we report the synthesis of 2-cyanoquinoxaline from quinoxaline under microwave irradiation and its molecular structure.

The bond lengths (Allen *et al.*, 1987) and angles in the title compound (Fig. 1) are within normal ranges. The quinoxaline ring (N1/N2/C1-C8) is essentially planar, with the maximum deviation of 0.012 (1) Å for atom C5. Short intermolecular distances between the centroids of the pyrazine (N1/N2/C1/C6-C8) and benzene rings (C1-C6) [3.6490 (5) Å] indicate the existence of $\pi\cdots\pi$ interactions.

In the crystal packing (Fig. 2), the molecules are linked *via* pairs of intermolecular C2—H2 \cdots N1 and C7—H7 \cdots N3 interactions, forming R^2_2 (8) and R^2_2 (10) ring motifs (Bernstein *et al.*, 1995) and these molecules are further linked into two-dimensional networks parallel to plane (1 0 2) *via* C4—H4 \cdots N2 interactions.

Experimental

A thoroughly ground mixture of selenium dioxide (1.32 g, 12 mmol) and malononitrile (0.26 g, 4 mmol) in 4-5 drops of DMSO was kept stirring in an open-mouth conical flask. The mixture became reddish-brown after 7 min. An exothermic reaction began in the next 10 minutes when triselenium dicyanide was formed. The heterocyclic substrate quinoxaline (0.39 g, 3 mmol) was added to the mixture after the termination of the exothermic reaction. The conical flask was placed in a domestic microwave oven at 240 W for 20 min. The progress of the reaction was monitored by TLC. After completion of the reaction, water was added and the mixture was extracted with chloroform. The organic layer was washed with saturated brine and dried over MgSO₄ and followed by evaporation with a rotary evaporator under low pressure to afford a light yellow substance. This was purified on silica gel (60-120 mesh) column chromatography eluting with petroleum ether (boiling point, 60-80° C) to give the compound (0.34 g, 74 %) as a crystalline solid.

Refinement

All H atoms were located in a difference Fourier map and refined freely.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound, showing 50% probability displacement ellipsoids for non-H atoms and the atom-numbering scheme.




Fig. 2. The crystal structure of the title compound viewed along the a axis. Intermolecular interactions are shown in dashed lines.

Quinoxaline-2-carbonitrile

Crystal data

$C_9H_5N_3$	$F(000) = 320$
$M_r = 155.16$	$D_x = 1.399 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 4710 reflections
$a = 3.8055 (1) \text{ \AA}$	$\theta = 2.9\text{--}32.7^\circ$
$b = 19.0466 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$c = 10.1845 (2) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 93.466 (1)^\circ$	Block, yellow
$V = 736.84 (3) \text{ \AA}^3$	$0.39 \times 0.28 \times 0.25 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	2716 independent reflections
Radiation source: fine-focus sealed tube	2183 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.023$
φ and ω scans	$\theta_{\text{max}} = 32.8^\circ, \theta_{\text{min}} = 2.1^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -5 \rightarrow 5$
$T_{\text{min}} = 0.966, T_{\text{max}} = 0.978$	$k = -29 \rightarrow 21$
11604 measured reflections	$l = -15 \rightarrow 14$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.047$	Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.135$	All H-atom parameters refined
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0782P)^2 + 0.0918P]$
2716 reflections	where $P = (F_o^2 + 2F_c^2)/3$
129 parameters	$(\Delta/\sigma)_{\max} = 0.001$
0 restraints	$\Delta\rho_{\max} = 0.53 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\min} = -0.23 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.12156 (18)	0.46098 (4)	0.84114 (7)	0.01481 (16)
N2	-0.12366 (19)	0.33470 (4)	0.71846 (8)	0.01659 (17)
N3	-0.2187 (2)	0.57757 (5)	0.60971 (9)	0.0269 (2)
C1	0.1884 (2)	0.39728 (4)	0.89844 (8)	0.01344 (17)
C2	0.3822 (2)	0.39456 (5)	1.02181 (9)	0.01744 (18)
C3	0.4514 (2)	0.33059 (5)	1.07937 (10)	0.02033 (19)
C4	0.3352 (2)	0.26756 (5)	1.01678 (10)	0.0210 (2)
C5	0.1474 (2)	0.26871 (5)	0.89805 (9)	0.01857 (19)
C6	0.0674 (2)	0.33382 (4)	0.83638 (9)	0.01441 (17)
C7	-0.1868 (2)	0.39663 (4)	0.66474 (9)	0.01675 (18)
C8	-0.0633 (2)	0.45941 (4)	0.72702 (8)	0.01496 (17)
C9	-0.1450 (2)	0.52598 (5)	0.66332 (9)	0.01913 (19)
H2	0.471 (4)	0.4375 (7)	1.0662 (14)	0.030 (3)*
H3	0.591 (4)	0.3282 (7)	1.1650 (16)	0.038 (4)*
H4	0.396 (3)	0.2229 (7)	1.0622 (13)	0.028 (3)*
H5	0.057 (4)	0.2262 (7)	0.8533 (14)	0.032 (3)*
H7	-0.328 (4)	0.4008 (7)	0.5793 (14)	0.028 (3)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0154 (3)	0.0145 (3)	0.0146 (3)	0.0003 (2)	0.0014 (2)	0.0010 (2)
N2	0.0168 (3)	0.0173 (3)	0.0157 (4)	-0.0006 (2)	0.0014 (3)	-0.0016 (3)

supplementary materials

N3	0.0328 (4)	0.0235 (4)	0.0241 (5)	0.0028 (3)	-0.0002 (3)	0.0057 (3)
C1	0.0128 (3)	0.0147 (3)	0.0130 (4)	0.0002 (2)	0.0019 (3)	0.0004 (3)
C2	0.0157 (4)	0.0218 (4)	0.0147 (4)	-0.0004 (3)	-0.0002 (3)	0.0005 (3)
C3	0.0164 (4)	0.0275 (4)	0.0170 (4)	0.0019 (3)	0.0003 (3)	0.0055 (3)
C4	0.0176 (4)	0.0208 (4)	0.0248 (5)	0.0030 (3)	0.0042 (3)	0.0088 (3)
C5	0.0179 (4)	0.0147 (4)	0.0235 (5)	0.0007 (3)	0.0036 (3)	0.0029 (3)
C6	0.0131 (3)	0.0152 (3)	0.0151 (4)	0.0000 (2)	0.0027 (3)	0.0000 (3)
C7	0.0164 (4)	0.0195 (4)	0.0142 (4)	-0.0003 (3)	0.0000 (3)	-0.0004 (3)
C8	0.0145 (3)	0.0168 (4)	0.0137 (4)	0.0005 (3)	0.0019 (3)	0.0015 (3)
C9	0.0204 (4)	0.0202 (4)	0.0167 (4)	0.0005 (3)	0.0005 (3)	0.0015 (3)

Geometric parameters (\AA , $^\circ$)

N1—C8	1.3220 (11)	C3—C4	1.4175 (14)
N1—C1	1.3636 (10)	C3—H3	0.995 (16)
N2—C7	1.3161 (11)	C4—C5	1.3670 (13)
N2—C6	1.3659 (11)	C4—H4	0.988 (13)
N3—C9	1.1506 (12)	C5—C6	1.4150 (11)
C1—C2	1.4190 (12)	C5—H5	0.980 (14)
C1—C6	1.4273 (11)	C7—C8	1.4202 (12)
C2—C3	1.3706 (12)	C7—H7	0.997 (14)
C2—H2	0.985 (14)	C8—C9	1.4495 (12)
C8—N1—C1	115.57 (7)	C4—C5—C6	119.57 (8)
C7—N2—C6	116.78 (7)	C4—C5—H5	123.2 (8)
N1—C1—C2	119.01 (7)	C6—C5—H5	117.2 (8)
N1—C1—C6	121.14 (8)	N2—C6—C5	119.33 (7)
C2—C1—C6	119.85 (7)	N2—C6—C1	121.28 (7)
C3—C2—C1	119.16 (8)	C5—C6—C1	119.38 (8)
C3—C2—H2	119.4 (8)	N2—C7—C8	121.44 (8)
C1—C2—H2	121.5 (8)	N2—C7—H7	120.7 (7)
C2—C3—C4	120.92 (9)	C8—C7—H7	117.9 (7)
C2—C3—H3	119.6 (8)	N1—C8—C7	123.77 (7)
C4—C3—H3	119.4 (8)	N1—C8—C9	117.52 (7)
C5—C4—C3	121.10 (8)	C7—C8—C9	118.71 (8)
C5—C4—H4	121.5 (8)	N3—C9—C8	177.49 (10)
C3—C4—H4	117.4 (8)		
C8—N1—C1—C2	-179.34 (7)	C4—C5—C6—C1	0.84 (13)
C8—N1—C1—C6	0.75 (12)	N1—C1—C6—N2	-0.95 (13)
N1—C1—C2—C3	-179.68 (8)	C2—C1—C6—N2	179.14 (7)
C6—C1—C2—C3	0.24 (13)	N1—C1—C6—C5	178.94 (7)
C1—C2—C3—C4	0.62 (13)	C2—C1—C6—C5	-0.97 (12)
C2—C3—C4—C5	-0.75 (14)	C6—N2—C7—C8	-0.32 (13)
C3—C4—C5—C6	0.00 (14)	C1—N1—C8—C7	-0.38 (12)
C7—N2—C6—C5	-179.20 (7)	C1—N1—C8—C9	179.15 (7)
C7—N2—C6—C1	0.69 (12)	N2—C7—C8—N1	0.18 (14)
C4—C5—C6—N2	-179.26 (7)	N2—C7—C8—C9	-179.36 (8)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
C2—H2 \cdots N1 ⁱ	0.984 (14)	2.619 (14)	3.5730 (12)	163.4 (12)
C4—H4 \cdots N2 ⁱⁱ	0.988 (13)	2.593 (13)	3.4268 (12)	142.0 (10)
C7—H7 \cdots N3 ⁱⁱⁱ	0.998 (14)	2.540 (15)	3.5225 (12)	168.3 (12)

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x+1, -y+1/2, z+1/2$; (iii) $-x-1, -y+1, -z+1$.

supplementary materials

Fig. 1




Fig. 2

